

Table S1 Main components in ERAD

Classification	Factor	Function	References
lectin	OS9/XTP3-B	Recognition for unfolded or misfolded protein	[1-3]
Chaperone	HSP/Bip	Recognition for unfolded or misfolded protein	[4-6]
Chaperone	EDEM _s	Recognition for unfolded or misfolded protein	[7-9]
Adaptor	SEL1	Hrd1 complex member	[3]
Adaptor	Herp	Hrd1 complex member	[10]
Dislocons	Derlin1,Derlin2, Derlin3	Hrd1 complex member	[11, 12]
ATP hydrolase	p97	Retrotranslocation	[13, 14]
p97 cofactors	Ufd1	Retrotranslocation	[15-17]
p97 cofactors	Npl4	Retrotranslocation	[15-17]
p97 cofactors	UBXD2	Retrotranslocation	[15-21]
p97 cofactors	UBXD8	Retrotranslocation	[15-21]
E1 Ubiquitin-activating enzyme	UBA1	Activate ubiquitin molecules	[22, 23]
E2 Ubiquitin-conjugating enzyme	UBC6e/Ube2j1, Ube2j2	Bound ubiquitin on E2 enzyme	[24]
E2 Ubiquitin-conjugating enzyme	UbcH5a/UBE2D1, UbcH5b /UBE2D2, UbcH5c /UBE2D3	Bound ubiquitin on E2 enzyme	[25-27]
E2 Ubiquitin-conjugating enzyme	UBE2K/E2-25K	Bound ubiquitin on E2 enzyme	[28]
E2 Ubiquitin-conjugating enzyme	UBE2G2	Bound ubiquitin on E2 enzyme	[29, 30]
E3 Ubiquitin ligase enzyme	Hrd1	Add ubiquitin chains to substrate proteins	[31]
E3 Ubiquitin ligase enzyme	TEB4/MARCH6	Add ubiquitin chains to substrate proteins	[32]
E3 Ubiquitin ligase enzyme	gp78	Add ubiquitin chains to substrate proteins	[33]
E3 Ubiquitin ligase enzyme	CHIP	Add ubiquitin chains to substrate proteins	[34, 35]
E3 Ubiquitin ligase enzyme	RMA1/RNF5	Add ubiquitin chains to substrate proteins	[36-38]
E3 Ubiquitin ligase enzyme	RNF4	Add ubiquitin chains to substrate proteins	[39]
E3 Ubiquitin ligase enzyme	RNF145	Add ubiquitin chains to substrate proteins	[40]

E3 Ubiquitin ligase enzyme	RNF170	Add ubiquitin chains to substrate proteins	[41-43]
E3 Ubiquitin ligase enzyme	RNF103/ KF-1	Add ubiquitin chains to substrate proteins	[44]
E3 Ubiquitin ligase enzyme	RNF185	Add ubiquitin chains to substrate proteins	[45]
E3 Ubiquitin ligase enzyme	Trc8	Add ubiquitin chains to substrate proteins	[46, 47]
E3 Ubiquitin ligase enzyme	RFP2	Add ubiquitin chains to substrate proteins	[48, 49]
E3 Ubiquitin ligase enzyme	SCF ^{Fbx2/Fbs1}	Add ubiquitin chains to substrate proteins	[50-52]
E3 Ubiquitin ligase enzyme	SCF ^{Fbx6/Fbs2}	Add ubiquitin chains to substrate proteins	[53, 54]
E3 Ubiquitin ligase enzyme	SCF ^{β-TrCP}	Add ubiquitin chains to substrate proteins	[55, 56]
E3 Ubiquitin ligase enzyme	Parkin	Add ubiquitin chains to substrate proteins	[57]
E3 Ubiquitin ligase enzyme	NRDP1/RNF41	Add ubiquitin chains to substrate proteins	[58]
E3 Ubiquitin ligase enzyme	NIXIN/ ZNRF4	Add ubiquitin chains to substrate proteins	[59]
E3 Ubiquitin ligase enzyme	TMEM129	Add ubiquitin chains to substrate proteins	[60-62]
E3 Ubiquitin ligase enzyme	SMURF1	Add ubiquitin chains to substrate proteins	[63]
E3 Ubiquitin ligase enzyme	RFP2/TRIM13	Add ubiquitin chains to substrate proteins	[64, 65]
E4 Ubiquitin extension enzyme	UBE4A/B	Extend the ubiquitin chain	[66]
Deubiquitinase	USP13	cleave the ubiquitin chain	[67]
Deubiquitinase	USP14	cleave the ubiquitin chain	[68]
Deubiquitinase	USP19	cleave the ubiquitin chain	[69]
Deubiquitinase	USP25	cleave the ubiquitin chain	[70]
Deubiquitinase	RPN11	cleave the ubiquitin chain	[68]
Deubiquitinase	UCH37	cleave the ubiquitin chain	[68]
Deubiquitinase	OTUB1	cleave the ubiquitin chain	[71]
Deubiquitinase	OTUD2 /YOD1	cleave the ubiquitin chain	[72]
Deubiquitinase	Ataxin-3	cleave the ubiquitin chain	[73, 74]
Ubiquitin-binding proteins	HR23a, HR23b	transfer the ubiquitinated substrate to proteasome	[75]
Ubiquitin-binding proteins	Rpn10	transfer the ubiquitinated substrate to proteasome	[76]

peptide: N-glycanase	NGLY1	Deglycosylation	[53]
----------------------	-------	-----------------	------

1. Cruciat, C.M., C. Hassler, and C. Niehrs, *The MRH protein Erlectin is a member of the endoplasmic reticulum synexpression group and functions in N-glycan recognition*. J Biol Chem, 2006. **281**(18): p. 12986-93.
2. Shenkman, M., et al., *A shared endoplasmic reticulum-associated degradation pathway involving the EDEM1 protein for glycosylated and nonglycosylated proteins*. J Biol Chem, 2013. **288**(4): p. 2167-78.
3. Christianson, J.C., et al., *OS-9 and GRP94 deliver mutant alpha1-antitrypsin to the Hrd1-SEL1L ubiquitin ligase complex for ERAD*. Nat Cell Biol, 2008. **10**(3): p. 272-82.
4. Kityk, R., J. Kopp, and M.P. Mayer, *Molecular Mechanism of J-Domain-Triggered ATP Hydrolysis by Hsp70 Chaperones*. Mol Cell, 2018. **69**(2): p. 227-237.e4.
5. Flynn, G.C., et al., *Peptide-binding specificity of the molecular chaperone BiP*. Nature, 1991. **353**(6346): p. 726-30.
6. Feige, M.J. and L.M. Hendershot, *Quality control of integral membrane proteins by assembly-dependent membrane integration*. Mol Cell, 2013. **51**(3): p. 297-309.
7. Cormier, J.H., et al., *EDEM1 recognition and delivery of misfolded proteins to the SEL1L-containing ERAD complex*. Molecular cell, 2009. **34**(5): p. 627-633.
8. Shenkman, M., et al., *Mannosidase activity of EDEM1 and EDEM2 depends on an unfolded state of their glycoprotein substrates*. Communications biology, 2018. **1**: p. 172-172.
9. Marin, M.B., et al., *Tyrosinase degradation is prevented when EDEM1 lacks the intrinsically disordered region*. PloS one, 2012. **7**(8): p. e42998-e42998.
10. Okuda-Shimizu, Y. and L.M. Hendershot, *Characterization of an ERAD pathway for nonglycosylated BiP substrates, which require Herp*. Mol Cell, 2007. **28**(4): p. 544-54.
11. Wu, X., et al., *Structural basis of ER-associated protein degradation mediated by the Hrd1 ubiquitin ligase complex*. Science, 2020. **368**(6489).
12. Neal, S., et al., *HRD Complex Self-Remodeling Enables a Novel Route of Membrane Protein Retrotranslocation*. iScience, 2020. **23**(9): p. 101493.
13. Vasic, V., et al., *Hrd1 forms the retrotranslocation pore regulated by auto-ubiquitination and binding of misfolded proteins*. Nature Cell Biology, 2020. **22**(3): p. 274-281.
14. Helenius, A. and M. Aebi, *Intracellular functions of N-linked glycans*. Science, 2001. **291**(5512): p. 2364-9.
15. Ye, Y., H.H. Meyer, and T.A. Rapoport, *The AAA ATPase Cdc48/p97 and its partners transport proteins from the ER into the cytosol*. Nature, 2001. **414**(6864): p. 652-6.
16. Alzayady, K.J., et al., *Involvement of the p97-Ufd1-Npl4 complex in the regulated endoplasmic reticulum-associated degradation of inositol 1,4,5-trisphosphate receptors*. J Biol Chem, 2005. **280**(41): p. 34530-7.
17. Tsai, B., Y. Ye, and T.A. Rapoport, *Retro-translocation of proteins from the endoplasmic reticulum into the cytosol*. Nat Rev Mol Cell Biol, 2002. **3**(4): p. 246-55.
18. Liang, J., et al., *Characterization of erasin (UBXD2): a new ER protein that promotes ER-associated protein degradation*. J Cell Sci, 2006. **119**(Pt 19): p. 4011-24.
19. Suzuki, M., et al., *Derlin-1 and UBXD8 are engaged in dislocation and degradation of lipidated ApoB-100 at lipid droplets*. Mol Biol Cell, 2012. **23**(5): p. 800-10.
20. Schuberth, C. and A. Buchberger, *UBX domain proteins: major regulators of the AAA ATPase Cdc48/p97*. Cell Mol Life Sci, 2008. **65**(15): p. 2360-71.

21. Bartz, R., et al., *Dynamic activity of lipid droplets: protein phosphorylation and GTP-mediated protein translocation*. J Proteome Res, 2007. **6**(8): p. 3256-65.
22. Borgo, C., et al., *Targeting the E1 ubiquitin-activating enzyme (UBA1) improves elexacaftor/tezacaftor/ivacaftor efficacy towards F508del and rare misfolded CFTR mutants*. Cell Mol Life Sci, 2022. **79**(4): p. 192.
23. Barghout, S.H. and A.D. Schimmer, *E1 Enzymes as Therapeutic Targets in Cancer*. Pharmacol Rev, 2021. **73**(1): p. 1-58.
24. Wang, X., et al., *Ube2j2 ubiquitinates hydroxylated amino acids on ER-associated degradation substrates*. The Journal of cell biology, 2009. **187**(5): p. 655-668.
25. Younger, J.M., et al., *A foldable CFTR{Delta}F508 biogenic intermediate accumulates upon inhibition of the Hsc70-CHIP E3 ubiquitin ligase*. The Journal of cell biology, 2004. **167**(6): p. 1075-1085.
26. Nadav, E., et al., *A novel mammalian endoplasmic reticulum ubiquitin ligase homologous to the yeast Hrd1*. Biochem Biophys Res Commun, 2003. **303**(1): p. 91-7.
27. Kaneko, M., et al., *Loss of HRD1-mediated protein degradation causes amyloid precursor protein accumulation and amyloid-beta generation*. The Journal of neuroscience : the official journal of the Society for Neuroscience, 2010. **30**(11): p. 3924-3932.
28. Flierman, D., et al., *E2-25K mediates US11-triggered retro-translocation of MHC class I heavy chains in a permeabilized cell system*. Proceedings of the National Academy of Sciences of the United States of America, 2006. **103**(31): p. 11589-11594.
29. Christianson, J.C. and Y. Ye, *Cleaning up in the endoplasmic reticulum: ubiquitin in charge*. Nat Struct Mol Biol, 2014. **21**(4): p. 325-35.
30. Chen, B., et al., *The activity of a human endoplasmic reticulum-associated degradation E3, gp78, requires its Cue domain, RING finger, and an E2-binding site*. Proc Natl Acad Sci U S A, 2006. **103**(2): p. 341-6.
31. Hampton, R.Y., R.G. Gardner, and J. Rine, *Role of 26S proteasome and HRD genes in the degradation of 3-hydroxy-3-methylglutaryl-CoA reductase, an integral endoplasmic reticulum membrane protein*. Mol Biol Cell, 1996. **7**(12): p. 2029-44.
32. Hassink, G., et al., *TEB4 is a C4HC3 RING finger-containing ubiquitin ligase of the endoplasmic reticulum*. The Biochemical journal, 2005. **388**(Pt 2): p. 647-655.
33. Jo, Y., P.V. Sguigna, and R.A. DeBose-Boyd, *Membrane-associated ubiquitin ligase complex containing gp78 mediates sterol-accelerated degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase*. J Biol Chem, 2011. **286**(17): p. 15022-31.
34. Fang, S., et al., *The tumor autocrine motility factor receptor, gp78, is a ubiquitin protein ligase implicated in degradation from the endoplasmic reticulum*. Proc Natl Acad Sci U S A, 2001. **98**(25): p. 14422-7.
35. Jiang, J., et al., *CHIP is a U-box-dependent E3 ubiquitin ligase: identification of Hsc70 as a target for ubiquitylation*. J Biol Chem, 2001. **276**(46): p. 42938-44.
36. Younger, J.M., et al., *Sequential quality-control checkpoints triage misfolded cystic fibrosis transmembrane conductance regulator*. Cell, 2006. **126**(3): p. 571-82.
37. Jeon, Y.J., et al., *Regulation of glutamine carrier proteins by RNF5 determines breast cancer response to ER stress-inducing chemotherapies*. Cancer cell, 2015. **27**(3): p. 354-369.
38. Tcherpakov, M., et al., *Regulation of endoplasmic reticulum-associated degradation by RNF5-dependent ubiquitination of JNK-associated membrane protein (JAMP)*. J Biol Chem, 2009. **284**(18): p. 12099-109.
39. Ahner, A., et al., *Small heat shock proteins target mutant cystic fibrosis transmembrane conductance*

- regulator for degradation via a small ubiquitin-like modifier-dependent pathway.* Molecular biology of the cell, 2013. **24**(2): p. 74-84.
- 40. Menzies, S.A., et al., *The sterol-responsive RNF145 E3 ubiquitin ligase mediates the degradation of HMG-CoA reductase together with gp78 and Hrd1.* eLife, 2018. **7**: p. e40009.
 - 41. Lu, J.P., et al., *RNF170 protein, an endoplasmic reticulum membrane ubiquitin ligase, mediates inositol 1,4,5-trisphosphate receptor ubiquitination and degradation.* J Biol Chem, 2011. **286**(27): p. 24426-33.
 - 42. Lee, M.K., et al., *Transmembrane Protein pUL50 of Human Cytomegalovirus Inhibits ISGylation by Downregulating UBE1L.* J Virol, 2018. **92**(15).
 - 43. Wright, F.A., et al., *A Point Mutation in the Ubiquitin Ligase RNF170 That Causes Autosomal Dominant Sensory Ataxia Destabilizes the Protein and Impairs Inositol 1,4,5-Trisphosphate Receptor-mediated Ca²⁺ Signaling.* J Biol Chem, 2015. **290**(22): p. 13948-57.
 - 44. Maruyama, Y., et al., *Ubiquitin ligase Kif1 is involved in the endoplasmic reticulum-associated degradation pathway.* Biochem Biophys Res Commun, 2008. **374**(4): p. 737-41.
 - 45. El Khouri, E., et al., *RNF185 is a novel E3 ligase of endoplasmic reticulum-associated degradation (ERAD) that targets cystic fibrosis transmembrane conductance regulator (CFTR).* J Biol Chem, 2013. **288**(43): p. 31177-91.
 - 46. Stefanovic-Barrett, S., et al., *MARCH6 and TRC8 facilitate the quality control of cytosolic and tail-anchored proteins.* EMBO reports, 2018. **19**(5): p. e45603.
 - 47. Stagg, H.R., et al., *The TRC8 E3 ligase ubiquitinates MHC class I molecules before dislocation from the ER.* The Journal of cell biology, 2009. **186**(5): p. 685-692.
 - 48. Lerner, M., et al., *The RBCC gene RFP2 (Leu5) encodes a novel transmembrane E3 ubiquitin ligase involved in ERAD.* Mol Biol Cell, 2007. **18**(5): p. 1670-82.
 - 49. Altier, C., et al., *The Cavβ subunit prevents RFP2-mediated ubiquitination and proteasomal degradation of L-type channels.* Nat Neurosci, 2011. **14**(2): p. 173-80.
 - 50. Yoshida, Y., et al., *E3 ubiquitin ligase that recognizes sugar chains.* Nature, 2002. **418**(6896): p. 438-42.
 - 51. Gong, B., et al., *SCFFbx2-E3-ligase-mediated degradation of BACE1 attenuates Alzheimer's disease amyloidosis and improves synaptic function.* Aging cell, 2010. **9**(6): p. 1018-1031.
 - 52. Yoshida, Y., et al., *Glycoprotein-specific ubiquitin ligases recognize N-glycans in unfolded substrates.* EMBO Rep, 2005. **6**(3): p. 239-44.
 - 53. Yoshida, Y., et al., *Loss of peptide:N-glycanase causes proteasome dysfunction mediated by a sugar-recognizing ubiquitin ligase.* Proc Natl Acad Sci U S A, 2021. **118**(27).
 - 54. Yoshida, Y., et al., *Fbs2 is a new member of the E3 ubiquitin ligase family that recognizes sugar chains.* J Biol Chem, 2003. **278**(44): p. 43877-84.
 - 55. Binette, J., et al., *Requirements for the selective degradation of CD4 receptor molecules by the human immunodeficiency virus type 1 Vpu protein in the endoplasmic reticulum.* Retrovirology, 2007. **4**: p. 75-75.
 - 56. Chowdhury, A.M.M.A., et al., *Multiple regulatory mechanisms of the biological function of NRF3 (NFE2L3) control cancer cell proliferation.* Scientific reports, 2017. **7**(1): p. 12494-12494.
 - 57. Imai, Y., M. Soda, and R. Takahashi, *Parkin Suppresses Unfolded Protein Stress-induced Cell Death through Its E3 Ubiquitin-protein Ligase Activity**. Journal of Biological Chemistry, 2000. **275**(46): p. 35661-35664.
 - 58. Fry, W.H.D., et al., *Quantity control of the ErbB3 receptor tyrosine kinase at the endoplasmic reticulum.* Molecular and cellular biology, 2011. **31**(14): p. 3009-3018.
 - 59. Neutzner, A., et al., *A systematic search for endoplasmic reticulum (ER) membrane-associated RING*

- finger proteins identifies Nixin/ZNRF4 as a regulator of calnexin stability and ER homeostasis.* The Journal of biological chemistry, 2011. **286**(10): p. 8633-8643.
60. van den Boomen, D.J., et al., *TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-induced degradation of MHC-I.* Proc Natl Acad Sci U S A, 2014. **111**(31): p. 11425-30.
61. van de Weijer, M.L., et al., *The E3 Ubiquitin Ligase TMEM129 Is a Tri-Spanning Transmembrane Protein.* Viruses, 2016. **8**(11).
62. van de Weijer, M.L., et al., *A high-coverage shRNA screen identifies TMEM129 as an E3 ligase involved in ER-associated protein degradation.* Nat Commun, 2014. **5**: p. 3832.
63. Guo, X., et al., *The E3 ligase Smurfl regulates Wolfram syndrome protein stability at the endoplasmic reticulum.* J Biol Chem, 2011. **286**(20): p. 18037-47.
64. Li, X., et al., *The transmembrane endoplasmic reticulum-associated E3 ubiquitin ligase TRIM13 restrains the pathogenic-DNA-triggered inflammatory response.* Sci Adv, 2022. **8**(4): p. eabh0496.
65. Bianchini, E., et al., *Unveiling the degradative route of the V247M α -sarcoglycan mutant responsible for LGMD-2D.* Hum Mol Genet, 2014. **23**(14): p. 3746-58.
66. Kaneko, C., et al., *Characterization of the mouse gene for the U-box-type ubiquitin ligase UFD2a.* Biochem Biophys Res Commun, 2003. **300**(2): p. 297-304.
67. Liu, Y., et al., *USP13 antagonizes gp78 to maintain functionality of a chaperone in ER-associated degradation.* Elife, 2014. **3**: p. e01369.
68. Lee, M.J., et al., *Trimming of ubiquitin chains by proteasome-associated deubiquitinating enzymes.* Molecular & cellular proteomics : MCP, 2011. **10**(5): p. R110.003871-R110.003871.
69. Hassink, G.C., et al., *The ER-resident ubiquitin-specific protease 19 participates in the UPR and rescues ERAD substrates.* EMBO reports, 2009. **10**(7): p. 755-761.
70. Jung, E.S., et al., *Acute ER stress regulates amyloid precursor protein processing through ubiquitin-dependent degradation.* Scientific reports, 2015. **5**: p. 8805-8805.
71. Zhu, D., et al., *Deubiquitinating enzyme OTUB1 promotes cancer cell immunosuppression via preventing ER-associated degradation of immune checkpoint protein PD-L1.* Cell Death Differ, 2021. **28**(6): p. 1773-1789.
72. Ernst, R., et al., *The otubain YOD1 is a deubiquitinating enzyme that associates with p97 to facilitate protein dislocation from the ER.* Molecular cell, 2009. **36**(1): p. 28-38.
73. Zhong, X. and R.N. Pittman, *Ataxin-3 binds VCP/p97 and regulates retrotranslocation of ERAD substrates.* Hum Mol Genet, 2006. **15**(16): p. 2409-20.
74. Hubener, J., et al., *N-terminal ataxin-3 causes neurological symptoms with inclusions, endoplasmic reticulum stress and ribosomal dislocation.* Brain, 2011. **134**(Pt 7): p. 1925-42.
75. Kang, Y., et al., *Defining how ubiquitin receptors hHR23a and S5a bind polyubiquitin.* Journal of molecular biology, 2007. **369**(1): p. 168-176.
76. van Nocker, S., et al., *The multiubiquitin-chain-binding protein Mcb1 is a component of the 26S proteasome in *Saccharomyces cerevisiae* and plays a nonessential, substrate-specific role in protein turnover.* Molecular and cellular biology, 1996. **16**(11): p. 6020-6028.