

Electronic supplementary materials

<https://doi.org/10.1631/jzus.B1900210>

Difference in survival between right- versus left-sided colorectal neuroendocrine neoplasms

Ge-han XU¹, Hua-wei ZOU^{†‡2}, Ashley B. GROSSMAN³

¹*The Second Clinical College of China Medical University, Shenyang 110004, China*

²*1st Department of Oncology, Shengjing Hospital Affiliated to China Medical University, Shenyang 110022, China*

³*NET Unit, ENETS Centre of Excellence, Royal Free Hospital, London NW3 2QG, UK*

[†]E-mail: zhw20181102@sina.com

Table S1 Multivariable Cox regression of independent variables for cancer-specific risk of death

Variable	HR	OR (95%CI)	P
Location			
Left	1.00		
Right	2.13	1.31-3.46	0.002
Sex			
Men	1.00		
Women	0.86	0.77-0.95	0.005
Race			
White	1.00		
Black	1.01	0.86-1.20	0.894
Asian or Pacific islander	0.91	0.73-1.14	0.410
Unknown	1.05	0.34-3.28	0.933
Age			
<60	1.00		
≥60	1.53	1.37-1.71	<0.001
Tumor stage			
I	1.00		
II	2.40	1.03-5.58	0.043
III	13.10	6.71-25.58	<0.001
IV	50.77	26.25-98.20	<0.001
Tumor grade			
Well differentiated	1.00		
Moderately differentiated	2.22	1.73-2.84	<0.001
Poorly differentiated	5.96	4.85-7.33	<0.001
Undifferentiated	6.14	4.93-7.64	<0.001
Surgery			
No/Unknown	1.00		
Yes	0.75	0.67-0.84	<0.001