



Quantum Yang-Baxter equation and constant R -matrix over Grassmann algebra

DUPLIJ Steven[†], KOTULSKA Olga, SADOVNIKOV Alexander

(Department of Physics and Technology, V.N. Karazin Kharkov National University, Svoboda Sq. 4, Kharkov 61077, Ukraine)

[†]E-mail: Steven.A.Duplij@univer.kharkov.ua

Received Mar. 10, 2005; revision accepted June 25, 2005

Abstract: Constant solutions to Yang-Baxter equation are investigated over Grassmann algebra for the case of 6-vertex R -matrix. The general classification of all possible solutions over Grassmann algebra and particular cases with 2,3,4 generators are studied. As distinct from the standard case, when R -matrix over number field can have a maximum 5 nonvanishing elements, we obtain over Grassmann algebra a set of new full 6-vertex solutions. The solutions leading to regular R -matrices which appear in weak Hopf algebras are considered.

Key words: Constant solution, Grassmann algebra, Regularity, R -matrix

doi:10.1631/jzus.2005.A1065

Document code: A

CLC number: O13

INTRODUCTION

The Yang-Baxter equation (Yang, 1967; Baxter, 1972) is one of the most valuable concepts in modern theoretical physics (Baxter, 1982). The importance of detailed study of Yang-Baxter equation solutions is due to its key role in constant solution models of statistics mechanics (Baxter, 1972; 1982) and field theory in lower dimensions (Yang, 1967), conformal field theory (Di Francesco *et al.*, 1997) and in quantum integrable systems (Faddeev *et al.*, 1990). From the group-theoretical viewpoint, when classical Yang-Baxter equation is closely connected with classical (semi-simple) group theory, the quantum Yang-Baxter equation is the basis of modern quantum group theory (Drinfeld, 1987; Shnider and Sternberg, 1993; Demidov, 1998; Chari and Pressley, 1996). There exist a constant, one-parameter and double-parameter forms of the quantum Yang-Baxter equation (Lambe and Radford, 1997). The corresponding constant (and commutative) solutions to the Yang-Baxter equation (Hietarinta, 1992) are used in evolution of non-linear equation quantization, quantum group theory (Etingof *et al.*, 1997; 1999; Lu *et al.*,

2000; Gu, 1997) and knot theory (Kauffman, 1991; Turaev, 1994). The solution of the quantum Yang-Baxter equation is R -matrix (Kassel, 1995; Majid, 1995) [corresponding to the transfer-matrix in the lattice statistical models (Baxter, 1982)].

Recently the unitary Yang-Baxter equation solutions found an application in quantum calculations (Holievo, 2002; Kitajev *et al.*, 1999), while unitary R -matrix of a special type, acting on quantum state of two qubits, according to Brylinskis' theorem (Brylinski and Brylinski, 1994) can be treated as the universal quantum gate (Kauffman and Lomonaco, 2004; Zhang *et al.*, 2005; Dye, 2003; Duplij *et al.*, 2005).

The generalization of inverse quantum scattering method on supersymmetric systems (Khoroshkin and Tolstoy, 1991) and corresponding R -matrices were considered in (Chang *et al.*, 1992; Zhang and Gould, 1991). The building of supersymmetric analogues of the given constructions demands thorough consideration of Yang-Baxter equation solutions over the Grassmann algebra.

The constant Yang-Baxter equation solutions for the case of 6 vertices used for the description of the two-parameter quantum plane (Manin, 1989) and the

quantum gates of special type were considered in this work (Chang *et al.*, 1992; Zhang and Gould, 1991; Manin, 1989). Classification of general solutions is made and some particular cases are considered. In contrast to the standard case, when the R -matrix over the standard number field (for example, \mathbb{R}, \mathbb{C}) can have not more than 5 non-zero elements (Hietarinta, 1992; 1993) (actually 5-vertex solution), in our case (over the Grassmann algebra), all 6 elements can be non-zero. Thus, a new type of solution appears, which is absent in standard case (Hietarinta, 1992; 1993), is really complete 6-vertex solution. In the conclusion, the solutions leading to regular R -matrices appearing in weak Hopf algebras (Li and Duplij, 2002; Duplij and Li, 2001a) are considered.

YANG-BAXTER EQUATION OVER GRASSMANN ALGEBRA

Let V be a vector space, then on the tensor product $V^{\otimes n}$ we define a linear operator $R: V^{\otimes n} \rightarrow V^{\otimes n}$ in the following way. Let $\{e_i\}$ be a basis in V . Then we assign to the operator R a numerical matrix R with n index pairs

$$R(e_{i_1} \otimes \dots \otimes e_{i_n}) = R_{i_1 \dots i_n}^{j_1 \dots j_n} (e_{j_1} \otimes \dots \otimes e_{j_n}), \quad (1)$$

where over repeated indices summation is supposed. Consider n -simplex equation over the tensor product $V^{[\otimes n(n+1)/2]}$, where the linear operators R act trivially, for example, $R_{12}(e_{i_1} \otimes e_{i_2} \otimes e_{i_3}) = r_{i_1 i_2}^{j_1 j_2} (e_{j_1} \otimes e_{j_2} \otimes e_{i_3})$. In the general case with $K_\alpha \in \{1, \dots, N\}$, $N=n(n+1)/2$, the R operators are:

$$(R_{K_1 \dots K_n})_{i_1 \dots i_n}^{j_1 \dots j_n} = r_{i_{K_1} \dots i_{K_n}}^{j_{K_1} \dots j_{K_n}} \prod_{k=1, k \neq K_\alpha, \forall \alpha}^N \delta_{i_k}^{j_k}, \quad (2)$$

where $r_{i_{K_1} \dots i_{K_n}}^{j_{K_1} \dots j_{K_n}}$ is the matrix element of R -matrix. For example, 2-simplex constant equation is defined by Eq.(3)

$$R_{12} R_{13} R_{23} = R_{23} R_{13} R_{12}, \quad (3)$$

and is called Yang-Baxter equation (Baxter, 1982;

Lambe and Radford, 1997), and 3-simplex equation below

$$R_{123} R_{145} R_{246} R_{356} = R_{356} R_{246} R_{145} R_{123}, \quad (4)$$

is called tetrahedron equation in (Hietarinta, 1997) and 4-simplex equations was considered

$$R_{1234} R_{1567} R_{2589} R_{3680} R_{4790} = R_{4790} R_{3680} R_{2589} R_{1567} R_{1234}. \quad (5)$$

In terms of multiindex matrices, defined in Eq.(2), operator Eqs.(3)~(5) are as follows:

$$r_{j_2 j_3}^{k_2 k_3} r_{j_1 k_3}^{k_1 k_3} r_{k_1 k_2}^{l_1 l_2} = r_{j_1 j_2}^{k_1 k_2} r_{k_1 j_3}^{l_1 k_3} r_{k_2 k_3}^{l_2 l_3}, \quad (6.1)$$

$$r_{j_3 j_5 j_6}^{k_3 k_5 k_6} r_{j_2 j_4 k_6}^{k_2 k_4 k_6} r_{j_1 k_4 k_5}^{k_1 k_4 k_5} r_{k_1 k_2 k_3}^{l_1 l_2 l_3} = r_{j_1 j_2 j_3}^{k_1 k_2 k_3} r_{k_1 k_4 k_5}^{l_1 k_4 k_5} r_{k_2 k_4 j_6}^{l_2 k_4 k_6} r_{k_3 k_5 k_6}^{l_3 l_5 k_6}, \quad (6.2)$$

$$r_{j_4 j_7 j_9 j_0}^{k_4 k_7 k_9 k_0} r_{j_3 j_6 j_8 k_0}^{k_3 k_6 k_8 k_0} r_{j_2 j_5 k_8 k_9}^{k_2 k_5 k_8 k_9} r_{j_1 k_5 k_6 k_7}^{k_1 k_5 k_6 k_7} r_{k_1 k_2 k_3 k_4}^{l_1 l_2 l_3 l_4} = r_{j_1 j_2 j_3 j_4}^{k_1 k_2 k_3 k_4} r_{k_1 j_5 j_6 j_7}^{l_1 k_5 k_6 k_7} r_{k_2 k_5 j_8 j_9}^{l_2 k_5 k_8 k_9} r_{k_3 k_6 k_8 j_0}^{l_3 k_6 k_8 k_0} r_{k_4 k_7 k_9 k_0}^{l_4 k_7 k_9 k_0}. \quad (6.3)$$

General formulation of similar (n -simplex) equations was given in (Carter and Saito, 1996), and permutation equations were studied in (Hietarinta, 1993; 1997).

Here we will consider constant solutions for 2-simplex Eq.(6.1) (Yang-Baxter equation) over Grassmann algebra, being a particular case of a superalgebra (Berezin, 1983; Kac, 1977), which is an important step on the way of consistent supersymmetric generalization of Yang-Baxter equation and its constant solutions (Khoroshkin and Tolstoy, 1991; Zhang, 1991; Links *et al.*, 1994).

Let A be a commutative superalgebra over a field K (where $K=\mathbb{R}, \mathbb{C}$) with the dissociation on the direct sum $A = A_0 \oplus A_1$ (Berezin, 1983; Kac, 1977).

Elements a from A_0 and A_1 are homogeneous with regard to parity $p(a) \stackrel{\text{def}}{=} \{\bar{i} \in \{\bar{0}, \bar{1}\} = Z_2 \mid a \in A_{\bar{i}}\}$. Supercommutator is defined as $[a, b] = ab - (-1)^{p(a)p(b)} ba$. In the particular case A_n is the Grassmann algebra with the element ξ_1, \dots, ξ_n , which satisfies $\xi_i \xi_j + \xi_j \xi_i = 0, 1 \leq i, j \leq n$, in particular $\xi_i^2 = 0$ (n can be infinite). Superalgebra structure in A_n is defined by the fact that parity of the element is considered equal to $p(\xi_i) = \bar{1}$ (Leites, 1980). Then even $x \in A_0$ and odd

$\chi \in \mathcal{A}_\Gamma$ elements of Grassmann algebra can be expanded into the series (which is finite under finite number of elements ξ_i)

$$x = x_{\text{numb}} + x_{\text{nil}} = x_0 + x_{12}\xi_1\xi_2 + x_{13}\xi_1\xi_3 + \dots$$

$$= x_{\text{numb}} + x_{i_1 \dots i_{2r}} \xi_{i_1} \dots \xi_{i_{2r}}, \tag{7.1}$$

$$\chi = \chi_{\text{nil}} + x_1\xi_1 + x_2\xi_2 + \dots + x_{123}\xi_1\xi_2\xi_3 + \dots$$

$$= x_{i_1 \dots i_{2r-1}} \xi_{i_1} \dots \xi_{i_{2r-1}}, \tag{7.2}$$

where $x_{i_1 \dots i_n} \in K$. The mapping ε , discarding odd elements, is called numerical mapping (Manin, 1984; De Witt, 1992) [canonical projection (Leites, 1983), body map (Rogers, 1980; Rabin, 1987; 1991)] and it acts on the elements in Eq.(7) as $\varepsilon(x) = x|_{\xi_i=0} = x_{\text{numb}}$, $\varepsilon(\chi) = \chi|_{\xi_i=0} = 0$. From Eq.(7) it follows that, for example, the equations $x^2=0$, $\chi x=0$ and $\chi\chi'=0$ can have non-zero and non-trivial solutions (zero divisors and nilpotents), which can essentially extend the number of solutions for different equations including the Yang-Baxter equation. For example, in \mathcal{A}_4 even non-zero nilpotents $x^2=0$ satisfy

$$x_0 = 0, x_{12}x_{34} - x_{13}x_{24} + x_{14}x_{23} = 0, \tag{8}$$

and for component non-zero zero dividers $\chi x=0$, we get

$$\left. \begin{aligned} x_0 &= 0, \\ x_1x_{23} - x_2x_{13} + x_3x_{12} &= 0, \\ x_1x_{24} - x_2x_{14} + x_4x_{12} &= 0, \\ x_1x_{34} - x_3x_{14} + x_4x_{13} &= 0, \\ x_2x_{34} - x_3x_{24} + x_4x_{23} &= 0. \end{aligned} \right\} \tag{9}$$

For $\chi\chi'=0$ we get the conditions

$$x_i x'_j - x_j x'_i = 0, \quad i, j = 1, 2, 3, 4 \tag{10}$$

which shows that such odd objects in Eq.(7.2) are nilpotents of the second nilpotency degree $n^2=0$.

Let us consider R -matrix over even part of Grassmann algebra, for example, with 4 elements, let us write its dissociation onto numerical and nilpotent parts

$$R = R^{(0)} + R^{(12)}\xi_1\xi_2 + R^{(13)}\xi_1\xi_3 + R^{(14)}\xi_1\xi_4 + R^{(23)}\xi_2\xi_3 + R^{(24)}\xi_2\xi_4 + R^{(34)}\xi_3\xi_4 + R^{(1234)}\xi_1\xi_2\xi_3\xi_4, \tag{11}$$

the components of Yang-Baxter equation can be represented in the following way

$$\left. \begin{aligned} R_{12} &= R_{12}^{(0)} + R_{12}^{(12)}\xi_1\xi_2 + R_{12}^{(13)}\xi_1\xi_3 + R_{12}^{(14)}\xi_1\xi_4 + \\ &R_{12}^{(23)}\xi_2\xi_3 + R_{12}^{(24)}\xi_2\xi_4 + R_{12}^{(34)}\xi_3\xi_4 + R_{12}^{(1234)}\xi_1\xi_2\xi_3\xi_4, \\ R_{13} &= R_{13}^{(0)} + R_{13}^{(12)}\xi_1\xi_2 + R_{13}^{(13)}\xi_1\xi_3 + R_{13}^{(14)}\xi_1\xi_4 + \\ &R_{13}^{(23)}\xi_2\xi_3 + R_{13}^{(24)}\xi_2\xi_4 + R_{13}^{(34)}\xi_3\xi_4 + R_{13}^{(1234)}\xi_1\xi_2\xi_3\xi_4, \\ R_{23} &= R_{23}^{(0)} + R_{23}^{(12)}\xi_1\xi_2 + R_{23}^{(13)}\xi_1\xi_3 + R_{23}^{(14)}\xi_1\xi_4 + \\ &R_{23}^{(23)}\xi_2\xi_3 + R_{23}^{(24)}\xi_2\xi_4 + R_{23}^{(34)}\xi_3\xi_4 + R_{23}^{(1234)}\xi_1\xi_2\xi_3\xi_4. \end{aligned} \right\} \tag{12}$$

Substitute the given expressions into Eq.(3) and we will get the system of equations for components

$$R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(0)} = R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(0)}, \tag{13.1}$$

$$R_{12}^{(0)} R_{13}^{(\alpha)} R_{23}^{(0)} + R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(\alpha)} + R_{12}^{(\alpha)} R_{13}^{(0)} R_{23}^{(0)}$$

$$= R_{23}^{(\alpha)} R_{13}^{(0)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(\alpha)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(\alpha)},$$

$$\alpha = 12, 13, 14, 23, 24, 34, \tag{13.2}$$

$$\begin{aligned} &R_{12}^{(0)} R_{13}^{(12)} R_{23}^{(34)} - R_{12}^{(0)} R_{13}^{(13)} R_{23}^{(24)} + R_{12}^{(0)} R_{13}^{(14)} R_{23}^{(23)} + \\ &R_{12}^{(0)} R_{13}^{(23)} R_{23}^{(14)} - R_{12}^{(0)} R_{13}^{(24)} R_{23}^{(13)} + R_{12}^{(0)} R_{13}^{(34)} R_{23}^{(12)} + \\ &R_{12}^{(12)} R_{13}^{(0)} R_{23}^{(34)} - R_{12}^{(13)} R_{13}^{(0)} R_{23}^{(24)} + R_{12}^{(14)} R_{13}^{(0)} R_{23}^{(23)} + \\ &R_{12}^{(23)} R_{13}^{(0)} R_{23}^{(14)} - R_{12}^{(24)} R_{13}^{(0)} R_{23}^{(13)} + R_{12}^{(34)} R_{13}^{(0)} R_{23}^{(12)} + \\ &R_{12}^{(12)} R_{13}^{(34)} R_{23}^{(0)} - R_{12}^{(13)} R_{13}^{(24)} R_{23}^{(0)} + R_{12}^{(14)} R_{13}^{(23)} R_{23}^{(0)} + \\ &R_{12}^{(23)} R_{13}^{(14)} R_{23}^{(0)} - R_{12}^{(24)} R_{13}^{(13)} R_{23}^{(0)} + R_{12}^{(34)} R_{13}^{(12)} R_{23}^{(0)} + \\ &R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(1234)} + R_{12}^{(0)} R_{13}^{(1234)} R_{23}^{(0)} + R_{12}^{(1234)} R_{13}^{(0)} R_{23}^{(0)} + \\ &= R_{23}^{(1234)} R_{13}^{(0)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(1234)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(1234)} + \\ &R_{23}^{(0)} R_{13}^{(12)} R_{12}^{(34)} - R_{23}^{(0)} R_{13}^{(13)} R_{12}^{(34)} + R_{23}^{(0)} R_{13}^{(14)} R_{12}^{(23)} + \\ &R_{23}^{(0)} R_{13}^{(23)} R_{12}^{(14)} - R_{23}^{(0)} R_{13}^{(24)} R_{12}^{(13)} + R_{23}^{(0)} R_{13}^{(34)} R_{12}^{(12)} + \\ &R_{23}^{(12)} R_{13}^{(0)} R_{12}^{(34)} - R_{23}^{(13)} R_{13}^{(0)} R_{12}^{(34)} + R_{23}^{(14)} R_{13}^{(0)} R_{12}^{(23)} + \\ &R_{23}^{(23)} R_{13}^{(0)} R_{12}^{(14)} - R_{23}^{(24)} R_{13}^{(0)} R_{12}^{(13)} + R_{23}^{(34)} R_{13}^{(0)} R_{12}^{(12)} + \\ &R_{23}^{(12)} R_{13}^{(34)} R_{12}^{(0)} - R_{23}^{(13)} R_{13}^{(24)} R_{12}^{(0)} + R_{23}^{(14)} R_{13}^{(23)} R_{12}^{(0)} + \\ &R_{23}^{(23)} R_{13}^{(14)} R_{12}^{(0)} - R_{23}^{(24)} R_{13}^{(13)} R_{12}^{(0)} + R_{23}^{(34)} R_{13}^{(12)} R_{12}^{(0)}, \end{aligned} \tag{13.3}$$

where Eq.(13.1) represents the standard constant Yang-Baxter equation for matrices over a numerical field. All possible solutions of Eq.(13.1) were derived in (Hietarinta, 1992; 1993). In general, using these

solutions by means of Eq.(13), all corresponding solutions classes over the Grassmann algebra can be obtained.

In the particular case, when there exists the symmetry

$$R^{(12)} = R^{(13)} = R^{(14)} = R^{(23)} = R^{(24)} = R^{(34)} = R^{(1)}, \quad (14)$$

then Eq.(13) simplifies and is represented in the following way

$$\left. \begin{aligned} &R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(0)} = R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(0)}, \\ &R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(1)} + R_{12}^{(0)} R_{13}^{(1)} R_{23}^{(0)} + R_{12}^{(1)} R_{13}^{(0)} R_{23}^{(0)} \\ &= R_{23}^{(1)} R_{13}^{(0)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(1)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(1)}, \\ &2(R_{12}^{(0)} R_{13}^{(1)} R_{23}^{(1)} + R_{12}^{(1)} R_{13}^{(0)} R_{23}^{(1)} + R_{12}^{(1)} R_{13}^{(1)} R_{23}^{(0)}) + \\ &R_{12}^{(0)} R_{13}^{(0)} R_{23}^{(1234)} + R_{12}^{(0)} R_{13}^{(1234)} R_{23}^{(0)} + R_{12}^{(1234)} R_{13}^{(0)} R_{23}^{(0)} \\ &= 2(R_{23}^{(1)} R_{13}^{(1)} R_{12}^{(0)} + R_{23}^{(1)} R_{13}^{(0)} R_{12}^{(1)} + R_{23}^{(0)} R_{13}^{(1)} R_{12}^{(1)}) + \\ &R_{23}^{(1234)} R_{13}^{(0)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(1234)} R_{12}^{(0)} + R_{23}^{(0)} R_{13}^{(0)} R_{12}^{(1234)}. \end{aligned} \right\} \quad (15)$$

Thus, there are solutions, satisfying the given system, but with non-zero nilpotent part.

From the formal point of view, purely nilpotent solutions [not containing numerical parts in Eq.(11)] exist with any number of Grassmann generators. For cubic nilpotent equation to be different from zero, it is necessary to have at least 6 generators of the Grassmann algebra. That is why for the number of generators less than 6 pure nilpotent solution [under $R^{(0)}=0$] will be any matrix, as the equations for the components of R -matrix are homogeneous of the third degree.

If we consider R -matrix of the general type with all non-zero elements, the Yang-Baxter equation gives the system of 64 equations with 16 unknowns (Hietarinta, 1997). Considering expansions over the Grassmann algebra Eq.(11), the number of equations and the number of variables grow essentially. That is why we will consider only 6-vertex solutions.

NONINVERTIBLE SOLUTIONS OF SOME EQUATIONS OVER GRASSMANN ALGEBRA

Let us consider the set of equations, which arise in finding the matrix solutions of Yang-Baxter equation over the even part of Grassmann algebra

$$ax=0, \quad (16)$$

$$abc=0, \quad (17)$$

$$a^2=b^2. \quad (18)$$

Consider the case when a is invertible. Then multiplying the left side of Eq.(16) by a^{-1} , we get the only solution $x=0$. Analogous arguments are true for Eq.(17) too. From Eq.(18) we write the equation onto numerical parts a and b : $a_0^2=b_0^2$. From here it follows, that a and b are invertible and noninvertible simultaneously. That is why the solutions of this equation for every value of the numerical part can differ in the nilpotent part only. Let it be assumed that for every a there exist two solutions b and $b' = b + \tilde{b}$, where \tilde{b} is nilpotent. Then $b^2 = b'^2 + 2b\tilde{b} + \tilde{b}^2$, from which it follows, that $\tilde{b}(2b + \tilde{b}) = 0$. The numerical part of the expression in brackets equals $2b_0$, i.e. it is invertible. That is why $\tilde{b} = 0$. Consequently, the solution of Eq.(18) will be $b = \pm a$.

Thus, new solutions can only exist in the class of purely nilpotent a . That is why we will consider further only nilpotent elements included into Eq.(16).

For Eq.(16) it will be further shown that for any a we can construct x . Then for Eq.(17) the derived result can be used by substitution into Eq.(16) a over ab , and x over c . That is, for arbitrary ab we can construct c .

Eq.(18) by substitution $a=e+f$, $b=e-f$ can be transformed into the type of Eq.(16): $ef=0$.

In Eq.(16) for every nilpotent a we will find all possible b . For the solution of Eq.(16) with a given a , it is convenient to represent it in the form of products sum of the Grassmann algebra generators with invertible parameter. For further computations it will be useful to write the solutions of Eq.(16) for some character values of a .

Type 1: a is represented by one term

$$a = \xi_{i_1} \cdots \xi_{i_n} \tilde{a}, \tilde{a}_0 \neq 0.$$

As \tilde{a} is invertible, then Eq.(16) is equivalent to the equation

$$\xi_{i_1} \cdots \xi_{i_n} x = 0. \quad (19)$$

For the solution of this equation, let us present x in the form of the sum of all possible ordered products of elements. Then in Eq.(19) the terms without ξ_i will

remain. As the product of generators included into x are ordered, then under each product of generators in Eq.(19) there will be only one numerical term. That is why the general solution of Eq.(19) will be

$$x = \sum_{k=1}^n \xi_{i_k} \tilde{x}_k,$$

where \tilde{x}_k are arbitrary, possibly invertible parameters of the Grassmann algebra. Then it is necessary that x contains at least one term included into $\xi_{i_1} \dots \xi_{i_l}$.

Type 2: variable a has the following presentation

$$a = \xi_{i_1} \dots \xi_{i_l} a_1 + \xi_{j_1} \dots \xi_{j_m} a_2, \tag{20}$$

There are sets of indices, which do not intersect $\{i_1, \dots, i_l\} \cap \{j_1, \dots, j_m\} = \emptyset$. Without loss of generality it can be assumed that $a_2=1$. Then x can be conveniently presented in the form of two terms $x=x_1+x_2$.

In the first of them two terms $\xi_{i_1} \dots \xi_{i_l} x_1 = 0$ and $\xi_{j_1} \dots \xi_{j_m} x_1 = 0$ vanish. Then using the result of the previous case, we get

$$x_1 = \sum_{p=1}^l \sum_{q=1}^m \xi_{i_p} \xi_{j_q} x_{pq},$$

where x_{pq} are arbitrary elements.

The variable x_2 will be defined from the condition that x_2 is not zero, when at least one of the terms in Eq.(20) does not vanish. For the sum to be nonzero, as $\{i_1, \dots, i_l\} \cap \{j_1, \dots, j_m\} = \emptyset$, it is necessary for x_2 to be

$$x_2 = \xi_{j_1} \dots \xi_{j_m} x_3 + \xi_{i_1} \dots \xi_{i_l} x_4, \tag{21}$$

where x_3 and x_4 are arbitrary parameters which do not contain $\xi_{i_1}, \dots, \xi_{i_l}, \xi_{j_1}, \dots, \xi_{j_m}$. Then Eq.(16) is equivalent to the equation

$$a'_1 x_3 + (-1)^{ml} x_4 = 0,$$

where $\xi_{j_1} \dots \xi_{j_m} a'_1 = a_1 \xi_{j_1} \dots \xi_{j_m}$ and a'_1 does not contain terms containing $\xi_{j_1}, \dots, \xi_{j_m}$. Then

$$x_2 = (\xi_{j_1} \dots \xi_{j_m} + \xi_{i_1} \dots \xi_{i_l} (-1)^{ml+1} a'_1) x_3.$$

Type 3: a is presented in the form of two terms, as in the previous case, except that sets of the indices intersect $\{i_1, \dots, i_l\} \cap \{j_1, \dots, j_m\} \neq \emptyset$. Therefore a can be written in the following form

$$a = \xi_{i_1} \dots \xi_{i_l} (\xi_{j_1} \dots \xi_{j_m} a_1 + \xi_{k_1} \dots \xi_{k_n}), \tag{22}$$

$\{i_1, \dots, i_l\} \cap \{j_1, \dots, j_m\} = \emptyset, \{i_1, \dots, i_l\} \cap \{k_1, \dots, k_n\} = \emptyset, \{j_1, \dots, j_m\} \cap \{k_1, \dots, k_n\} = \emptyset$. Then parameter x can be presented in the form of three terms

$$x = x_1 + x_2 + x_3, \tag{23}$$

In the first term $\xi_{i_1} \dots \xi_{i_l}$ vanishes, i.e.

$$\xi_{i_1} \dots \xi_{i_l} x_1 = 0.$$

Then using the result of the first case, we get

$$x_1 = \sum_{p=1}^l \xi_{i_p} \tilde{x}_p.$$

In the second term both terms in the bracket vanish simultaneously, i.e. Eq.(22)

$$\xi_{j_1} \dots \xi_{j_m} x_2 = 0 \wedge \xi_{k_1} \dots \xi_{k_n} x_2 = 0.$$

Then in analogy with the previous case

$$x_2 = \sum_{p=1}^m \sum_{q=1}^n \xi_{j_p} \xi_{k_q} x_{pq}.$$

In the third term the multiplier in the bracket vanishes as in the previous case. The type x_3 looks similar to Eq.(21)

$$x_3 = \xi_{k_1} \dots \xi_{k_n} x_4 + \xi_{j_1} \dots \xi_{j_m} x_5,$$

where x_4 and x_5 do not contain $\xi_{j_1}, \dots, \xi_{j_m}, \xi_{k_1}, \dots, \xi_{k_n}$.

Then Eq.(16) is equivalent to the equation

$$a'_1 x_4 + (-1)^{ml} x_5 = 0,$$

where $\xi_{k_1} \dots \xi_{k_n} a'_1 = a_1 \xi_{k_1} \dots \xi_{k_n}$ and a'_1 does not have

terms containing $\xi_{k_1}, \dots, \xi_{k_n}$. Then, finally

$$x_3 = (\xi_{j_1} \cdots \xi_{j_m} + \xi_{i_1} \cdots \xi_{i_l} (-1)^{ml+1} a') x_4.$$

Using the obtained results we can construct the general type of solution for Eq.(16) under the given type a

$$a = \sum_{k=1}^n \alpha_k a_k, \tag{24}$$

where $\alpha_k = \xi_{i_1} \cdots \xi_{i_l}$, i_1, \dots, i_l are in increasing order, a_k are invertible parameters not containing $\xi_{i_1} \cdots \xi_{i_l}$.

The general process of finding solutions is as follows: let us present x in the form of the sum $n+1$ of terms, where n is the number of terms in Eq.(24)

$$x = \sum_{i=0}^n x_i.$$

Every x_i vanishes simultaneously in the i terms in Eq.(24), that such terms are chosen as any, i.e., each x_i

is the sum $x_i = \sum_{j=1}^{\binom{n}{i}} x_{ij}$, in each term of which vanishes

some concrete selection C_{ij} from n over i terms in Eq.(24)

$$\forall k \in C_{ij}, a_k x_{ij} = 0, \sum_{k \in C_{ij}} a_k x_{ij} = 0. \tag{25}$$

Cases $j=n$ and $j=n-1$ coincide, as all the sum vanishes and the sum of $n-1$ of its terms also vanishes, then n th term vanishes from each term.

The situation can be such that for some selection C_{ij} , some terms will vanish in the sum in Eq.(25). Then such case will become the case with greater i . That is why it is necessary to begin with greater i in the search for x . The same order of finding x is defined by the way of finding the solutions for quotient types a .

Thus, the given scheme allows getting all possible solutions of Eq.(16).

Consider the examples. Let us find some concrete even nilpotent equation solutions of Eq.(16)

over Grassmann algebra with 2,3,4,5 elements.

Case 1: For elements of the type

$$a = a_{12} \xi_1 \xi_2, \quad b = b_{12} \xi_1 \xi_2,$$

Eq.(16) is satisfied $\forall a_{12}, b_{12}$, as $\xi_i^2=0$. Identical situation is true for 3 elements

$$a = a_{12} \xi_1 \xi_2 + a_{13} \xi_1 \xi_3 + a_{23} \xi_2 \xi_3, \\ b = b_{12} \xi_1 \xi_2 + b_{13} \xi_1 \xi_3 + b_{23} \xi_2 \xi_3,$$

and all the coefficients are arbitrary.

Case 2: For elements of the type

$$a = a_{12} \xi_1 \xi_2 + a_{13} \xi_1 \xi_3 + a_{14} \xi_1 \xi_4 + a_{23} \xi_2 \xi_3 + \\ a_{24} \xi_2 \xi_4 + a_{34} \xi_3 \xi_4 + a_{1234} \xi_1 \xi_2 \xi_3 \xi_4, \\ b = b_{12} \xi_1 \xi_2 + b_{13} \xi_1 \xi_3 + b_{14} \xi_1 \xi_4 + b_{23} \xi_2 \xi_3 + \\ b_{24} \xi_2 \xi_4 + b_{34} \xi_3 \xi_4 + b_{1234} \xi_1 \xi_2 \xi_3 \xi_4,$$

Eq.(16) comes to the following expression for the coefficients

$$a_{12} b_{34} - a_{13} b_{24} + a_{14} b_{23} + a_{23} b_{14} - a_{24} b_{13} + a_{34} b_{12} = 0,$$

but a_{1234}, b_{1234} remain undefined. Thus, the solution will be 13-parametrical. Analogous situation exists for 5 elements, then

$$a = a_{12} \xi_1 \xi_2 + a_{13} \xi_1 \xi_3 + a_{14} \xi_1 \xi_4 + a_{15} \xi_1 \xi_5 + \\ a_{23} \xi_2 \xi_3 + a_{24} \xi_2 \xi_4 + a_{25} \xi_2 \xi_5 + a_{34} \xi_3 \xi_4 + \\ a_{35} \xi_3 \xi_5 + a_{45} \xi_4 \xi_5 + a_{1234} \xi_1 \xi_2 \xi_3 \xi_4 + a_{1235} \xi_1 \xi_2 \xi_3 \xi_5 + \\ a_{1245} \xi_1 \xi_2 \xi_4 \xi_5 + a_{1345} \xi_1 \xi_3 \xi_4 \xi_5 + a_{2345} \xi_2 \xi_3 \xi_4 \xi_5, \\ b = b_{12} \xi_1 \xi_2 + b_{13} \xi_1 \xi_3 + b_{14} \xi_1 \xi_4 + b_{15} \xi_1 \xi_5 + \\ b_{23} \xi_2 \xi_3 + b_{24} \xi_2 \xi_4 + b_{25} \xi_2 \xi_5 + b_{34} \xi_3 \xi_4 + \\ b_{35} \xi_3 \xi_5 + b_{45} \xi_4 \xi_5 + b_{1234} \xi_1 \xi_2 \xi_3 \xi_4 + b_{1235} \xi_1 \xi_2 \xi_3 \xi_5 + \\ b_{1245} \xi_1 \xi_2 \xi_4 \xi_5 + b_{1345} \xi_1 \xi_3 \xi_4 \xi_5 + b_{2345} \xi_2 \xi_3 \xi_4 \xi_5.$$

Eq.(16) gives the system

$$a_{12} b_{34} - a_{13} b_{24} + a_{14} b_{23} + a_{23} b_{14} - a_{24} b_{13} + a_{34} b_{12} = 0, \\ a_{12} b_{35} - a_{13} b_{25} + a_{15} b_{23} + a_{23} b_{15} - a_{25} b_{13} + a_{35} b_{12} = 0, \\ a_{12} b_{45} - a_{14} b_{25} + a_{15} b_{24} + a_{24} b_{15} - a_{25} b_{14} + a_{45} b_{12} = 0, \\ a_{13} b_{45} - a_{14} b_{35} + a_{15} b_{34} + a_{34} b_{15} - a_{35} b_{14} + a_{45} b_{13} = 0, \\ a_{23} b_{45} - a_{24} b_{35} + a_{25} b_{34} + a_{34} b_{25} - a_{35} b_{24} + a_{45} b_{23} = 0,$$

where coefficients $a_{1234}, a_{1235}, a_{1245}, a_{1345}, a_{2345}, b_{1234}, b_{1235}, b_{1245}, b_{1345}, b_{2345}$ are undefined. The solution will be 25-parametrical. It is easy to see that in the case of 6 elements there will be equations on coefficients of the new type.

Thus, it can be assumed that the solutions of Eq.(16) over Grassmann algebra with $2n$ and $2n+1$ generators are defined by the same conditions.

Then we consider the Yang-Baxter equation solution over Grassmann algebra.

CLASSIFICATION OF YANG-BAXTER EQUATION SOLUTIONS WITH THE NUMBER OF VERTICES, LESS OR EQUAL TO SIX

According to (Kassel, 1995), 6-vertex solution for the Yang-Baxter equation is R -matrix of the type

$$R = \begin{pmatrix} p & \cdot & \cdot & \cdot \\ \cdot & c & d & \cdot \\ \cdot & a & b & \cdot \\ \cdot & \cdot & \cdot & q \end{pmatrix}, \tag{26}$$

where a, b, c, d, p, q are even elements of the Grassmann algebra. From Yang-Baxter Eq.(3) it follows, that R -matrix is defined precisely up to a constant (scale symmetry), so that it is always possible to define the normalization of the element, whose numerical part differs from zero. As it is unknown beforehand (before classification), which of the elements in Eq.(3) has non-zero numerical part, we are not going to do the normalization Eq.(26).

From Eq.(2) and Eq.(3), the evident type of matrix R_{12}, R_{13}, R_{23} leads to

$$R_{12} = \begin{pmatrix} p & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & p & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & c & \cdot & d & \cdot & \cdot \\ \cdot & \cdot & \cdot & c & \cdot & d & \cdot \\ \cdot & \cdot & a & \cdot & b & \cdot & \cdot \\ \cdot & \cdot & \cdot & a & \cdot & b & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q \end{pmatrix},$$

$$R_{13} = \begin{pmatrix} p & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & c & \cdot & \cdot & d & \cdot & \cdot \\ \cdot & \cdot & p & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & c & \cdot & \cdot & d \\ \cdot & a & \cdot & \cdot & b & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & q & \cdot \\ \cdot & \cdot & \cdot & a & \cdot & \cdot & b \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q \end{pmatrix},$$

$$R_{23} = \begin{pmatrix} p & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & c & d & \cdot & \cdot & \cdot & \cdot \\ \cdot & a & b & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & q & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & p & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & c & d \\ \cdot & \cdot & \cdot & \cdot & \cdot & a & b \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q \end{pmatrix}. \tag{27}$$

Let us substitute R_{12}, R_{13}, R_{23} into Eq.(6.1) and get the following (essentially redefined) system of equations

$$cda = 0, \tag{28}$$

$$bda = 0, \tag{29}$$

$$da(d - a) = 0, pd(d - p) + cbd = 0, \tag{30}$$

$$\left. \begin{aligned} qd(d - q) + cbd &= 0, \\ pa(a - p) + cba &= 0, \\ qa(a - q) + cba &= 0. \end{aligned} \right\} \tag{31}$$

Let us present all the parameters in the form of sum of numerical part and even nilpotent part as $x = x_0 + \tilde{x}$, where $x = a, b, c, d, p, q$. It is clear that Eqs.(28)~(31) must be separately satisfied by the numerical part, so that the classification will be done according to it.

Because in Eq.(28) $cda=0$, it is convenient to classify the solutions for the numerical part with vanishing a_0 and d_0 .

(1) Both a_0 and d_0 differ from zero: $d_0, a_0 \neq 0 \rightarrow b_0 = c_0 = 0, a_0 = d_0, \{p_0, q_0\} = \{0, a_0\}$, where $\{\}$ means the set of elements. Then for the numerical part of R -matrix we derive 4-vertex solution

$$R^{(0)} = \begin{pmatrix} \{0, a_0\} & \cdot & \cdot & \cdot \\ \cdot & \cdot & a_0 & \cdot \\ \cdot & a_0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \{0, a_0\} \end{pmatrix} \sim \begin{pmatrix} \{0, 1\} & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \{0, 1\} \end{pmatrix}, \tag{32}$$

where the last equivalency follows from the normalization because $a_0 \neq 0$.

(2) One of the elements a_0 or d_0 differs from zero:

(a) $a_0 = 0$,

$$d_0 \neq 0 \rightarrow \{p_0, q_0\} = \left\{ \frac{d_0}{2} \pm \sqrt{\frac{d_0^2}{4} + b_0 c_0} \right\},$$

then for the numerical part of R -matrix we get 5-vertex solution

$$R^{(0)} = \begin{pmatrix} \frac{d_0}{2} \pm \sqrt{\frac{d_0^2}{4} + b_0 c_0} & \cdot & \cdot & \cdot \\ \cdot & c_0 & d_0 & \cdot \\ \cdot & \cdot & b_0 & \cdot \\ \cdot & \cdot & \cdot & \frac{d_0}{2} \pm \sqrt{\frac{d_0^2}{4} + b_0 c_0} \end{pmatrix}. \tag{33}$$

(b) Or for $d_0 = 0, a_0 \neq 0 \rightarrow \{p_0, q_0\} = \{a_0/2 \pm \sqrt{a_0^2/4 + b_0 c_0}\}$, then again we get 5-vertex solution

$$R^{(0)} = \begin{pmatrix} \frac{a_0}{2} \pm \sqrt{\frac{a_0^2}{4} + b_0 c_0} & \cdot & \cdot & \cdot \\ \cdot & c_0 & \cdot & \cdot \\ \cdot & a_0 & b_0 & \cdot \\ \cdot & \cdot & \cdot & \frac{a_0}{2} \pm \sqrt{\frac{a_0^2}{4} + b_0 c_0} \end{pmatrix}. \tag{34}$$

(3) Both numerical parts of the elements a and d equal zero: $a_0=d_0=0 \rightarrow p_0, q_0, b_0, c_0$ are any, and the numerical part of R -matrix becomes diagonal (4-vertex numerical solution)

$$R^{(0)} = \begin{pmatrix} p_0 & \cdot & \cdot & \cdot \\ \cdot & c_0 & \cdot & \cdot \\ \cdot & \cdot & b_0 & \cdot \\ \cdot & \cdot & \cdot & q_0 \end{pmatrix}. \tag{35}$$

Now let us continue the classification based on nilpotent parts. Using the fact that the element is invertible, if its numerical part differs from zero, we get the final answer for the first case:

(1) $a=d$ is any even invertible: $b=c=0, \{p, q\}=\{0, a\}$, then we have 2- and 4-vertex solutions

$$R = \begin{pmatrix} \{0, a\} & \cdot & \cdot & \cdot \\ \cdot & \cdot & a & \cdot \\ \cdot & a & \cdot & \cdot \\ \cdot & \cdot & \cdot & \{0, a\} \end{pmatrix} \sim \begin{pmatrix} \{0, 1\} & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot \\ \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \{0, 1\} \end{pmatrix}, \tag{36}$$

where the last equivalency is derived from Yang-Baxter equation scale symmetry.

In the second case the invertibility d leads to the vanishing of nilpotent parameter a from Eq.(28). Then the remaining simplified system of Eqs.(30)~(31) can be solved according to parameters p and q . Equations on these parameters coincide

$$p^2=pd+bc, q^2=qd+bc. \tag{37}$$

If the quadratic equation discriminant has a nonzero numerical part $d_0^2/4+b_0c_0 \neq 0$, then the solution can be written in the following way.

(2) Under $a=0$ we have:

(a) $d_0^2/4+b_0c_0 \neq 0 \rightarrow \{p, q\} = d/2 \pm \sqrt{d^2/4+bc}$, then we get 5-vertex solution

$$R = \begin{pmatrix} \frac{d}{2} \pm \sqrt{\frac{d^2}{4} + bc} & \cdot & \cdot & \cdot \\ \cdot & c & d & \cdot \\ \cdot & \cdot & b & \cdot \\ \cdot & \cdot & \cdot & \frac{d}{2} \pm \sqrt{\frac{d^2}{4} + bc} \end{pmatrix}. \tag{38}$$

If the discriminant is equal to zero, i.e., $d_0^2/4+b_0c_0=0$, then Eq.(38) is useless, as it is impos-

sible to find the square root from nilpotent expression. Then from Eqs.(28)~(31) and Eq.(37) it is seen that the vanishing discriminant numerical part leads to invertibility of parameters b, c, p, q . Suppose (using scale symmetry) $d=1$. The equations on the remaining nilpotent parts p and q are given as

$$\tilde{p}^2 = \tilde{q}^2. \tag{39}$$

Then we get:

(b) $d_0^2/4+b_0c_0=0 \rightarrow b, c, p, q$ is invertible and $p_0 = q_0 = d_0/2, c = (\tilde{p}^2 - 1/4)/b, b_0c_0 = -1/4, p_0 = q_0 = 1/2$.

So we get again 5-vertex solution

$$R = \begin{pmatrix} 1/2 + \tilde{p} & \cdot & \cdot & \cdot \\ \cdot & \frac{\tilde{p}^2 - 1/4}{b} & 1 & \cdot \\ \cdot & \cdot & b & \cdot \\ \cdot & \cdot & \cdot & 1/2 + \tilde{q} \end{pmatrix}. \tag{40}$$

In this case we have the set of solutions in comparison with the previous case due to nilpotency. The main difference from the previous case Eq.(38) is that the parameters p and q are now connected by Eq.(39) on nilpotent parts, which gives great freedom in choosing their meanings.

(3) In the third case, when the numerical part has a diagonal form, we can point out four sub-cases.

(a) $p_0^2 - b_0c_0 \neq 0$ or $q_0^2 - b_0c_0 \neq 0 \rightarrow a=d=0, p, q, b, c$ are any, then we get diagonal R -matrix (4-vertex solution)

$$R = \begin{pmatrix} p & \cdot & \cdot & \cdot \\ \cdot & c & \cdot & \cdot \\ \cdot & \cdot & b & \cdot \\ \cdot & \cdot & \cdot & q \end{pmatrix}. \tag{41}$$

(b) $p_0^2 - b_0c_0 = 0$ and $q_0^2 - b_0c_0 = 0$. Suppose $p=1$, then $p_0^2 = q_0^2 = b_0c_0 = 1$,

$$ad=0, \tag{42}$$

$$a \left(a + \frac{\tilde{c}}{c_0} + \frac{\tilde{b}}{b_0} + \frac{\tilde{b} \tilde{c}}{b_0 c_0} \right) = 0, \tag{43}$$

$$d \left(d + \frac{\tilde{c}}{c_0} + \frac{\tilde{b}}{b_0} + \frac{\tilde{b} \tilde{c}}{b_0 c_0} \right) = 0. \tag{44}$$

The numerical part of the parameter q because of $q_0^2=1$ can have the meaning of ± 1 . Consequently the equations for the nilpotent part \tilde{q} lead to two different systems of equations

$$q_0 = +1 \rightarrow a\tilde{q} = 0, d\tilde{q} = 0, \tag{45}$$

$$q_0 = -1 \rightarrow a(a - \tilde{q}) = 0, d(d - \tilde{q}) = 0. \tag{46}$$

We rewrite Eq.(46), using Eqs.(43)~(44), in linear form over a and d as

$$\left. \begin{aligned} q_0 = +1 &\rightarrow a\tilde{q} = 0, d\tilde{q} = 0, \\ q_0 = -1 &\rightarrow a(\tilde{q} - f) = 0, d(\tilde{q} - f) = 0. \end{aligned} \right\} \tag{47}$$

where $f = -\frac{\tilde{b}}{b_0} - \frac{\tilde{c}}{c_0} - \frac{\tilde{b} \tilde{c}}{b_0 c_0}$.

(c) $p_0^2 - b_0c_0 = 0$ and $q_0^2 - b_0c_0 = 0 \rightarrow p_0 = q_0 = b_0 = 0, c_0 \neq 0 \rightarrow c = 1$ and Eqs.(42)~(44) have the following form

$$\left. \begin{aligned} ad &= 0, \\ a(p^2 - pa - b) &= 0, d(p^2 - pd - b) = 0, \\ a(q^2 - qa - b) &= 0, d(q^2 - qd - b) = 0. \end{aligned} \right\} \tag{48}$$

(d) All numerical parts are equal to zero: $p_0 = q_0 = b_0 = c_0 = a_0 = d_0 = 0$.

Eqs.(42)~(44) and Eq.(48) reveal that only in Cases 3(b) and 3(c) we can expect full 6-vertex solutions, which are absent in the classification (Hietarinta, 1993; 1997).

Now let us consider the concrete type of solutions when the number of Grassmann algebra generators is 2, 3, and 4.

YANG-BAXTER EQUATION SOLUTION OVER GRASSMANN ALGEBRA WITH TWO ELEMENTS

We present the expansion of each element of R -matrix Eq.(26) over 2 Grassmann generators as

$$\left. \begin{aligned} c &= c_0 + c_{12}\xi_1\xi_2, \quad b = b_0 + b_{12}\xi_1\xi_2, \quad d = d_0 + d_{12}\xi_1\xi_2, \\ p &= p_0 + p_{12}\xi_1\xi_2, \quad q = q_0 + q_{12}\xi_1\xi_2, \quad a = a_0 + a_{12}\xi_1\xi_2. \end{aligned} \right\} \quad (49)$$

$$\left. \begin{aligned} d_0 &= 0, \quad a_0 = 0, \\ p_0^2 d_{12} - c_0 b_0 d_{12} &= 0, \quad q_0^2 d_{12} - c_0 b_0 d_{12} = 0, \\ p_0^2 a_{12} - c_0 b_0 a_{12} &= 0, \quad q_0^2 a_{12} - c_0 b_0 a_{12} = 0. \end{aligned} \right\} \quad (53)$$

Substitution of these equations into the system of Eqs.(28)~(31) yields the system of equations below

$$\begin{aligned} c_0 d_0 a_0 &= 0, \quad b_0 d_0 a_0 = 0, \quad d_0 a_0 (d_0 - a_0) = 0, \quad (50) \\ p_0 d_0 (d_0 - p_0) + c_0 b_0 d_0 &= 0, \quad q_0 d_0 (d_0 - q_0) + c_0 b_0 d_0 = 0, \\ p_0 a_0 (a_0 - p_0) + c_0 b_0 a_0 &= 0, \quad q_0 a_0 (a_0 - q_0) + c_0 b_0 a_0 = 0, \\ c_0 d_0 a_{12} + c_0 a_0 d_{12} + d_0 a_0 c_{12} &= 0, \\ b_0 d_0 a_{12} + b_0 a_0 d_{12} + d_0 a_0 b_{12} &= 0, \\ d_0 a_0 (d_{12} - a_{12}) + d_0 (d_0 - a_0) a_{12} + a_0 (d_0 - a_0) d_{12} &= 0, \\ p_0 d_0 (d_{12} - p_{12}) + p_0 (d_0 - p_0) d_{12} + \\ d_0 (d_0 - p_0) p_{12} + c_0 b_0 d_{12} + c_0 d_0 b_{12} + b_0 d_0 c_{12} &= 0, \\ q_0 d_0 (d_{12} - q_{12}) + q_0 (d_0 - q_0) d_{12} + d_0 (d_0 - q_0) q_{12} + \\ c_0 b_0 d_{12} + c_0 d_0 b_{12} + b_0 d_0 c_{12} &= 0, \\ p_0 a_0 (a_{12} - p_{12}) + p_0 (a_0 - p_0) a_{12} + a_0 (a_0 - p_0) p_{12} + \\ c_0 b_0 a_{12} + c_0 a_0 b_{12} + b_0 a_0 c_{12} &= 0, \\ q_0 a_0 (a_{12} - q_{12}) + q_0 (a_0 - q_0) a_{12} + a_0 (a_0 - q_0) q_{12} + \\ c_0 b_0 a_{12} + c_0 a_0 b_{12} + b_0 a_0 c_{12} &= 0. \end{aligned} \quad (51)$$

Eq.(50) will be considered as the key ones. We find the solutions for all non-zero elements c, b, d, p, q, a as numerical and nilpotent parts do not vanish simultaneously, which essentially narrows the solutions classes.

Case I: $d_0=0$. Eqs.(50)~(51) have the following form:

$$\left. \begin{aligned} d_0 &= 0, \quad p_0 a_0 (a_0 - p_0) + c_0 b_0 a_0 = 0, \\ q_0 a_0 (a_0 - q_0) + c_0 b_0 a_0 &= 0, \end{aligned} \right\} \quad (52.1)$$

$$a_0^2 d_{12} = 0, \quad (52.2)$$

$$c_0 a_0 d_{12} = 0, \quad b_0 a_0 d_{12} = 0, \quad (52.3)$$

$$p_0^2 d_{12} - c_0 b_0 d_{12} = 0, \quad q_0^2 d_{12} - c_0 b_0 d_{12} = 0, \quad (52.4)$$

$$\begin{aligned} p_0 a_0 (a_{12} - p_{12}) + p_0 (a_0 - p_0) a_{12} + a_0 (a_0 - p_0) p_{12} + \\ c_0 b_0 a_{12} + c_0 a_0 b_{12} + b_0 a_0 c_{12} &= 0, \end{aligned} \quad (52.5)$$

$$\begin{aligned} q_0 a_0 (a_{12} - q_{12}) + q_0 (a_0 - q_0) a_{12} + a_0 (a_0 - q_0) q_{12} + \\ c_0 b_0 a_{12} + c_0 a_0 b_{12} + b_0 a_0 c_{12} &= 0. \end{aligned} \quad (52.6)$$

From Eq.(52.2) it follows, that $a_0=0$ and

From Eq.(53) it follows, that $p_0^2=q_0^2=c_0 b_0$ (because of the assumption, none of the 6 elements equals zero). Insert the parameters $p_0=q_0=t, c_0=r \neq 0$, then $b_0=t^2/r$, as nilpotent parts are not defined from Eq.(53), insert 6 more parameters $c_{12}=v, b_{12}=w, d_{12}=y, p_{12}=l, q_{12}=m, a_{12}=n$. The final solution will be an 8-parametric one

$$\left. \begin{aligned} c &= r + v\xi_1\xi_2, \quad b = t^2 / r + w\xi_1\xi_2, \quad d = y\xi_1\xi_2, \\ p &= t + l\xi_1\xi_2, \quad q = t + m\xi_1\xi_2, \quad a = n\xi_1\xi_2. \end{aligned} \right\} \quad (54)$$

Let us write the resulting 6-vertex R -matrix

$$\begin{aligned} R &= \begin{pmatrix} t + l\xi_1\xi_2 & 0 & 0 & 0 \\ 0 & r + v\xi_1\xi_2 & y\xi_1\xi_2 & 0 \\ 0 & n\xi_1\xi_2 & t^2 / r + w\xi_1\xi_2 & 0 \\ 0 & 0 & 0 & t + m\xi_1\xi_2 \end{pmatrix} \\ &= \begin{pmatrix} t & 0 & 0 & 0 \\ 0 & r & 0 & 0 \\ 0 & 0 & t^2 / r & 0 \\ 0 & 0 & 0 & t \end{pmatrix} + \begin{pmatrix} l & 0 & 0 & 0 \\ 0 & v & y & 0 \\ 0 & n & w & 0 \\ 0 & 0 & 0 & m \end{pmatrix} \xi_1\xi_2. \end{aligned} \quad (55)$$

According to the classification of the previous section, such solution refers to Case 3(b). Evidently, if we considered R -matrix over numerical field, this solution would be 4-vertex, but not 6 (Hietarinta, 1997). Under $t=0$ we obtain 6-vertex R -matrix, in which only 1 element is invertible, that in our classification corresponds to Case 3(c).

Case II: $a_0=0$. Eqs.(50)~(51) lead to the following:

$$a_0 = 0, \quad (56.1)$$

$$\left. \begin{aligned} p_0 d_0 (d_0 - p_0) + c_0 b_0 d_0 &= 0, \\ q_0 d_0 (d_0 - q_0) + c_0 b_0 d_0 &= 0, \end{aligned} \right\} \quad (56.2)$$

$$c_0 d_0 a_{12} = 0, \quad b_0 d_0 a_{12} = 0, \quad (56.3)$$

$$d_0^2 a_{12} = 0, \quad (56.4)$$

$$p_0 d_0 (d_{12} - p_{12}) + p_0 (d_0 - p_0) d_{12} + d_0 (d_0 - p_0)$$

$$\times p_{12} + c_0 b_0 d_{12} + c_0 d_0 b_{12} + b_0 d_0 c_{12} = 0, \quad (56.5)$$

$$q_0 d_0 (d_{12} - q_{12}) + q_0 (d_0 - q_0) d_{12} + d_0 (d_0 - q_0) q_{12} +$$

$$c_0 b_0 d_{12} + c_0 d_0 b_{12} + b_0 d_0 c_{12} = 0, \quad (56.6)$$

$$p_0^2 a_{12} - c_0 b_0 a_{12} = 0, \quad q_0^2 a_{12} - c_0 b_0 a_{12} = 0. \quad (56.7)$$

From Eq.(56.4) it follows, that $d_0=0$, then Eq.(56) becomes to Eq.(53) and the solutions coincide with Case I.

Case III: $c_0=b_0=0, d_0=a_0$. Then Eqs.(50)~(51) are as follows:

$$c_0 = 0, \quad b_0 = 0, \quad d_0 = a_0, \quad (57.1)$$

$$p_0 d_0 (d_0 - p_0) = 0, \quad q_0 d_0 (d_0 - q_0) = 0, \quad (57.2)$$

$$d_0 a_0 c_{12} = 0, \quad d_0 a_0 b_{12} = 0, \quad (57.3)$$

$$d_0 a_0 (d_{12} - a_{12}) = 0, \quad (57.4)$$

$$p_0 d_0 (d_{12} - p_{12}) + p_0 (d_0 - p_0) d_{12} + d_0 (d_0 - p_0) p_{12} = 0, \quad (57.5)$$

$$q_0 d_0 (d_{12} - q_{12}) + q_0 (d_0 - q_0) d_{12} + d_0 (d_0 - q_0) q_{12} = 0. \quad (57.6)$$

Eq.(57.3) shows that $a_0=b_0=c_0=d_0=0$, then the coefficients $p_0, q_0, a_{12}, b_{12}, c_{12}, d_{12}, p_{12}, q_{12}$ remain undefined. The solution will also be 8-parametrical $p_0=t, q_0=r, a_{12}=l, b_{12}=m, c_{12}=n, d_{12}=k, p_{12}=w, q_{12}=s$ and can be represented in the following form

$$R = \begin{pmatrix} t + w\xi_1\xi_2 & 0 & 0 & 0 \\ 0 & n\xi_1\xi_2 & k\xi_1\xi_2 & 0 \\ 0 & l\xi_1\xi_2 & m\xi_1\xi_2 & 0 \\ 0 & 0 & 0 & r + s\xi_1\xi_2 \end{pmatrix} = \begin{pmatrix} t & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & r \end{pmatrix} + \begin{pmatrix} w & 0 & 0 & 0 \\ 0 & n & k & 0 \\ 0 & l & m & 0 \\ 0 & 0 & 0 & s \end{pmatrix} \xi_1\xi_2. \quad (58)$$

This solution is 6-vertex **R**-matrix with 2 invertible elements, which is not contained in the Hietarinta classification (Hietarinta, 1992; 1993).

SOLUTIONS OF YANG-BAXTER EQUATIONS OVER GRASSMAN ALGEBRA WITH THREE GENERATORS

Let us write the expansion of each element of **R**-matrix Eq.(26) over 3 generators as

$$c = c_0 + c_{12}\xi_1\xi_2 + c_{13}\xi_1\xi_3 + c_{23}\xi_2\xi_3, \\ b = b_0 + b_{12}\xi_1\xi_2 + b_{13}\xi_1\xi_3 + b_{23}\xi_2\xi_3, \\ d = d_0 + d_{12}\xi_1\xi_2 + d_{13}\xi_1\xi_3 + d_{23}\xi_2\xi_3, \\ p = p_0 + p_{12}\xi_1\xi_2 + p_{13}\xi_1\xi_3 + p_{23}\xi_2\xi_3, \\ q = q_0 + q_{12}\xi_1\xi_2 + q_{13}\xi_1\xi_3 + q_{23}\xi_2\xi_3, \\ a = a_0 + a_{12}\xi_1\xi_2 + a_{13}\xi_1\xi_3 + a_{23}\xi_2\xi_3. \quad (59)$$

Substituting Eq.(59) into Eqs.(28)~(31) yields the equations in the components

$$c_0 d_0 a_0 = 0, \quad b_0 d_0 a_0 = 0, \quad d_0 a_0 (d_0 - a_0) = 0, \quad (60)$$

$$\left. \begin{aligned} p_0 d_0 (d_0 - p_0) + c_0 b_0 d_0 &= 0, \\ q_0 d_0 (d_0 - q_0) + c_0 b_0 d_0 &= 0, \\ p_0 a_0 (a_0 - p_0) + c_0 b_0 a_0 &= 0, \\ q_0 a_0 (a_0 - q_0) + c_0 b_0 a_0 &= 0, \\ c_0 d_0 a_{12} + c_0 a_0 d_{12} + d_0 a_0 c_{12} &= 0, \\ c_0 d_0 a_{13} + c_0 a_0 d_{13} + d_0 a_0 c_{13} &= 0, \\ c_0 d_0 a_{23} + c_0 a_0 d_{23} + d_0 a_0 c_{23} &= 0, \\ b_0 d_0 a_{12} + b_0 a_0 d_{12} + d_0 a_0 b_{12} &= 0, \\ b_0 d_0 a_{13} + b_0 a_0 d_{13} + d_0 a_0 b_{13} &= 0, \\ b_0 d_0 a_{23} + b_0 a_0 d_{23} + d_0 a_0 b_{23} &= 0 \end{aligned} \right\} \quad (61)$$

$$d_0 a_0 (d_{12} - a_{12}) + d_0 (d_0 - a_0) a_{12} + a_0 (d_0 - a_0) d_{12} = 0, \\ d_0 a_0 (d_{13} - a_{13}) + d_0 (d_0 - a_0) a_{13} + a_0 (d_0 - a_0) d_{13} = 0, \\ d_0 a_0 (d_{23} - a_{23}) + d_0 (d_0 - a_0) a_{23} + a_0 (d_0 - a_0) d_{23} = 0. \quad (62)$$

and

$$\left. \begin{aligned} p_0 d_0 (d_\beta - p_\beta) + p_0 (d_0 - p_0) d_\beta + d_0 (d_0 - p_0) p_\beta + \\ c_0 b_0 d_\beta + c_0 d_0 b_\beta + b_0 d_0 c_\beta &= 0, \\ q_0 d_0 (d_\gamma - q_\gamma) + q_0 (d_0 - q_0) d_\gamma + d_0 (d_0 - q_0) q_\gamma + \\ c_0 b_0 d_\gamma + c_0 d_0 b_\gamma + b_0 d_0 c_\gamma &= 0, \\ p_0 a_0 (a_\delta - p_\delta) + p_0 (a_0 - p_0) a_\delta + a_0 (a_0 - p_0) p_\delta + \\ c_0 b_0 a_\delta + c_0 a_0 b_\delta + b_0 a_0 c_\delta &= 0, \\ q_0 a_0 (a_\varepsilon - q_\varepsilon) + q_0 (a_0 - q_0) a_\varepsilon + a_0 (a_0 - q_0) q_\varepsilon + \\ c_0 b_0 a_\varepsilon + c_0 a_0 b_\varepsilon + b_0 a_0 c_\varepsilon &= 0, \\ \delta, \gamma, \beta, \varepsilon &= 12, 13, 23. \end{aligned} \right\} \quad (63)$$

In symmetric case, when components under 2 generators are equal in all elements $a_{12}=a_{13}=a_{23}$ etc., the system simplifies and comes to the case of 2 elements, as nilpotency index of non-numerical part in both cases equals 2.

As for two Grassmann generators, Eq.(60) will be clarifying, mark 3 cases.

Case I: $d_0=0$, then the system has the form

$$\left. \begin{aligned} d_0 &= 0, \\ p_0 a_0 (a_0 - p_0) + c_0 b_0 a_0 &= 0, \\ q_0 a_0 (a_0 - q_0) + c_0 b_0 a_0 &= 0, \\ c_0 a_0 d_{12} = 0, c_0 a_0 d_{13} = 0, c_0 a_0 d_{23} &= 0, \\ b_0 a_0 d_{12} = 0, b_0 a_0 d_{13} = 0, b_0 a_0 d_{23} &= 0, \\ a_0^2 d_{12} = 0, a_0^2 d_{13} = 0, a_0^2 d_{23} &= 0, \\ p_0^2 d_{12} - c_0 b_0 d_{12} = 0, p_0^2 d_{13} - c_0 b_0 d_{13} &= 0, \\ p_0^2 d_{23} - c_0 b_0 d_{23} = 0, q_0^2 d_{12} - c_0 b_0 d_{12} &= 0, \\ q_0^2 d_{13} - c_0 b_0 d_{13} = 0, q_0^2 d_{23} - c_0 b_0 d_{23} &= 0. \end{aligned} \right\} \quad (64)$$

and

$$\begin{aligned} p_0 a_0 (a_\tau - p_\tau) + p_0 (a_0 - p_0) a_\tau + a_0 (a_0 - p_0) p_\tau \\ + c_0 b_0 a_\tau + c_0 a_0 b_\tau + b_0 a_0 c_\tau &= 0, \\ q_0 a_0 (a_\psi - q_\psi) + q_0 (a_0 - q_0) a_\psi + a_0 (a_0 - q_0) q_\psi \\ + c_0 b_0 a_\psi + c_0 a_0 b_\psi + b_0 a_0 c_\psi &= 0, \end{aligned} \quad (65)$$

$\tau, \psi = 12, 13, 23.$

From Eq.(64) it follows, that $a_0=0$, as we suppose, that **R**-matrix contains 6 non-zero elements. The system is as follows:

$$\left. \begin{aligned} d_0 &= 0, a_0 = 0, \\ p_0^2 d_{12} - c_0 b_0 d_{12} = 0, p_0^2 d_{13} - c_0 b_0 d_{13} &= 0, \\ p_0^2 d_{23} - c_0 b_0 d_{23} &= 0, \\ q_0^2 d_{12} - c_0 b_0 d_{12} = 0, q_0^2 d_{13} - c_0 b_0 d_{13} &= 0, \\ q_0^2 d_{23} - c_0 b_0 d_{23} &= 0, \\ p_0^2 a_{12} - c_0 b_0 a_{12} = 0, p_0^2 a_{13} - c_0 b_0 a_{13} &= 0, \\ p_0^2 a_{23} - c_0 b_0 a_{23} &= 0, \\ q_0^2 a_{12} - c_0 b_0 a_{12} = 0, q_0^2 a_{13} - c_0 b_0 a_{13} &= 0, \\ q_0^2 a_{23} - c_0 b_0 a_{23} &= 0. \end{aligned} \right\} \quad (66)$$

As d_{12}, d_{13}, d_{23} simultaneously cannot be equal to zero (we suppose that all 6 elements of **R**-matrix do

not come to zero, including the complete element $d \neq 0$), then Eq.(66) shows that $p_0^2 = q_0^2 = c_0 b_0$. Insert parameters $p_0 = q_0 = t, c_0 = r$, then $b_0 = t^2/r$, and the components $d_{12}, d_{13}, d_{23}, a_{12}, a_{13}, a_{23}, c_{12}, c_{13}, c_{23}, b_{12}, b_{13}, b_{23}, p_{12}, p_{13}, p_{23}, q_{12}, q_{13}, q_{23}$ remain undefined, i.e. the solutions will be 20-parametrical.

By analogy with expansion on 2 Grassmann generators Case II coincides with Case I, and Case III is obtained from Case I by putting $c_0 = b_0 = 0$.

SOLUTIONS OF YANG-BAXTER EQUATION OVER GRASSMANN ALGEBRA WITH FOUR ELEMENTS

The classification of all solutions was obtained earlier and Cases 1, 2(a), 3(a) require no further study, as the solutions are given in evident form Eqs.(36), (38), (40), (41) for any number of Grassmann algebra generators. The remaining Cases 2(b), 3(b), 3(c), 3(d) contain nilpotent equations. In the general case, for nilpotents it is impossible to give concrete expressions for their solutions. That is why these cases require further analysis.

As expressions of the third degree on Grassmann algebra with four elements equal zero, Eqs.(28)~(31) are simplified in the following way. Case 2(b) contains the equation of the second degree that is why it remains without changes

$$\tilde{p}^2 = \tilde{q}^2. \quad (67)$$

For Case 3(b) the system of Eqs.(45)~(46) also remains unchanged, but the expressions are changed for $f = -\tilde{b}/b_0 - \tilde{c}/c_0$.

In Case 3(c), Eq.(48) is quadratic, which is why they do not change

$$ab=0, db=0, ad=0. \quad (68)$$

In Case 3(d) all parameters p, q, a, b, c, d can be arbitrary, even nilpotent.

For the solutions of these systems, let us write all the parameters in the component form

$$x = \sum_{\substack{i,j=1 \\ i < j}}^4 x_{ij} \xi_i \xi_j + x_{1234} \xi_1 \xi_2 \xi_3 \xi_4, \quad (69)$$

where $x=a, b, c, d, p, q$. Coordinate x_{1234} can be arbitrary, as it cancels any such nilpotent. Then all the components in the equation will be as follows

Case 2(b)

$$p_{12}p_{34} + p_{14}p_{23} - p_{13}p_{24} = q_{12}q_{34} + q_{14}q_{23} - q_{13}q_{24}. \quad (70)$$

The solution can be, for example, under $p_{12} \neq 0$

$$p_{34} = (-p_{14}p_{23} + p_{13}p_{24} + q_{12}q_{34} + q_{14}q_{23} - q_{13}q_{24}) / p_{12}. \quad (71)$$

Case 3(b)

$$\begin{aligned} q_{12}a_{34} - q_{13}a_{24} + q_{14}a_{23} + q_{23}a_{14} - q_{24}a_{13} + q_{34}a_{12} &= 0, \\ q_{12}d_{34} - q_{13}d_{24} + q_{14}d_{23} + q_{23}d_{14} - q_{24}d_{13} + q_{34}d_{12} &= 0, \\ [2(a_{12}q'_{13} - a_{13}q'_{12}) + (q_{12}f_{13} - q_{13}f_{12})]a_{24} - \\ [2(a_{12}q'_{14} - a_{14}q'_{12}) + (q_{12}f_{14} - q_{14}f_{12})]a_{23} - \\ [2a_{12}q'_{23} + (q_{12}f_{23} - q_{23}f_{12})]a_{14} + \\ [2a_{12}q'_{24} + (q_{12}f_{24} - q_{24}f_{12})]a_{13} - \\ [2a_{12}q'_{34} + (q_{12}f_{34} - q_{34}f_{12})]a_{12} &= 0, \\ [2(d_{12}q'_{13} - d_{13}q'_{12}) + (q_{12}f_{13} - q_{13}f_{12})]d_{24} - \\ [2(d_{12}q'_{14} - d_{14}q'_{12}) + (q_{12}f_{14} - q_{14}f_{12})]d_{23} - \\ [2d_{12}q'_{23} + (q_{12}f_{23} - q_{23}f_{12})]d_{14} + \\ [2d_{12}q'_{24} + (q_{12}f_{24} - q_{24}f_{12})]d_{13} - \\ [2d_{12}q'_{34} + (q_{12}f_{34} - q_{34}f_{12})]d_{12} &= 0, \\ (a_{12}q'_{13} - a_{13}q'_{12})d_{24} - (a_{12}q'_{14} - a_{14}q'_{12})d_{23} - \\ (a_{12}q'_{23} - a_{23}q'_{12})d_{14} + (a_{12}q'_{24} - a_{24}q'_{12})d_{13} + \\ [-2a_{12}q'_{34} + (a_{13}q'_{24} + a_{24}q'_{13}) - (a_{14}q'_{23} + a_{23}q'_{14})]d_{12} &= 0, \end{aligned}$$

here $q' = \tilde{q}$ in the case “+” and $q' = \tilde{q} - f$ in the case “-”.

Case 3(c)

$$\begin{aligned} -b_{12}a_{34} + b_{13}a_{24} - b_{14}a_{23} - b_{23}a_{14} + b_{24}a_{13} - b_{34}a_{12} &= 0, \\ -b_{12}d_{34} + b_{13}d_{24} - b_{14}d_{23} - b_{23}d_{14} + b_{24}d_{13} - b_{34}d_{12} &= 0, \\ (a_{12}b_{13} - a_{13}b_{12})d_{24} - (a_{12}b_{14} - a_{14}b_{12})d_{23} - \\ (a_{12}b_{23} - a_{23}b_{12})d_{14} + (a_{12}b_{24} - a_{24}b_{12})d_{13} + \\ [-2a_{12}b_{34} + (a_{13}b_{24} + a_{24}b_{13}) - (a_{14}b_{23} + a_{23}b_{14})]d_{12} &= 0. \end{aligned}$$

This system of equations can be solved, for example, if $b_{12} \neq 0$ and $(a_{12}b_{13} - a_{13}b_{12}) \neq 0$. Then we can express a_{34}, d_{34}, d_{24} through other variables. The obtained systems for such coordinates are linear, that is

why their solutions are rational functions of the rest of the components. Final expressions are too cumbersome to be presented here.

Only in Cases 3(b), 3(c), 3(d) there exist 6-vertex solutions, in contrast to the solutions over the numerical field in which only 5-vertex solutions are possible (Hietarinta, 1992; 1993).

REGULAR YANG-BAXTER EQUATION SOLUTIONS

In (Duplij and Sadovnikov, 2002) set-theoretical (Drinfeld, 1992; Etingof *et al.*, 1999; Gu, 1997) regular (according to von Neumann) solutions for Yang-Baxter equation were found. The consideration of similar solutions for 6-vertex R -matrix is interesting. Let us remind that matrix R is regular (von Neumann), if there exists matrix \bar{R} such that $R\bar{R}R = R, \bar{R}R\bar{R} = \bar{R}$ (Penrose, 1955; Rao and Mitra, 1971; Clifford and Preston, 1972). Such R -matrices appeared during the study of weak Hopf algebras (Li and Duplij, 2002; Duplij and Li, 2001a; 2001b). Properties of regular supermatrices were considered in (Duplij, 2000; Duplij and Kotulskaja, 2002).

For 6-vertex R -matrix we will use, instead of the “reversibility” condition

$$R^{21}R = R R^{21} = id, \quad (72)$$

which is called the unitarity in (Etingof *et al.*, 1997; 1999), the Von Neumann regularity in the form

$$R\bar{R}^{21}R = R, \bar{R}^{21}R\bar{R}^{21} = \bar{R}^{21}, \quad (73)$$

where

$$\bar{R}^{21} = \begin{pmatrix} p & \cdot & \cdot & \cdot \\ \cdot & b & a & \cdot \\ \cdot & d & c & \cdot \\ \cdot & \cdot & \cdot & q \end{pmatrix}. \quad (74)$$

Notice that non-unitary [with violation Eq.(72)] set-theoretical solutions to Yang-Baxter equation of another form were considered in (Lu *et al.*, 2000; Soloviev, 2002).

Substitution of Eq.(74) into Eq.(73) gives limitations, on the condition of regularity on the elements of 6-vertex R -matrix Eq.(26)

$$\begin{aligned}
 p^3 &= p, q^3 = q, \\
 (ab + bd)d + (a^2 + cb)b &= b, \\
 (ab + bd)c + (a^2 + cb)a &= a, \\
 (cb + d^2)d + (ca + dc)b &= d, \\
 (cb + d^2)c + (ca + dc)a &= c.
 \end{aligned}
 \tag{75}$$

The analysis shows that regular noninvertible solutions (satisfying $R^{21}R \neq id$) can appear in cases: (1) under $a = \pm 1$, p and q are simultaneously not equal to a ; 2(a) under $d = \pm 1$, $a = 0$, $bc = 0$, $p = \{0, \pm 1\}$, $q = \{0, \pm 1\}$. For example, for a particular case from Case 1 we have

$$\begin{aligned}
 R &= \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \bar{R}^{21} = \begin{pmatrix} a & 0 & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\
 R\bar{R}^{21} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a^2 & 0 & 0 \\ 0 & 0 & a^2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \neq id.
 \end{aligned}
 \tag{76}$$

but regularity Eq.(73) is fulfilled. In a particular case from Case 2(a) we get

$$\begin{aligned}
 R &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & c & 1 & 0 \\ 0 & 0 & b & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \bar{R}^{21} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 1 & c & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
 R\bar{R}^{21} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & c & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \neq id.
 \end{aligned}
 \tag{77}$$

and regularity Eq.(73) is fulfilled.

Thus, we presented the classification of constant 6-vertex Yang-Baxter equation solutions. Concrete examples were given for Cases 2, 3, 4 of Grassmann algebra elements and the corresponding analysis was done. The obtained solutions can be used for supersymmetric quantum group generalizations and quantum computations.

ACKNOWLEDGMENTS

The authors are grateful to G.C. Kourinnoy, B.V. Novikov, S.D. Sinelshchikov, L.L. Vaksman for their fruitful discussions and to L. Kauffman, Fang Li, Jian-Zu Zhang for useful comments during correspondence. We are thankful to T.L. Kudryashova for language checking.

References

- Baxter, R.J., 1972. Partition function for the eight-vertex model. *Ann. Phys.*, **70**:193-228.
- Baxter, R.J., 1982. *Exactly Solved Models in Statistical Mechanics*. Academic Press, London.
- Berezin, F.A., 1983. *Vvedeniye v Algebru i Analiz s Antikommutirujushimi Peremennymi*. Izd-vo MGU, Moskva, p.208.
- Brylinski, J.L., Brylinski, R., 1994. *Universal Quantum Gates. Mathematics of Quantum Computation*. Chapman & Hall/CRC Press, Boca Raton, p.124-134.
- Carter, J.S., Saito, M., 1996. On formulation and solutions of simplex equations. *J. Mod. Phys.*, **A11**:4453-4463.
- Chang, D., Phillips, I., Rozansky, I., 1992. R -matrix approach to quantum superalgebras. *J. Math. Phys.*, **33**(11):3710-3715.
- Chari, V., Pressley, A., 1996. *A Guide to Quantum Groups*. Cambridge University Press, Cambridge.
- Clifford, A., Preston, G., 1972. *Algebraic Theory of Semi-Groups*. Mir, Moskva, p.283.
- De Witt, B.S., 1992. *Supermanifolds*. 2nd Edition. Cambridge University Press, Cambridge, p.407.
- Demidov, E.E., 1998. *Quantum Groups*. Faktorial, Moskva, p.146.
- Di Francesco, P., Mathieu, P., Sénéchal, D., 1997. *Conformal Field Theory*. Springer-Verlag, Berlin, p.890.
- Drinfeld, V.G., 1987. *Quantum Groups*. Proceedings of the ICM. Phode Island. AMS, Berkeley, p.798-820.
- Drinfeld, V.G., 1992. On some unsolved problems in quantum group theory. *Lect. Notes. Math.*, **1510**:1-8.
- Duplij, S.A., 2000. *Semisupermanifolds and Semigroups*. Krok, Kharkov, p.220.
- Duplij, S.A., 2003. On supermatrix idempotent operator semigroups. *Linear Algebra Appl.*, **360**:59-81.
- Duplij, S.A., Li, F., 2001a. On regular solutions of quantum Yang-Baxter equation and weak Hopf algebras. *Journal of Kharkov National University, ser. Nuclei, Particles and Fields*, **521**(2):15-30.
- Duplij, S.A., Li, F., 2001b. Regular solutions of quantum Yang-Baxter equation from weak Hopf algebras. *Czech. J. Phys.*, **51**(12):1306-1311.
- Duplij, S.A., Kotulskaja, O.I., 2002. Supermatrix structures and generalized inverses. *Journal of Kharkov National University, ser. Nuclei, Particles and Fields*, **548**(1):3-14.
- Duplij, S.A., Sadovnikov, A.S., 2002. Regular supermatrix

- solutions for quantum Yang-Baxter equation. *Journal of Kharkov National University, ser. Nuclei, Particles and Fields*, **569**(3):15-22.
- Duplij, S.A., Kalashnikov, V.V., Maslov, E.A., 2005. Quantum information, qubits and quantum algorithms. *Journal of Kharkov National University, ser. Nuclei, Particles and Fields*, **657**(1):99-104.
- Dye, H.A., 2003. Unitary solutions to the Yang-Baxter equation in dimension four. *Quantum Information Processing*, **2**:117-150.
- Etingof, P., Schedler, T., Soloviev, A., 1997. On Set-theoretical Solutions to the Quantum Yang-Baxter Equation. Cambridge, p.4 (Preprint /MIT, q-alg/9707027).
- Etingof, P., Schedler, T., Soloviev, A., 1999. Set-theoretical solutions to the quantum Yang-Baxter equation. *Duke Math. J.*, (2):169-209.
- Faddeev, L.D., Reshetikhin, N.Y., Takhtajan, L.A., 1990. Quantum Lie groups and Lie algebras. *Leningrad Math. J.*, **1**:193-236.
- Gu, P., 1997. A set-theoretical solution of the Yang-Baxter equation and "metahomomorphisms" of groups. *Chinese Sci. Bull.*, **42**(15):1602-1606.
- Hietarinta, J., 1992. All solutions to the constant quantum Yang-Baxter equation in two dimensions. *Phys. Lett.*, **A165**:245-251.
- Hietarinta, J., 1993. Solving the two-dimensional constant quantum Yang-Baxter equation. *J. Math. Phys.*, **34**:1725-1756.
- Hietarinta, J., 1997. Permutation-type solutions to the Yang-Baxter and other n -simplex equations. *J. Phys.*, **A30**:4757-4771.
- Holievo, A.S., 2002. Introduction to Quantum Information Theory. MCNMO, Moskva.
- Kac, V.G., 1977. Lie superalgebras. *Adv. Math.*, **26**(1):8-96.
- Kassel, C., 1995. Quantum Groups. Springer-Verlag, New York, p.531.
- Kauffman, L.H., 1991. Knots and Physics. World Sci., Singapore.
- Kauffman, L.H., Lomonaco, S.J., 2004. Braiding operators are universal quantum gates. *New J. Phys.*, **6**:134-139.
- Khoroshkin, S.M., Tolstoy, V.N., 1991. Universal R -matrix for quantized (super) algebras. *Comm. Math. Phys.*, **141**(3):599-617.
- Kitajev, A., Shen, A., Vialy, M., 1999. Klassi-cheskije i Kvantovyje Vychislienija. Mcnmo, Moskva.
- Lambe, L.A., Radford, D.E., 1997. Introduction to the Quantum Yang-Baxter Equation and Quantum Groups: an Algebraic Approach. Kluwer, Dordrecht, p.292.
- Leites, D.A., 1980. Vvedenije v teoriju supermnogoobrazij. *Uspiehi Mat. Nauk*, **35**(1):3-57.
- Leites, D.A., 1983. A Theory of Supermanifolds. Karelskij filial AN SSSR, Petrozavodsk, p.199.
- Li, F., Duplij, S., 2002. Weak Hopf algebras and singular solutions of quantum Yang-Baxter equation. *Commun. Math. Phys.*, **225**(1):191-217.
- Links, J., Scheunert, M., Gould, M.D., 1994. Diagonalization of the braid generator on unitary irreps of quantum supergroups. *Lett. Math. Phys.*, **32**(3):231-240.
- Lu, J.H., Yan, M., Zhu, Y.C., 2000. On set-theoretical Yang-Baxter equation. *Duke Math. J.*, **104**(1):1-18.
- Majid, S., 1995. Foundations of Quantum Group Theory. Cambridge University Press, Cambridge.
- Manin, U.I., 1984. Kalibrovochnyje Polia I Kompleksnaja Geometrija. Nauka, Moskva, p.335.
- Manin, Y., 1989. Multiparametric quantum deformation of the general linear supergroup. *Comm. Math. Phys.*, **123**:123-135.
- Penrose, R., 1955. A generalized inverse for matrices. *Math. Proc. Cambridge Phil. Soc.*, **51**:406-413.
- Rabin, J.M., 1987. Supermanifolds and Super Riemann Surfaces. Super Field Theories. Plenum Press, New York, p.557-569.
- Rabin, J.M., 1991. Status of the Algebraic Approach to Super Riemann Surfaces. Physics and Geometry. Plenum Press, New York, p.653-668.
- Rao, C.R., Mitra, S.K., 1971. Generalized Inverse of Matrices and Its Application. Wiley, New York, p.251.
- Rogers, A., 1980. A global theory of supermanifolds. *J. Math. Phys.*, **21**(5):1352-1365.
- Shnider, S., Sternberg, S., 1993. Quantum Groups. International Press, Boston, p.371.
- Soloviev, A., 2002. Non-unitary set-theoretical solutions to the quantum Yang-Baxter equation. *Math. Res. Lett.*, **7**:577-596.
- Turaev, V.G., 1994. Quantum Invariants of Knots and 3-Manifolds. W. de Greuter, Berlin.
- Yang, C.N., 1967. Some exact results for the many-body problem in one dimension with repulse delta-function interaction. *Phys. Rev. Lett.*, **19**:1312-1315.
- Zhang, R.B., 1991. Graded representations of the Temperley-Lieb algebra, quantum supergroups, and the Jones polynomial. *J. Math. Phys.*, **32**(10):2605-2613.
- Zhang, R.B., Gould, M.D., 1991. Universal R -matrices and invariants of quantum supergroups. *J. Math. Phys.*, **32**(12):3261-3267.
- Zhang, Y., Kauffman, L.H., Ge, M.L., 2005. Yang-Baxterization, Universal Quantum Gate, and Hamiltonians. Chicago, p.36 (Preprint/Univ. Illinois, quant-ph/0502015).