



Statistical culture-based strategies to enhance chlamyospore production by *Trichoderma harzianum* SH2303 in liquid fermentation^{*}

Ya-qian LI^{†1,2}, Kai SONG¹, Ya-chai LI³, Jie CHEN^{††1,2}

⁽¹⁾School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China)

⁽²⁾Key Laboratory of Urban Agriculture (South), Ministry of Agriculture, Shanghai 200240, China)

⁽³⁾Affiliated Hospital of Hebei University, Baoding 071100, China)

[†]E-mail: lauren@sjtu.edu.cn; jiechen59@sjtu.edu.cn

Received Sept. 17, 2015; Revision accepted Jan. 30, 2016; Crosschecked July 13, 2016

Abstract: *Trichoderma*-based formulations are applied as commercial biocontrol agents for soil-borne plant pathogens. Chlamyospores are active propagules in *Trichoderma* spp., but their production is currently limited due to a lack of optimal liquid fermentation technology. In this study, we explored response surface methodologies for optimizing fermentation technology in *Trichoderma* SH2303. Our initial studies, using the Plackett-Burman design, identified cornmeal, glycerol, and initial pH levels as the most significant factors ($P < 0.05$) for enhancing the production of chlamyospores. Subsequently, we applied the Box-Behnken design to study the interactions between, and optimal levels of, a number of factors in chlamyospore production. These statistically predicted results indicated that the highest number of chlamyospores (3.6×10^8 spores/ml) would be obtained under the following condition: corn flour 62.86 g/L, glycerol 7.54 ml/L, pH 4.17, and 6-d incubation in liquid fermentation. We validated these predicted values via three repeated experiments using the optimal culture and achieved maximum chlamyospores of 4.5×10^8 spores/ml, which approximately a 8-fold increase in the number of chlamyospores produced by *T. harzianum* SH2303 compared with that before optimization. These optimized values could help make chlamyospore production cost-efficient in the future development of novel biocontrol agents.

Key words: *Trichoderma harzianum* SH2303, Chlamyospore, Plackett-Burman screening, Box-Behnken design, Fermentation optimization

<http://dx.doi.org/10.1631/jzus.B1500226>

CLC number: S476

1 Introduction

Trichoderma are fast growing filamentous fungi found in many ecosystems (Marra *et al.*, 2006) that kill other pathogenic fungi and consume them using hydrolytic enzymes (Zeilinger *et al.*, 1999; Harman

et al., 2004). This antagonistic behavior has led to their use (especially the mycoparasites *T. atroviride*, *T. harzianum*, and *T. virens*) as biological control agents (BCAs) for agricultural applications (Schuster and Schmoll, 2010; Carreras-Villasenor *et al.*, 2011). Over the past decade, researchers have made significant progress towards understanding the genetic, developmental, and regulatory signaling elements of the secondary metabolites involved in the interaction of *Trichoderma*-plant-pathogens (Mach and Zeilinger, 2003; Harman, 2006; Woo *et al.*, 2006).

Trichoderma species produce three major types of propagules (mycelia, conidia, and chlamyospores) (Verma *et al.*, 2007) that possess distinct physiological

[‡] Corresponding author

^{*} Project supported by the National Natural Science Foundation of China (Nos. 31201557 and 31270155), the Natural Science Foundation of Shanghai (No. 12ZR1414100), the Foundation of Basic Science and Technology of China (No. 2014FY20900), the Ministry of Education University Doctoral Foundation (No. 20120073120070), and the Shanghai Jiao Tong University Medical-Engineering Cross Research Fund (No. YG2015MS37), China

ORCID: Ya-qian LI, <http://orcid.org/0000-0001-9622-994X>

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2016

characteristics in terms of production, stability, and BCA activity. For efficient BCA action, it is imperative to select the most suitable type of propagule. Most commercial *Trichoderma*-based formulations use conidiospores because they are affordable and can be propagated efficiently in standing liquid or solid form (Fravel, 2005). Chlamydospore-based formulations have other beneficial features, including resistance to drying and low temperatures, insensitivity to soil antibiotics, and extended preservation time, which in turn simplifies processing, storage, and transport of the BCA (Mishra *et al.*, 2012). For instance, studies on the shelf life of the *Trichoderma* chlamydospore-based Tricoguard™ have shown that it can be safely stored for up to 270 d and can survive in the soil of natural ecosystems better than conidia (Jagtap and Bhatnagar, 2000; Mishra *et al.*, 2012). However, a better understanding of the factors involved in the morphogenic switch from mycelia to chlamydospore production is still essential for its use in commercial formulations.

To date, researchers have focused primarily on the effect of environmental conditions, such as carbon (C) and nitrogen (N), C:N ratio, pH, light, and induction signals, on the initiation of conidiation in *Trichoderma* (Gao *et al.*, 2007; Friedl *et al.*, 2008; Steyaert *et al.*, 2010b; Tisch and Schmoll, 2010). A few studies have reported the use of liquid-culture technology for the mass production and application of *Trichoderma* chlamydospores (Lewis and Papavizas, 1983; 1984), but there is still a barrier to their commercial use because of low fermentation yield and high cost. The objective of the present research, therefore, was to develop a cost-effective medium and optimal conditions for the production of chlamydospores in *T. harzianum* SH2303 (effective against corn stalk rot and *Fusarium* Wilt of Cucumber) (Lin *et al.*, 2012) and thus to lay a solid foundation for the development of *Trichoderma*-based active chlamydospore formulations in agriculture.

2 Materials and methods

2.1 Strains and inoculation

We identified and isolated the *T. harzianum* strain SH2303 (CGMCC No. 4963) from corn fields in the Shanghai district based on morphology and internal transcribed spacer (ITS) sequence analysis;

its NCBI accession number is KJ755188. The strain was subsequently grown on potato dextrose agar (PDA) plates, incubated at 28 °C for 5 d, and then the mature spores were harvested and adjusted to 1×10^7 conidia/ml with sterile distilled water. The conidial concentration was determined using a haemocytometer. A total of 100 µl homogeneous conidial suspension (about 1×10^6 conidia) was inoculated into 250 ml Erlenmeyer flasks containing 100 ml of sterilized medium and incubated in a rotary shaker at 28 °C and 200 r/min for 72 h. The initial pH value of the cultures was not controlled.

2.2 Medium and growth conditions

Six different kinds of medium were chosen for the initial screening experiment (Table 1): (1) potato dextrose (PD): potato extract 1000 ml and dextrose (glucose) 20 g/L; (2) potato sucrose (PS): potato extract 1 L and sucrose 20 g/L, pH 6.0–6.5; (3) Richard: sucrose 50 g/L, KNO₃ 10 g/L, K₂HPO₄ 5 g/L, MgSO₄ 0.5 g/L, FeCl₃ 0.02 g/L; (4) Czapek: sucrose 30 g/L, KNO₃ 2 g/L, KH₂PO₄ 1 g/L, FeSO₄·7H₂O 0.5 g/L, KCl 0.5 g/L, FeSO₄ 0.01 g/L; (5) Gorodkova: peptone 10 g/L, beef extract 10 g/L, glucose 2.5 g/L, NaCl 5 g/L; (6) cornmeal medium: cornmeal and water (20:100 (w/v); corn flour 200 g/L). All prepared liquid media were sterilized at 115 °C for 15 min and stored at room temperature until further use. For the potato extracts we sliced the potatoes very thinly, immediately added 1000 ml water to prevent their oxidation, boiled them until soft for 10 min, and finally filtered them through a cotton-cloth.

The pH value of the fermentation media was adjusted by adding either 1 mol/L sodium hydroxide or 1 mol/L hydrochloric acid as required.

2.3 Experimental design and data analysis

The Plackett-Burman (PB) experimental design is a two-level, fractional, factorial design method that is based on the non-perfectly balanced block principle and intended to test for the dependence of some measured quantity on a number of independent variables (Lai *et al.*, 2003). This design filters out the most important factors involved in the fermentation process in as few experiments as possible. We varied each factor ($n=8$) over 2 levels of specialized design (Table 2). The positive and negative effects of the various factors were determined according to the *t*-value test and displayed on a Pareto chart (Fig. 1).




Fig. 1 Pareto chart (a) and normal plot graph (b) identifying the screening factors

Hollow shapes represent the 8 factors of PB test, and among them, orange (F, H, C, G) denotes positive effects; blue (A, B, D, E) denotes negative effects; solid shapes represent 3 blank tests (Note: for interpretation of the references to color in this figure legend, the reader is referred to the web version of this article)

2.4 Box-Behnken design for chlamyospore optimization

The Box-Behnken design (BBD) is a statistical method that is used to study the maximum response corresponding to the best interactions in a multi-factor system (Bae and Shoda, 2005). Each factor, or independent variable, is placed at one of three equal spaced values, usually coded as -1 , 0 , $+1$. The quadratic regression equation was used to calculate the best fit of the main and interaction effects of the various factors.

We studied the optimization, interaction, and combined effects of the three major factors (corn flour, initial pH, and glycerol) on *Trichoderma*-derived chlamyospore production using 17 experimental runs corresponding to the Box-Behnken response surface method. This method was divided into two

categories: (1) the factorial points of the variable values in X_1, X_2, X_3 that constitute the three-dimensional vertex containing a total of 12 factorial points, and (2) zero (the center of the region); the zero experiment was repeated three times to estimate the experimental error. Subsequent to the regression analysis, the experimental factors (R_1) were expressed with the second order polynomial equation, as follows:

$$R_1 = \lambda_0 + \lambda_1 X_1 + \lambda_2 X_2 + \lambda_3 X_3 + \lambda_{12} X_1 X_2 + \lambda_{13} X_1 X_3 + \lambda_{23} X_2 X_3 + \lambda_{11} X_1^2 + \lambda_{22} X_2^2 + \lambda_{33} X_3^2.$$

Optimization of the experimental data identified the model coefficient, while the simple multi-function trait analysis determined the extreme points and the maximum values of the corresponding independent variable.

2.5 Chlamyospore and conidiospore assay

The chlamyospores and conidiospores were assayed with a hemocytometer. Spore suspensions were harvested from 6-d cultures with 5 ml sterile water containing 0.01% Triton X-100. The suspensions were passed through four layers of cheese cloth to remove hyphal fragments, vortexed for 1 min to break up the clumps of cells, and counted microscopically (Olympus BX51TF, Japan). All experiments were carried out three times.

2.6 Statistical analysis

A logarithmic transformation was applied on the sporulation data before statistical analysis. Statistical analysis of all data was subjected to Design Expert 7.0 software (Stat-Ease, Inc., Minneapolis, MN, USA). The statistical significance of the regression coefficients was 95%. The optimum levels of variables were obtained by graphical analysis.

3 Results

3.1 Effect of the C:N ratio on chlamyospore formation

Five kinds of medium with different concentrations of C and N were chosen to identify the optimal C:N ratio (Table 1). Results showed that conidiospores, but not chlamyospores, were produced on the Richard medium (C:N=17.5:1) while the Gorodkova

Table 1 Effects of different media on chlamyospore and conidiospore production by *T. harzianum* SH2303 in liquid culture

No.	Medium	Main C/N source (g/L)	C:N (mol:mol)	ChlS ($\times 10^6$)	ConS ($\times 10^6$)
1	PD	Glucose (20)	(3–4):1	0.27 \pm 0.10	4.20 \pm 0.32
2	PS	Sucrose (20)	(3–4):1	2.20 \pm 0.13	2.40 \pm 0.21
3	Richard	Sucrose (50), KNO ₃ (10)	17.5:1	0	3.40 \pm 0.20
4	Czapek	Sucrose (30), KNO ₃ (2)	52.5:1	0.62 \pm 0.05	1.20 \pm 0.18
5	Gorodkova	Glucose (2.5), peptone (10), beef extract (10)	1:2	2.50 \pm 0.18	0
6	Cornmeal	Corn flour (200)	9:1	52.00 \pm 5.00	61.00 \pm 2.20

ChlS: chlamyospores; ConS: conidiospores; PD: potato dextrose; PS: potato sucrose. Carbon and nitrogen content in natural and synthetic ingredients. Peptone (12% nitrogen), beef extract (13% nitrogen), corn flour (8.7% nitrogen and 75.2% carbon)

medium (C:N=1:2) was able to produce chlamyospores only. Similarly, higher levels of glucose (PD) compared to sucrose (PS) promote conidiospore production. It is worth noting that the natural material cornmeal (C:N=9:1) is highly beneficial for the formation of both spores. Compared with synthetic media, cornmeal contains a variety of amino acids, trace elements, vitamins, and some unknown growth factors that are advantageous for the production of chlamyospores. As corn flour is cheap and easily acquired, we adopted corn flour as providing the optimal C and N for enhanced chlamyospore production.

3.2 Plackett-Burman screen of crucial factors

We also analyzed the following factors as possible candidates for improving chlamyospore formation: peptone, glycerol, (NH)₄SO₄, KH₂PO₄, MgSO₄·7H₂O, plus the inoculation volume and the initial pH of the medium. In Table 2, each row represents an experiment and each column represents an independent variable while the signs represent the two different levels (high and low) of the independent variable. The significant levels of each medium were determined by *t*-test; the predicted value was consistent with the real values as identified on the fitted first-order model (Table 3).

Further analysis, analysis of variance (ANOVA), revealed that cornmeal, initial pH, and glycerol all had a significant positive influence on the formation of chlamyospores. Other factors such as inoculums and peptone were found to have negative effects. The calculated model *F*-value of 72.27 implies that the model is significant; there is only a 1.37% chance that such a large value could occur due to noise. Values of “Prob>*F*” that are less than 0.01 indicate that the model terms are extremely significant. In this case *F*, *H*, and *C* are significant (Table 4). The calculated

Table 2 Code and actual values of all 8 variables used in the Plackett-Burman design

Variable	Culture factor	Code level	
		Low (-1)	High (+1)
<i>A</i>	Inoculums (ml)	2	8
<i>B</i>	Peptone (g/L)	0	4
<i>C</i>	Initial PH	3	7
<i>D</i>	(NH) ₄ SO ₄ (g/L)	0	1
<i>E</i>	KH ₂ PO ₄ (g/L)	0	3
<i>F</i>	Corn flour (g/L)	40	80
<i>G</i>	MgSO ₄ ·7H ₂ O (g/L)	0	3
<i>H</i>	Glycerol (g/L)	0	5

Table 3 Experimental design of the Plackett-Burman method used for screening the culture

Run	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>	<i>F</i>	<i>G</i>	<i>H</i>	<i>Y</i>	\hat{Y}
1	+1	-1	-1	-1	+1	-1	+1	+1	6.62	6.59
2	+1	-1	+1	+1	+1	-1	-1	-1	6.32	6.30
3	-1	+1	+1	-1	+1	+1	+1	-1	7.08	7.05
4	-1	+1	-1	+1	+1	-1	+1	+1	6.45	6.48
5	-1	-1	-1	-1	-1	-1	-1	-1	6.56	6.57
6	-1	-1	+1	-1	+1	+1	-1	+1	7.49	7.52
7	+1	+1	-1	+1	+1	+1	-1	-1	6.08	6.10
8	+1	+1	+1	-1	-1	-1	+1	-1	6.38	6.41
9	+1	-1	+1	+1	-1	+1	+1	+1	7.38	7.39
10	+1	+1	-1	-1	-1	+1	-1	+1	6.87	6.86
11	-1	-1	-1	+1	-1	+1	+1	-1	6.92	6.91
12	-1	+1	+1	+1	-1	-1	-1	+1	6.79	6.76

A–H correspond to the different factors shown in Table 2, i.e., inoculums, peptone, initial pH, (NH)₄SO₄, KH₂PO₄, corn flour, MgSO₄·7H₂O, and glycerol in alphabetical order. *Y*, experimental responses; \hat{Y} , predicted values calculated from the fitted first-order model

“Pred *R*-Squared” of 0.8552 is in reasonable agreement with the “Adj *R*-Squared” of 0.9668. The “Adeq Precision” measures the signal to noise ratio; a ratio greater than 4 is desirable while the calculated ratio of 21.101 indicates an adequate signal. This model can be used to navigate the design space.

Table 4 Analysis of variance for the Plackett-Burman design

Factor	Sum of squares	df	Mean square	F-value	P-value (Prob>F)
Model	2.005533	8	0.222837	72.27147	0.0137*
A-A	0.224133	1	0.224133	72.69189	0.0135*
B-B	0.224133	1	0.224133	72.69189	0.0135*
C-C	0.313633	1	0.313633	101.71890	0.0097**
D-D	0.093633	1	0.093633	30.36757	0.0314*
E-E	0.061633	1	0.061633	19.98919	0.0466*
F-F	0.607500	1	0.607500	197.02700	0.0050**
G-G	0.043200	1	0.043200	14.01081	0.0645
H-H	0.425633	1	0.425633	138.04320	0.0072**
Residual	0.006167	2	0.003083		
Cor total	2.011700	10			

** P<0.01, * P<0.05

3.3 Box-Behnken design optimization

We next optimized the production of *T. harzi-anum* chlamydo spores using corn flour (X_1), initial pH (X_2), and glycerol (X_3) as the crucial parameters in our Box-Behnken analysis (Table 5). The experimental results were fitted with a second order polynomial equation. The values of the regression coefficients were calculated and the fitted equations (in terms of the coded values) for predicting log chlamydo spores (Y_C) were as follows:

$$Y_C = 8.61 + 0.08X_1 - 0.00625X_2 - 0.054X_3 + 0.020X_1X_2 - 0.030X_1X_3 + 0.0025X_2X_3 - 0.097X_1^2 - 0.084X_2^2 - 0.074X_3^2.$$

Based on our results, the “Pred R-Squared” of 0.9172 is in reasonable agreement with the “Adj R-Squared” of 0.9494 and the “Adeq Precision” of 15.391 indicates an adequate signal. The analysis of variance for the refined model is summarized in Table 6. The model F-value of 34.35 implies that the model was significant; there is only a 0.01% chance that this value could occur due to noise. Values of “Prob>F” less than 0.05 indicate that the model terms are significant. For our experiments, X_1 (cornmeal) and X_3 (initial pH) were both significant. The calculated “Lack of Fit F-value” of 0.24 implies that it is not significantly relative to pure error. We found that there is a 86.69% chance that this value could

Table 5 Results of the central composition experiment

Run	Code variable level			Real variable level			log Chl	
	X_1	X_2	X_3	Corn flour	Initial pH	Glycerol	Y	\hat{Y}
1	1	1	0	65	4.0	9.0	8.43	8.37
2	0	0	0	60	4.0	8.0	8.59	8.49
3	1	0	-1	65	3.8	8.0	8.49	8.32
4	0	1	-1	60	3.8	9.0	8.40	8.52
5	-1	-1	0	55	4.0	7.0	8.38	8.38
6	0	0	0	60	4.0	8.0	8.65	8.60
7	0	0	0	60	4.0	8.0	8.57	8.33
8	0	-1	-1	60	3.8	7.0	8.52	8.43
9	-1	0	-1	55	3.8	8.0	8.36	8.51
10	-1	0	1	55	4.2	8.0	8.32	8.49
11	0	-1	1	60	4.2	7.0	8.49	8.40
12	0	1	1	60	4.2	9.0	8.38	8.39
13	0	0	0	60	4.0	8.0	8.61	8.61
14	1	0	1	65	4.2	8.0	8.53	8.61
15	1	-1	0	65	4.0	7.0	8.59	8.61
16	0	0	0	60	4.0	8.0	8.61	8.61
17	-1	1	0	55	4.0	9.0	8.34	8.61

Y, experimental responses; \hat{Y} , predicted values calculated from the fitted second-order model; Chl: chlamydo spore

Table 6 ANOVA analysis of the regression equation

Factor	Sum of squares	df	Mean square	F-value	P-value (Prob>F)
Model	0.180000	9	0.020000	34.35000	<0.0001**
X_1	0.051200	1	0.051200	86.46562	<0.0001**
X_2	0.000312	1	0.000312	0.52774	0.4911
X_3	0.023113	1	0.023113	39.03197	0.0004**
X_1X_2	0.001600	1	0.001600	2.70205	0.1442
X_1X_3	0.003600	1	0.003600	6.07961	0.0431*
X_2X_3	0.000025	1	0.000025	0.04222	0.8430
X_1^2	0.039413	1	0.039413	66.55977	<0.0001**
X_2^2	0.029887	1	0.029887	50.47191	0.0002**
X_3^2	0.023213	1	0.023213	39.20151	0.0004**
Residual	0.004145	7	0.000592		
Lack of fit	0.000625	3	0.000208	0.23674	0.8669
Pure error	0.003520	4	0.000880		
Cor total	0.187212	16			

** P<0.01, * P<0.05

occur due to noise. These results are consistent with the model. Chlamydo spore production, as predicted by the final quadratic model together with the corresponding observed values, is shown in Table 5. Results show excellent agreement between the predicted value (\hat{Y}) and experimental data (Y). The optimum

values for achieving the maximum level of chlamyospore production are $X_1=62.86$ g/L, $X_2=4.17$, and $X_3=7.54$ ml/L, and the predicted chlamyospore production corresponding to these values is 3.6×10^8 spores/ml. This concentration is the maximum value bounded by the experimental values.

These experiments were performed in triplicate to confirm the accuracy of the model, yielding an average of 4.5×10^8 spores/ml. The agreement between the predicted and experimental results verifies the validity of the model and the existence of optimal conditions. For response contour plots (Fig. 2), based on the final model, we held two variables constant at their optimum levels while varying the other two within their experimental range. As shown in Fig. 2a, the maximum response occurs when cornmeal and initial pH are at approximately 62.86 g/L and 4.17, respectively. Moreover, these two factors were found to have synergistic effects on chlamyospore production. A 3D surface graph representation of the data corresponding to the contour plot of the predicted values at the optimal concentrations of corn flour and glycerol versus their desirability can be seen in Fig. 3.

4 Discussion

Trichoderma has the potential to produce chlamyospores cost-effectively and maintain long periods of vigorous vegetative growth during use (Fravel, 2005). We optimized statistically the fermentation conditions for the production of *Trichoderma* SH2303-derived chlamyospores in flask experiments using cornmeal as the basal medium. We also used a two-level Plackett-Burman and an optimal Box-Behnken design to determine the important factors and optimal concentrations for chlamyospore production. We were able to achieve maximum production of 4.5×10^8 chlamyospores/ml using an optimal medium consisting of cornmeal (62.86 g/L), glycerol (7.54 ml/L), and an initial pH of 4.17, in liquid culture. This is the first publication describing how to optimize the production of *T. harzaium*-derived chlamyospores using a Box-Behnken design with cornmeal as the primary nutrient.

Biotic factors (especially carbon and nitrogen status) influence conidiospore production in many species of *Trichoderma* (Gao et al., 2007). We found




Fig. 2 Contour plots of the interaction effect of two specific factors on chlamyospore production

(a) Effects of corn meal (X_1) and pH (X_2), and their mutual interaction; (b) Effects of corn meal (X_1) and glycerol (X_3), and their mutual interaction; (c) Effects of glycerol (X_3) and pH (X_2), and their mutual interaction




Fig. 3 3D surface graph and its corresponding contour plot of prediction of chlamydospore production (b) and its desirability (a) at the optimal concentrations of corn flour (X_1) and glycerol (X_3)

that the production of chlamydospores (thick-walled mitotic spores that generally act as “resting spores”) also relies heavily on different levels of nutrient (carbon and nitrogen) and environmental conditions (pH). Natural carbon and nitrogen sources, such as corn flour, were shown to benefit preferentially sporulation of both chlamydospores and conidiospores. These data disagree with a previous report which found that *T. viride*-derived conidiation is primarily dependent on the level of the carbon (Gao *et al.*, 2007). Our data indicate that C:N ratio affects the production of *T. harzianum* SH2303-derived spores, while natural corn meal seems to maximize sporulation. The influence of C:N ratio on sporulation is strain-specific; however, hence consideration must be given to the complexity of nutrient requirements

for improving spore yields for fungal biocontrol agents.

Several growth/production media using molasses, D-glucose, cellulose, and/or soluble starch have been shown to increase the production of *Trichoderma* spp. spores (Lewis and Papavizas, 1983). However, these substrates are not economically efficient, owing to high raw material costs and moderate sporulation (10^8 spores). In the current study, cornmeal, an important natural source of carbon and nitrogen with a relatively low cost, was shown to promote significantly the formation of SH2303-derived chlamydospores in liquid fermentation. Several other studies have reported on the use of alternative cheap raw materials for *Trichoderma* biomass production, including bran, and peanut hull meal (Singh *et al.*, 2007; Maurya *et al.*, 2012).

Glycerol is a rapid carbon source that benefits mycelial growth during the early stages of fermentation and has been found to increase the capacity of *T. harzianum* talc formulations to retain water, thus increasing their longevity (Sriram *et al.*, 2011). Other environmental parameters such as light, temperature, and especially pH are also critical for chlamydospore production. Although conidial fungi can grow over a wide range of pH, maximum growth and sporulation occur near neutral pH. Our study clearly demonstrates that an acidic pH (4.17) maximizes chlamydospore formation. Similarly, a low ambient pH has been shown to favor conidiation in *Trichoderma* (Moreno-Mateos *et al.*, 2007) and photoconidiation in *T. pleuroticola*, *T. atroviride*, and *T. hamatum*, suggesting that photoconidiation may also be regulated in the different species (Steyaert *et al.*, 2010a). Recent research into the molecular regulation of conidiation by VELVET in *T. virens*, suggests that VeA acts as a positive regulator of conidiation and chlamydospore production in a nutrient-rich liquid media, while the deletion of *vell* strongly promotes chlamydospore production (Mukherjee and Kenerley, 2010). These data suggest how *vell* gene-mediated signaling may be manipulated to improve the production of chlamydospores.

We found that the efficient and effective use of *Trichoderma* to produce BCAs involves achieving a balance between total cost and the overall growth rate of the chlamydospores in a liquid medium. An understanding of liquid growth culture conditions will

also be useful for future genome-wide analyses of gene expression changes during chlamyospore development. Future studies on other parameters, such as dissolved oxygen and carbon dioxide, and construction of *vel1* over-expression strain, will further help maximize the production of chlamyospore-based BCAs during fermentation.

Compliance with ethics guidelines

Ya-qian LI, Kai SONG, Ya-chai LI, and Jie CHEN declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Bae, S., Shoda, M., 2005. Statistical optimization of culture conditions for bacterial cellulose production using Box-Behnken design. *Biotechnol. Bioeng.*, **90**(1):20-28. <http://dx.doi.org/10.1002/bit.20325>
- Carreras-Villasenor, N., Sanchez-Arreguin, J.A., Herrera-Estrella, A.H., 2011. *Trichoderma*: sensing the environment for survival and dispersal. *Microbiology*, **158**(1):3-16. <http://dx.doi.org/10.1099/mic.0.052688-0>
- Fravel, D.R., 2005. Commercialization and implementation of biocontrol. *Annu. Rev. Phytopathol.*, **43**(1):337-359. <http://dx.doi.org/10.1146/annurev.phyto.43.032904.092924>
- Friedl, M.A., Kubicek, C.P., Druzhinina, I.S., 2008. Carbon source dependence and photostimulation of conidiation in *Hypocrea atroviridis*. *Appl. Environ. Microbiol.*, **74**(1):245-250. <http://dx.doi.org/10.1128/AEM.02068-07>
- Gao, L., Sun, M.H., Liu, X.Z., et al., 2007. Effects of carbon concentration and carbon to nitrogen ratio on the growth and sporulation of several biocontrol fungi. *Mycol. Res.*, **111**(1):87-92. <http://dx.doi.org/10.1016/j.mycres.2006.07.019>
- Harman, G.E., 2006. Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*, **96**(2):190-194. <http://dx.doi.org/10.1094/PHTO-96-0190>
- Harman, G.E., Howell, C.R., Viterbo, A., et al., 2004. *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nat. Rev. Microbiol.*, **2**(1):43-56. <http://dx.doi.org/10.1038/nrmicro797>
- Jagtap, G.P., Bhatnagar, S.S., 2000. *Trichoderma* chlamyospores-based formulation (Tricoguard™): impact on shelf life. *Pestology*, **24**(9):70-71.
- Lai, L.S.T., Pan, C.C., Tzeng, B.K., 2003. The influence of medium design on lovastatin production and pellet formation with a high-producing mutant of *Aspergillus terreus* in submerged cultures. *Process Biochem.*, **38**(9):1317-1326. [http://dx.doi.org/10.1016/S0032-9592\(02\)00330-8](http://dx.doi.org/10.1016/S0032-9592(02)00330-8)
- Lewis, J.A., Papavizas, G.C., 1983. Production of chlamyospores and conidia by *Trichoderma* spp. in liquid and solid growth media. *Soil. Biol. Biochem.*, **15**(3):351-357. [http://dx.doi.org/10.1016/0038-0717\(83\)90083-4](http://dx.doi.org/10.1016/0038-0717(83)90083-4)
- Lewis, J.A., Papavizas, G.C., 1984. Chlamyospore formation by *Trichoderma* spp. in natural substrates. *Can. J. Microbiol.*, **30**(1):1-7. <http://dx.doi.org/10.1139/m84-001>
- Lin, Z., Ma, J., Yi, S., et al., 2012. Effect of salinity on the biocontrol *Trichoderma* strain SH2303. *J. Shanghai Jiaotong Univ. (Agric. Sci.)*, **30**(5):51-53 (in Chinese).
- Mach, R.L., Zeilinger, S., 2003. Regulation of gene expression in industrial fungi: *Trichoderma*. *Appl. Microbiol. Biotechnol.*, **60**(5):515-522. <http://dx.doi.org/10.1007/s00253-002-1162-x>
- Marra, R., Ambrosino, P., Carbone, V., et al., 2006. Study of the three-way interaction between *Trichoderma atroviride*, plant and fungal pathogens by using a proteomic approach. *Curr. Genet.*, **50**(5):307-321. <http://dx.doi.org/10.1007/s00294-006-0091-0>
- Maurya, D.P., Singh, D., Pratap, D., et al., 2012. Optimization of solid state fermentation conditions for the production of cellulase by *Trichoderma reesei*. *J. Environ. Biol.*, **33**(1):5-8.
- Mishra, D.S., Prajapati, C.R., Gupta, A.K., et al., 2012. Relative bio-efficacy and shelf-life of mycelial fragments, conidia and chlamyospores of *Trichoderma harzianum*. *Vegetos*, **25**(1):225-232.
- Moreno-Mateos, M.A., Delgado-Jarana, J., Codón, A.C., et al., 2007. pH and *Pac1* control development and antifungal activity in *Trichoderma harzianum*. *Fungal Genet. Biol.*, **44**(12):1355-1367. <http://dx.doi.org/10.1016/j.fgb.2007.07.012>
- Mukherjee, P.K., Kenerley, C.M., 2010. Regulation of morphogenesis and biocontrol properties in *Trichoderma virens* by a VELVET protein, Vel1. *Appl. Environ. Microbiol.*, **76**(7):2345-2352. <http://dx.doi.org/10.1128/AEM.02391-09>
- Schuster, A., Schmoll, M., 2010. Biology and biotechnology of *Trichoderma*. *Appl. Microbiol. Biotechnol.*, **87**(3):787-799. <http://dx.doi.org/10.1007/s00253-010-2632-1>
- Singh, A., Srivastava, S., Singh, H.B., 2007. Effect of substrates on growth and shelf life of *Trichoderma harzianum* and its use in biocontrol of diseases. *Bioresour. Technol.*, **98**(2):470-473. <http://dx.doi.org/10.1016/j.biortech.2006.01.002>
- Sriram, S., Roopa, K.P., Savitha, M.J., 2011. Extended shelf-life of liquid fermentation derived talc formulations of *Trichoderma harzianum* with the addition of glycerol in the production medium. *Crop Prot.*, **30**(10):1334-1339. <http://dx.doi.org/10.1016/j.cropro.2011.06.003>
- Steyaert, J.M., Weld, R.J., Stewart, A., 2010a. Ambient pH intrinsically influences *Trichoderma* conidiation and colony morphology. *Fungal Biol.*, **114**(2-3):198-208. <http://dx.doi.org/10.1016/j.funbio.2009.12.004>
- Steyaert, J.M., Weld, R.J., Stewart, A., 2010b. Isolate-specific conidiation in *Trichoderma* in response to different

- nitrogen sources. *Fungal Biol.*, **114**(2-3):179-188.
<http://dx.doi.org/10.1016/j.funbio.2009.12.002>
- Tisch, D., Schmoll, M., 2010. Light regulation of metabolic pathways in fungi. *Appl. Microbiol. Biotechnol.*, **85**(5): 1259-1277.
<http://dx.doi.org/10.1007/s00253-009-2320-1>
- Verma, M., Brar, S.K., Tyagi, R.D., et al., 2007. Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochem. Eng. J.*, **37**(1):1-20.
<http://dx.doi.org/10.1016/j.bej.2007.05.012>
- Woo, S.L., Scala, F., Ruocco, M., et al., 2006. The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology*, **96**(2): 181-185.
<http://dx.doi.org/10.1094/PHTO-96-0181>
- Zeilinger, S., Galhaup, C., Payer, K., et al., 1999. Chitinase gene expression during mycoparasitic interaction of *Trichoderma harzianum* with its host. *Fungal Genet. Biol.*, **26**(2):131-140.
<http://dx.doi.org/10.1006/fgbi.1998.1111>

中文概要

题目: 哈茨木霉菌 SH2303 高产厚垣孢子液体发酵优化
目的: 鉴于厚垣孢子为主要活性成分的木霉菌剂具有抗逆性强、存活期长、易贮藏等特点, 更容易满足实际生产需求, 本文利用系统的液体发酵策略, 优化木霉菌高产厚垣孢子液体发酵条件,

同时降低发酵成本, 指导未来木霉菌厚垣孢子生防菌剂研发。

创新点: 采用从开放到闭合的递进式液体发酵筛选优化模式, 通过 Plackett-Burman 设计和 Box-Behnken 优化设计确定影响木霉菌产厚垣孢子的关键因子及最佳发酵条件, 确定玉米粉作为基础培养基, 添加甘油和调节 pH 相结合的液体发酵策略, 在降低发酵成本的同时提高厚垣孢子产量。

方法: 以玉米粉为基础培养基, 通过 Plackett-Burman 筛选实验, 从 8 个候选的发酵因子中初步确定玉米粉、甘油和发酵液起始 pH 值三个显著因子。进而利用响应面中的 Box-Behnken 设计 (表 5) 三个因素为自变量, 厚垣孢子产量为响应值, 通过 Design Expert 7.0 软件对实验数据进行多项式回归分析, 建立多元二次回归方程。通过方差分析, 最终确定木霉菌 SH2303 高产厚垣孢子液体发酵工艺的最佳参数。

结论: 确定了影响厚垣孢子形成的三个关键的发酵因子: 玉米粉、甘油和起始 pH 值。获得厚垣孢子形成最佳的液体发酵条件: 玉米粉 62.86 g/L, 甘油 7.54 ml/L, 起始 pH 4.17, 发酵 6 天。实际最大厚垣孢子产量 4.5×10^8 孢子/ml, 与模型预测值 3.6×10^8 孢子/ml 相当。厚垣孢子产量与优化前相比提高 8 倍, 可用于指导低成本的木霉菌厚垣孢子新型生防菌剂开发。

关键词: 哈茨木霉菌 SH2303 厚垣孢子; Plackett-Burman 筛选; Box-Behnken 设计; 液体发酵优化