



Review

<https://doi.org/10.1631/jzus.B2300779>

Advancements in mechanisms and drug treatments for fibrodysplasia ossificans progressiva

Yijun ZHOU¹, Ce SHI²✉, Hongchen SUN²✉

¹Department of Endodontics, Hospital of Stomatology, Jilin University, Changchun, 130021, China

²Department of Oral Pathology, Hospital of Stomatology, Jilin University, Changchun, 130021, China

Abstract: Fibrodysplasia ossificans progressiva (FOP) is a rare genetic disorder characterized by congenital bilateral malformation of the large toe and progressive, extensive, and irreversible heterotopic ossification (HO) of soft tissues throughout the body, leading to severe disabilities. FOP is caused primarily by mutations in activin A receptor type I (ACVR1), also known as activin-like kinase 2 (ALK2), a receptor belonging to the bone morphogenetic protein (BMP) type I family. However, the continuous and complex process of HO in FOP is not yet fully understood, which has impeded the development of therapeutic drugs. Despite surgical removal of HO, which often results in recurrence and expansion of ossification, there is currently no definitive drug treatment available to completely prevent, halt, or reverse the progression of HO in FOP. Currently, researchers are intensively studying the pathogenesis of FOP at various stages and developing promising drug candidates, including Saracatinib, Palovarotene, and Rapamycin. This review provides an overview of progress in understanding the mechanism of FOP and the development of therapeutic drugs, with the goal of providing insights for further research and the development of new treatment methods.

Key words: Fibrodysplasia ossificans progressive (FOP); Heterotopic ossification (HO); Mechanism; Drug treatment; Activin A receptor type 1 (ACVR1)

1 Introduction

Fibrodysplasia ossificans progressiva (FOP, OMIM135100) is a rare genetic heterotopic ossification (HO) disease (Towler and Shore, 2022) with a prevalence of about 1 in 2 million individuals in the Western world (She and Zhang, 2018). It is characterized by congenital bilateral malformation of the large toe, presenting as short, thick, and valgus, and extensive, progressive, and irreversible HO of the soft tissues throughout the body. Even minor local inflammation, such as vaccination, can cause HO. Progressive HO leads to trunk deformities and joint contractures in the extremities, and systemic ankyloses can result in walking difficulties and respiratory dysfunction in some patients (Kitoh, 2020), significantly impairing both their quality of life and life expectancy. The life expectancy of patients is generally only about 40 years (Pignolo et al., 2022a).

Currently, there is no definitive drug treatment available to completely prevent, halt, or reverse HO in FOP, and trauma prevention remains a primary therapeutic focus (Kitoh, 2020). Glucocorticoids, non-steroidal anti-inflammatory drugs (NSAIDs), leukotriene inhibitors, and mast cell stabilizers are used to control inflam-

✉ Hongchen SUN, hcsun@jlu.edu.cn;

Ce SHI, ceshi@jlu.edu.cn

Yijun ZHOU, <https://orcid.org/0009-0002-4035-8157>

Hongchen SUN, <https://orcid.org/0000-0002-2533-8150>

Ce SHI, <https://orcid.org/0000-0002-1033-1626>

Received Oct. 29, 2023; Revision accepted Jan. 12, 2024;

Crosschecked xxx. xx, 20xx; Published online xxx. xx, 20xx

mation (Brennan et al., 2018; Wentworth et al., 2019). However, the effect of current drug therapy is limited, leading to recurrence and expansion of ossification after surgical removal of HO. Moreover, the three main current requirements for clinical medicine, namely (1) drugs that can be safely used to prevent HO on a daily basis, (2) managing pain, and (3) removing/reducing the HO that has already formed to relieve fused joints, are not being adequately met.

Research has shown that all reported FOP cases show mutations in the activin A receptor type I (ACVR1) gene and 95% show the R206H mutation (Shore et al., 2006), resulting in ACVR1 becoming an activin receptor and activating the downstream SMAD1/5/9 signaling pathway (Hatsell et al., 2015). However, HO in FOP is a complex process. This review provides an overview of the researches focused on the pathogenesis of FOP and the development of promising drug candidates in order to provide reference for further research.

2 Mutation of the ACVR1 gene and changes in TGF- β /BMP signaling pathways

2.1 Mechanisms

FOP is caused by mutations in the *ACVR1* gene at 2q23-24 (Shore, et al., 2006). ACVR1 is a type I BMP transmembrane serine/threonine kinase receptor. Type I BMP receptors, which include ACVR1 (ALK2), BMPRIA (ALK3), and BMPRIB (ALK6), can form heterotetramers with type II BMP receptors such as BMPRII, ACVR2A, and ACVR2B upon BMP binding (Hu et al., 2021). Type II BMP receptors subsequently phosphorylate the glycine/serine region of type I BMP receptors, triggering downstream signaling pathways (Jia and Meng, 2021). The R206H mutation in ACVR1 results in its aberrant function as an activin receptor, leading to perpetual activation of the downstream SMAD1/5/9 signaling pathway (Hatsell, et al., 2015). This mutation also reduces the binding affinity of ACVR1 to its negative regulator FKBP12, thereby promoting the onset and progression of HO in FOP (Song et al., 2010; Hatsell, et al., 2015).

Activin A (ACTA) is a secreted factor implicated in the development of HO in FOP. Experiments involving *Acvr1*^{[R206H]^{FlEx}+} transgenic mice as an FOP model have shown that these mice do not develop HO either spontaneously or in response to local injury once the activation of ACVR1 by ACTA is blocked (Hatsell, et al., 2015). While some studies have reported increased ACTA production in fibroblasts from FOP patients (De Rooter et al., 2022), others have found no significant differences in serum ACTA levels between FOP patients and healthy controls (Ye et al., 2023). This discrepancy suggests a potential tissue/cell-specific effect of ACTA that warrants further investigation. ACTA has been shown to either engage ACVR1B to activate Smad2/3 or bind ACVR1 and type II Activin/BMP receptors via its finger 2 tip loop to form a non-signaling complex (NSC) in the BMP signaling pathway. FOP-associated ACVR1 missense mutations can convert this NSC into a signaling complex. The NSC containing wild-type ACVR1 exerts a negative regulatory effect on the signaling that drives HO in FOP (Aykul et al., 2020), and overexpression of wild-type ACVR1 in FOP mice inhibits HO (Yamamoto et al., 2022). Overexpression of ACVR1 in fibro/adipogenic progenitors (FAPs) has a cell-autonomous effect, reducing ACTA-induced osteogenic signaling (Yamamoto, et al., 2022). These findings underscore a regulatory mechanism of the levels of BMP signaling abnormality induced by ACTA with the ACVR1 R206H mutation.

In addition to ACVR1 and ACTA, other molecules in the BMP pathway have attracted research interest. For instance, in zebrafish, Wentworth et al. (2022) found that the knockdown of wild-type *BMPRIA* and *ACVR2A* could weaken ACVR1 R206H signaling, particularly in response to ACTA. FOP also implicates signaling cascades within the TGF- β superfamily. Notably, TGF- β /BMP family ligands can also transmit signals via non-SMAD cascades, such as TGF- β -activated kinase 1 (TAK1) (Zhang, 2017; Hwang et al., 2022). In FOP patients, macrophages have been observed to secrete increased amounts of TGF- β and exhibit prolonged NF- κ B and p38 MAPK activation without p-SMAD1/5 changes, indicating dysregulated TAK1 activation (Barruet et al., 2018). Recent studies have shown that in primary dermal fibroblasts from FOP patients, TGF- β 1

acted as an FOP-specific stimulant of ACTA, as shown by the upregulation of INHBA expression (De Ruiter et al., 2023). Therefore, TGF- β signaling pathways may play a previously unexpected role in FOP, and we expect further insights from ongoing research.

2.2 Drug Treatments

2.2.1 Gene Therapy

Gene therapy, a promising avenue for diseases with genetic origins such as inherited retinal dystrophy (Chiu et al., 2021), is similarly being explored for FOP. Several gene therapy approaches are under investigation for the treatment of FOP due to the highly consistent genetic etiology, the severe lifetime progression of ectopic ossification, and the heavy burden of lifetime medication.

2.2.1.1 Si-RNA

Kaplan et al. (2012) used allele-specific RNAi (ASP-RNAi) to suppress disease-related alleles without inhibiting the normal alleles and reported that ASP-RNAi could mediate selective suppression of the mutant c.617A *ACVR1*, decrease the increased BMP pathway signaling, and restore enhanced osteogenic differentiation of FOP SHED cells Yang et al. (2022) used a combined approach of gene replacement and silencing (by an AAV-compatible artificial miRNA) to suppress ACTA-induced aberrant BMP signaling of human FOP iPSCs and mouse *ACVR1*(R206H) KI skeletal cells. This approach prevented both traumatic and spontaneous HO in *Acvr1* (R206H)KI FOP mice without causing detrimental effects on cartilage development, bone growth, or bone remodeling (Yang et al., 2022). They also tried AAV combinations targeting *ACVR1* and *ACTA*, and the local delivery of this AAV gene therapy resulted in a significant decrease in endochondral bone formation in *Acvr1*R206H/+ mice (Yang et al., 2023). We expect further insights concerning optimization of the combination, vector biodistribution, toxicity, dose-ranging, and especially side effects.

2.2.1.2 Antisense oligonucleotides (ASOs)

ASOs are synthetic single-stranded DNA or RNA sequences, typically 15-25 nucleotides in length, that downregulate gene expression by inducing RNaseH-mediated degradation of the target transcript (gapmers) or preventing the binding of proteins to RNA (steric-blocking) (Lim et al., 2020). Recently, the Food and Drug Administration (FDA) approved nusinersen, an ASO, for the treatment of spinal muscular atrophy (Nagar et al., 2023).

Shi et al. (2013) designed ASOs targeting exon 8 of the wild-type mouse *Acvr1* gene. These ASOs knocked down mouse *Acvr1* expression through exon skipping, promoting myogenic differentiation of C2C12 myoblasts and inhibiting BMP6-induced osteoblast differentiation in mouse endothelial cells.

Maruyama and Yokota (2018) created a phosphorodiamidate morpholino oligomer (PMO), a morpholine moiety connected through methylene phosphorodiamidate that replaces the ribose, and successfully skipped the FOP mutation (exon 6) of *ACVR1* in FOP patient fibroblasts

Building on ASO technology, locked nucleic acids (LNAs) offer increased stability by connecting the 2'-oxygen with the 4'-carbon on the same ribose through a methylene bridge (Xiong et al., 2021). Maruyama et al. (2022) developed LNA gapmers, short DNA oligonucleotides with LNA modifications at both ends, which efficiently knocked down *ACVR1* R206H expression at the RNA level, while *ACVR1* WT was mainly unaffected in human FOP fibroblasts. Furthermore, some LNA gapmers suppressed ACTA-induced osteogenic differentiation in C2C12 (*ACVR1* R206H) cells.

While this strategy shows promise for modulating the activity of the *ACVR1* gene, its efficacy in FOP mouse models remains to be studied.

2.2.2 Inhibitors of Type I BMP receptor/*ACVR1*

A common strategy in developing FOP drugs is to inhibit the function of *ACVR1* or the BMP Type I receptor. The first inhibitor discovered was Dorsomorphin, identified during in vivo screening of BMP inhibitors (Yu et al., 2008b). Dorsomorphin exerts its inhibitory effect on *ACVR1* by directly binding to the ATP-binding

pocket of ACVR1's intracellular kinase domain through its pyrazolo[1,5-a]pyrimidine core (Chaikuad et al., 2012). However, concerns regarding its clinical safety arose due to the poor specificity of Dorsomorphin (Luo et al., 2016). Subsequently, more selective inhibitors, namely DMH1 (Hao et al., 2010) and LDN-193189 (Cuny et al., 2008; Yu et al., 2008a), were developed and shown to be effective in FOP animal models. To better distinguish ACVR1 (ALK2) from other BMP type I receptors (ALK1/ACVRL1, ALK3/BMPR1A, and ALK6) and inhibit ACVR1 more specifically while reducing side effects, a series of inhibitors were developed, including LDN-212854 (Mohedas et al., 2013), ML347 (Engers et al., 2013), OD36 (Sanchez-Duffhues et al., 2019), and "Compound 23" (Ullrich et al., 2022).

Among these modified compounds, LDN-212854 exhibits enhanced specificity in targeting ACVR1 (Mohedas, et al., 2013) and has undergone extensive research. LDN-212854 was developed from LDN-193189, and the modification increases the specificity for ACVR1 (ALK2) over BMPR1A (ALK3) by 66-fold and the specificity for ACVR1 over ALK5 by over 9000-fold (Mohedas, et al., 2013). LDN-212854 was shown to inhibit the formation of HO in FOP mouse models, which are transgenic mice with inducible constitutively active (ca) ACVR1 (*Acvr1^{Q207D}*) (Mohedas, et al., 2013; Dey et al., 2016). In vitro studies revealed that LDN-212854 can partly inhibit the mineralization of pericytes from human induced pluripotent stem cells (hiPSCs) from FOP patients with the *ACVR1* R206H mutation (Cai et al., 2015). However, under the same drug concentration (6 mg kg⁻¹, I.P., twice daily), there was no significant difference in the range of motion score between LDN-212854 and LDN-193189 in caACVR1 mice (Mohedas, et al., 2013). These results indicate that the difference in inhibition specificity between the two compounds is not significant in inhibiting HO formation in FOP, but may have some significance in reducing side effects.

Sanchez-Duffhues et al. (2019) used a chemistry-based platform of macrocyclic kinase inhibitors and FOP-derived endothelial cells to develop drugs targeting ACVR1. They found that OD36 and OD52 displayed improved specificity compared with LDN-193189, efficiently inhibited BMP-6 induced phosphorylation of Smad1/5/9 in KS483 cells, exclusively inhibited caACVR1 in C2C12 cells, and inhibited ACTA-induced osteogenic and chondrogenic differentiation of blood-derived endothelial colony-forming cells (ECFCs) from ACVR1^{R206H} FOP donors.

Jiang et al. (2018) identified a new ACVR1 inhibitor, "Compound 23", with 4(sulfamoyl)naphthyl as the pendant substituent at the 3-position of the pyrazolo[1,5-a] pyrimidine core, which exhibited better specificity for ACVR1 compared with other BMP type I receptor kinases and showed acceptable in vivo pharmacokinetics

In addition to the pyrazolo[1,5-a]pyrimidine scaffold, some researchers identified the imidazo[1,2-a]pyridine core as a suitable replacement system for potent ALK inhibitors and explored ways for improving its metabolic stability. As a result, "Compound 2" was developed, which has a specificity profile similar to that of LDN-193189, good oral bioavailability in rats, and a favorable brain: plasma ratio of 1.6 (Engers et al., 2020).

On the other hand, through screening 250 recombinant human kinases, Sanvitale et al. (2013) identified K02288 with a pyridine core, showing in vitro activity against ACVR1 at low nanomolar concentrations similar to LDN-193189. K02288 specifically inhibited the BMP-induced Smad pathway without affecting TGF- β signaling (Sanvitale, et al., 2013). However, similar to LDN-193189, K02288 shows low specificity for ACVR1 versus BMPR1A and ALK5 (Mohedas, et al., 2013).

Ullrich discovered several compounds with strikingly similar structural characteristics to K02288 analogs and detailed the optimization of a novel series of ACVR1 inhibitors based on a 2-aminopyrazine-3-carboxamide scaffold. Among these compounds, "Compound 23" demonstrated high aqueous solubility, membrane permeability, and microsomal stability, along with biochemical and cellular potency. It also showed good selectivity over other BMP and TGF- β receptor kinases (Ullrich, et al., 2022).

Sato et al. (2020) reported a series of bis-heteroaryl pyrazoles as ACVR1 inhibitors, including RK-59638. "Compound 8", based on RK-59638, was reported to bind to the ATP binding pocket of ACVR1^{R206H}, showing good membrane permeability and a strong ability to inhibit ALP activity in C2C12 (ACVR1 R206H) cells (Yamamoto et al., 2021).

Witten et al. (2022) designed a novel series of potent and selective ACVR1 inhibitors based on a bicyclic lactam scaffold. Their specific roles in cells remain to be studied.

Through in-house high-throughput screening and structure-guided drug design, Nguyen et al. (2022) recently discovered a series of pyrazolopyrimidines that act as selective ALK2 inhibitors. Among these, “Compound 16” showed high selectivity against other ALK subtypes, a favorable pharmacokinetic profile, and significant in vivo suppression of hepcidin which is regulated by ALK2 in hepatocytes.

Saracatinib, also known as AZD0530, is an oral SRC inhibitor developed by AstraZeneca for the treatment of ovarian adenocarcinoma (Hennequin et al., 2006), with favorable pharmacokinetic parameters and a suitable safety profile at a daily dosage of up to 250 mg (Hannon et al., 2010). Williams et al. (2021) screened a library of clinically tested small-molecule inhibitors and found that Saracatinib showed a good specific inhibitory effect on ACVR1. Saracatinib has been shown to inhibit the formation of HO in the *Acvr1*^{Q207D} and *Acvr1*^{R206H} transgenic mouse models of FOP (Williams, et al., 2021). Following treatment with Saracatinib, the transcriptome induced by ACTA in an FOP-derived endothelial cell model generated from induced pluripotent stem cells (iECs) was rescued to wild type levels. However, the transcription of SMAD2/3 target genes induced by ACTA was preserved in both wild type and FOP iECs (Hildebrandt et al., 2021). Saracatinib is currently undergoing a phase II clinical trial for FOP (NCT04307953) with a daily dose of 100 mg administered over a treatment period of 6 to 18 months.

To specifically target FOP-mutated ACVR1 without affecting wild-type ACVR1, Blueprint Medicines Inc. developed IPN60130 (also known as BLU-782) (Meng et al., 2022). In a Phase I clinical trial of IPN60130 in healthy volunteers (NCT03858075), the drug demonstrated good tolerability and favorable pharmacokinetic and pharmacodynamic properties (National Library of Medicine, 2019). IPN60130 is currently in Phase II clinical trials (NCT05039515).

INCB000928 is a potent and orally bioavailable small molecule inhibitor, showing high specificity for ACVR1 over ACVRL1 and BMPR1A in biochemical enzyme assays. INCB00928 also showed favorable pharmacokinetic properties in in vivo rodent studies. Serine/threonine kinase and ACVR1 are known to regulate serum ferrimodulin levels. Knockdown or loss of *Acvr1* in preclinical animal studies has been shown to result in decreased ferrimodulin production and elevated serum iron levels. INCB00928 inhibits BMP-induced ferrimodulin expression in both immortalized human liver cell lines and primary human liver cells. Furthermore, it reduces the levels of phosphorylated SMAD1/5 in the liver tissues of drug-treated mice (Chen et al., 2020). These findings suggest that INCB000928 holds promise as a potential treatment for FOP, and Phase II clinical trials for FOP (NCT05090891) are currently underway (National Library of Medicine, 2021).

Several research groups are working to identify natural inhibitors of ACVR1, often derived from marine organisms. For instance, Ohte et al. (2020) isolated a new diketopiperazine-like compound, named protuboxepin K, from the culture broth of the marine-derived fungal strain *Aspergillus* sp. BFM-0085. They also identified a unique sesquiterpene lactone, bicyclolamellolactone A, from the Indonesian marine sponge *Lamellodysidea* sp. (cf. *L. herbacea*) (Ohte et al., 2021). Similarly, Yamazaki et al. (2020) isolated dysidenin, herbassterol, and stelletasterol from EtOH extracts of the marine sponge *Dysidea* sp. (No. 256), following the screening of extracts from 188 Indonesian marine invertebrates. All these compounds demonstrated potent ALP inhibitory activity in C2C12 (R206H) cells. However, the specificity and effectiveness of these natural inhibitors in FOP mouse models is unknown. Nonetheless, the discovery of new natural inhibitors may provide new insights into the mechanism of action of ACVR1.

Several related inhibitors have shown their effectiveness in preclinical studies and results from others developed or identified more recently are yet to be disclosed.

2.2.3 Anti-ACVR1 antibodies

Building on the concept of blocking ACVR1, Daiichi Sankyo is developing the anti-ACVR1 monoclonal antibody DS-6016a in collaboration with Saitama Medical University in Japan. A Phase I clinical trial of DS-6016a has recently been completed (registered under NCT04818398), confirming the safety, tolerability, and pharmacokinetics of DS-6016a in healthy Japanese participants (Ifopa, 2022). However, specific experi-

mental results for DS-6016a are not yet available.

Another anti-ACVR1 monoclonal antibody, Rm0443, was isolated from a rat hybridoma cell line. This antibody binds to mouse and human ACVR1 through the residue H64, selectively inhabits ACVR1 signaling, and prevents HO in BMP7-induced HO mouse models and in one FOP mouse model (hALK2(R206H) FIE_x KI; CAG-cre/Esr1). However, it increased HO in another FOP mouse model (mAlk2(R206H) FIE_x KI). The contrasting effects of Rm0443 on human and mouse ACVR1 are attributed to a single residue difference at position 330. As Rm0443 is a rat antibody, researchers have developed humanized antibodies, including the aforementioned DS-6016a (Katagiri et al., 2023).

Conversely, some studies have reported that certain anti-ACVR1 antibodies exacerbated HO in FOP (ACVR1 R206H) mouse models, even though the Fab fragments of these antibodies still inhibited ACVR1 signaling in vitro (Aykul et al., 2022; Lees-Shepard et al., 2022). In view of these contradictory in vivo results, there is speculation that the divalent antibody for type I receptors prompts dimer formation to activate BMP signaling, which explains why the monovalent Fab fragment inhibits ACVR1 signaling in vitro as expected. The single residue difference at position 330 between human and mouse ACVR1 may provide clues to support this. Therefore, further studies are needed to elucidate the blocking mechanisms of anti-ACVR1 antibodies with ACVR1 from different species.

2.2.4 Anti-ACTA antibodies

As previously mentioned, ACTA, a secreted factor, is essential for the development of HO in FOP. Studies conducted on *Acvr1*^{[R206H]FIE_x+} transgenic mice, used as an FOP model, demonstrate that these mice do not develop HO either spontaneously or in response to local injury when treated with antibodies against ACTA. These antibodies block the activation of ACVR1 by ACTA (Hatsell, et al., 2015). Consequently, ongoing research is exploring the potential of anti-ACTA antibodies for the treatment of FOP.

Research has shown that treatment with Regeneron's anti-ACTA antibody, Garetosmab (also known as REGN247711), significantly inhibits HO in the FOP mouse model. Moreover, it not only inhibits the development of HO after the formation of bone tissue but also promotes the resorption of pre-existing heterotopic bone tissue to some extent (Upadhyay et al., 2017). According to reports, this antibody has recently completed a Phase II clinical trial (NCT04307953). FOP patients enrolled in the 28-week Garetosmab treatment trial exhibited a 25% reduction in overall heterotopic bone tissue and nearly 90% reduction in newly developed heterotopic bone tissue compared to patients treated with placebo. However, further research and analysis are needed to address the occurrence of serious adverse events, such as pneumonia and abscess, during the trial period (Ventura et al., 2021).

2.2.5 TGF- β related inhibitors

Given the established role of the TGF- β signaling pathway in FOP, several inhibitors of the pathway have been explored. NG-25, a small molecule inhibitor of TAK1, was evaluated and proven to limit heterotopic bone formation in a rat blast-associated trauma HO model and a murine burn tenotomy injury model (Strong et al., 2020). Similarly, Galunisertib (also known as LY2157299), a TGF- β receptor I kinase inhibitor that has shown good tolerability in patients with malignant tumors, has been demonstrated to prevent HO development in an acquired HO model (Mao et al., 2022). However, their efficacy in FOP remains to be studied and the results of relevant clinical studies are still to be disclosed.

3 Inflammation in FOP

3.1 Mechanisms

FOP lesions can form spontaneously or be triggered by inflammation or trauma. In the early stages of FOP, macrophages and mast cells infiltrate the flare-up sites. Depletion of these cells has been shown to impair HO in

an FOP (ACVR1 R206H) mouse model (Convente et al., 2018). However, other studies have shown that monocyte depletion, using liposome-encapsulated clodronate in injured skeletal muscle, can actually enhance the contribution of ECs to chondrogenesis and HO formation (Tirone et al., 2019). While the mechanisms of HO formation may vary between different subclasses, the heterogeneity of macrophages in FOP is an issue that requires further investigation.

As previously mentioned, macrophages in FOP patients exhibit increased secretion of TGF- β and prolonged activation of NF- κ B and p38 MAPK, with no significant changes in p-SMAD1/5. Barruet et al. (2018) reported increased levels of IL-3, IL-7, IL-8, and IL-10 cytokines in FOP. Hrkac et al. (2022) compared the proteomic profiles of EVs derived from pooled plasma of FOP patients during flare-up and remission phases. They found that HO occurred via processes of innate immunity and the Ephrin B signaling pathway, which may contribute to HO through macrophage chemotaxis and activation (Hrkac et al., 2022). Haviv et al. (2019) observed a marked increase in IL-1 β plasma levels during “flare-ups” in FOP patients. Del Zotto et al. (2017) characterized peripheral blood mononuclear cells (PBMC) by immunophenotyping in FOP and suggested that the increased expression of DNAX accessory molecule-1 (DNAM1, also known as CD226) in FOP monocytes may indicate an up-regulation of monocyte migration towards inflammation sites during “flare-ups”. A case-control study reported correlations between FOP genotype, acute flare-up status, and levels of adiponectin, tenascin-C, and kallikrein-7, suggesting a complex interplay between acute inflammation, chronic inflammation, and genetic background in FOP (Pignolo et al., 2022b).

Using RNA sequencing, Schoenmaker et al. (2023) described the transcriptomic changes induced by ACTA during osteoclast formation in CD14-positive monocytes from healthy controls and FOP patients. They found that “Inflammatory response” (GO: 0006954) was one of the GO terms most significantly associated with upregulated genes, and the class II major histocompatibility complex transactivator (CIITA) was one of the molecules associated with this term. The researchers speculated that ACTA-induced upregulation of CIITA may drive osteoclasts towards a more immunological role under high local concentrations of ACTA, but the specific significance of this finding remains to be explored (Schoenmaker et al., 2023).

In addition, mTOR signaling has been shown to play a crucial role in the early inflammatory phase of HO and is required for chondrogenesis and osteogenesis (Hino et al., 2018). Hypoxia, a known promoter of FOP, increases the intensity and duration of canonical BMP signaling through Rabaptin 5 (RABEP1)-mediated retention of ACVR1 in FOP connective tissue progenitor cells (CTPCs) (Wang et al., 2016). These findings suggest potential targets for new treatments.

3.2 Drug treatments

3.2.1 Anti-inflammatory drugs

Given the findings discussed above, anti-inflammatory drugs represent the cornerstone of current FOP treatment. High-dose corticosteroids are considered a preventative measure following traumatic injury or prior to unavoidable surgery, and prompt treatment following the first flare-up in a major joint is recommended (Smilde et al., 2022). Nonsteroidal anti-inflammatory drugs (NSAIDs) are also frequently used in FOP, as levels of prostaglandins are elevated in FOP patients experiencing a flare-up (Smilde, et al., 2022). However, results from clinical trials of these drugs in FOP are still lacking.

FOP also occurs in cats. A recent case report indicated that a regimen of 5 mg/kg BID enrofloxacin and hydrotherapy helped a cat with FOP-like manifestation demonstrate consistent improvement in endurance, quality of life, and range of motion over a period of 3 years. However, the cat's ACVR1 sequence was not determined, so they could not fully confirm that it had FOP. Regrettably, the cat ultimately succumbed to sudden death (Jacobsen et al., 2023).

3.2.2 Anti-IL-1 agents

In a case report, Haviv et al. (2019) diagnosed a boy with FOP who did not respond to anti-inflammatory

therapy. Upon detecting elevated IL-1 β plasma levels during a “flare-up”, they administered the anti-IL-1 agents anakinra and canakinumab. This treatment reduced the frequency of paroxysmal episodes and shortened their duration. However, there is still a relative lack of basic research data on this approach.

3.2.3 mTOR inhibitor Rapamycin.

Rapamycin, a potent and specific inhibitor of mTOR, is commonly used to prevent transplant rejection and lymphangiomyomatosis (Ventura, et al., 2021). Hino et al. (2017) identified Rapamycin as a candidate drug for FOP using an in vitro screening model and FOP-iPSC-derived iMSCs to screen for chemical compounds that could suppress the enhanced chondrogenesis of cells. Rapamycin has been reported to suppress HO in several FOP mouse models, including ACVR1^{R206H} and caACVR1^{Q207D} transgenic mice, as well as in mice with FOP-iPSC transplantation (Hino et al., 2017) and trauma-induced HO mice (Hino, et al., 2018).

However, recent studies showed that Rapamycin did not show clear benefits in reducing HO in two young patients with the ACVR1 R206H mutation (Kaplan et al., 2018b). It was reported that they continued to develop new flares during Rapamycin treatment. However, assessing the response to therapy is challenging since the therapeutic dose may not have been sufficient. Currently, Rapamycin is undergoing clinical trials (UMIN000028429) to evaluate its efficacy and safety in patients with FOP, and the outcomes have not been publicly released.

3.2.4 HIF1 α inhibitors

Based on studies of the HIF signaling pathway in FOP, several HIF1 α inhibitors, such as apigenin, PX-478, or imatinib, have been reported to prevent HO in FOP mouse models (Wang, et al., 2016; Zitvogel et al., 2016).

Imatinib is a tyrosine kinase inhibitor that was approved in the 1990s for the treatment of chronic myelogenous leukemia (Capdeville et al., 2002). The drug is well-tolerated with minimal side effects (Hochhaus et al., 2017). Wang et al. (2016) confirmed its effectiveness in inhibiting BMP signaling in SHED cells from FOP patients and reducing the formation of HO in the caACVR1^{Q207D} FOP mouse model. As a result, imatinib was used off-label in seven children with severe FOP with continuous flare-ups, and who were refractory to standard-of-care treatments (Kaplan et al., 2018a). A decrease in the intensity of flare-ups was reported in six of the patients who regularly took imatinib. However, since this was not a randomized controlled trial, clinical trials are needed to provide further evidence regarding the effectiveness of imatinib in preventing HO.

3.2.5 Other related drugs

Some researchers have considered FOP to be an autoinflammatory disease and have explored the potential effectiveness of disease-modifying antirheumatic drugs (DMARDs) for FOP treatment. One such drug is Tofacitinib, a Janus kinase (JAK) inhibitor approved by the FDA for the treatment of polyarticular juvenile idiopathic arthritis (JIA). In a retrospective observational study, Nikishina et al. (2023) gathered data from 13 genetically confirmed FOP patients who were treated with Tofacitinib at a dosage of 5 mg twice a day. They found that Tofacitinib reduced the number of flares, improved the range of motion in large joints, and improved imaging characteristics (Nikishina et al., 2023).

4 Other mechanisms and drug treatments in FOP

4.1 Mechanisms

4.1.1 Vascularization in FOP

The role of blood vessels and associated signaling pathways in promoting trauma-induced HO is well-established (Hwang et al., 2019), and histologic images of HE-stained sections from FOP patients demonstrate a richer vascular environment than that seen in nonhereditary HO (Ware et al., 2019). Monocytes

from FOP patients have been shown to secrete increased levels of VEGF in response to inflammation, compared with controls (Tirone, et al., 2019). Using the previously mentioned FOP iEC, RNA-Seq analysis revealed that ACTA-induced pSMAD1/5 was upregulated exclusively in the FOP iECs. This resulted in a FOP-specific transcriptome characterized by the involvement of genes in blood vessel formation and the activation of BMP and NOTCH pathways (Hildebrandt, et al., 2021). As a result, vascularization has become a new area of focus in FOP research.

4.1.2 Chondrogenesis and endochondral ossification in FOP

Chondrogenesis and endochondral ossification represent one of the primary mechanisms of HO in FOP. Following muscle injury in FOP mice, a subset of FAPs organize and undergo chondrogenic differentiation, followed by hypertrophy and mineralization of the cartilage matrix. Subsequently, the cartilage is resorbed by osteoclasts and remodeled into bone (Pierce and Perrien, 2021). The "seed cells" of HO differ across tissues, but the underlying mechanism remains similar. Retinoid signaling, mediated by nuclear retinoic acid receptors (RARs), including RAR α , β , and γ , plays a crucial role in chondrogenesis and skeletal formation (Weston et al., 2003; Pacifici, 2018). This includes decreased retinoid signaling required in chondrogenic cell differentiation and cartilage maturation (Pignolo and Pacifici, 2021), providing new insights for treatment. Moreover, Yang et al. (2021) found that increased BMP signaling via caACVR1 induced cranial neural crest cells (CNCCs) in mice towards a chondrogenic fate by stimulating mTORC1 activity and subsequently suppressing autophagic degradation of β -catenin. As mentioned earlier, the Ephrin B signaling pathway, identified in the EVs of FOP patients, could enhance endochondral ossification in fracture repair and may play a similar role in FOP (Hrkac, et al., 2022).

4.1.3 FOP: a potential segmental progeroid syndrome

Pignolo et al. (2020) observed similarities between certain features related to dysregulated BMP signaling in FOP and those seen in accelerated aging, leading them to suggest that FOP might be considered a segmental progeroid syndrome. They proposed that hypoxia, cellular damage, and inflammation resulting from soft tissue injury could lead to the accumulation of senescent cells, similar to what is observed in senescent tissues (Pignolo, et al., 2020). They also posited that aging could promote tissue transformation towards cartilage formation in FOP muscle (Wang et al., 2021b). This perspective provides valuable insights for potential therapeutic interventions in FOP.

4.1.4 Dysregulated temperature response in FOP

A study by Wang et al. found that FOP CTPCs exhibited a dysregulated temperature response, and chondrogenesis was temperature-sensitive and amplified in FOP CTPCs (Wang et al., 2021a). This discovery adds temperature to the list of microenvironmental factors that likely influence FOP and suggests potential therapeutic avenues, such as antipyretics.

4.1.5 Decreased apoptosis levels in FOP

Stanley et al. (2022) building on previous studies of FAP, found that Acvr1^{R206H/+} FAPs failed to decline due to significantly decreased levels of apoptosis during regeneration in FOP mouse models. This suggests that apoptosis-related modulators targeting FAP may be new therapeutic candidates (Stanley et al., 2022).

4.2 Drug treatments

4.2.1 Retinoid agonists

4.2.1.1 NRX195183

Based on the result that atRA treatment of limb mesenchymal skeletal precursor cells blocks chondrogenesis and cartilage formation, Shimono et al.(2010) tested the RAR α agonist NRX195183 in an acquired HO

mouse model with subcutaneous implantation of rhBMP-2/Matrigel scaffolds, and found that NRX195183 reduced HO formation. Although NRX195183 has completed a Phase 1 clinical trial (Herbert et al., 2007), further testing at high doses is required.

4.2.1.2 Palovarotene

Shimono et al. (2011) developed the RAR γ agonist Palovarotene (also known as R667) based on the usefulness of NRX195183. Palovarotene is a highly specific RAR γ agonist, effectively inhibiting spontaneous HO in ACVR1 R206H mice (Chakkalakal et al., 2016). Several Phase II (NCT02190747, NCT02279095 and NCT02979769) and III (NCT03312634) clinical trials have recently been completed, and some results are pending. The single-arm, open-label, Phase III MOVE trial (NCT03312634) showed that mean annualized new HO volume was 60% lower in the MOVE group versus the NHS (Pignolo et al., 2023). Consequently, Palovarotene (Sohonos) was recently approved by the USFDA for reducing the volume of HO in adults and children aged 8 years and older for females, and 10 years and older for males with FOP. However, it has been observed to cause premature maturation and closure of growth plates in some FOP patients (Pignolo and Pacifici, 2021), and there is an increased risk of vertebral fracture in patients treated with palovarotene (Pignolo, et al., 2023). Other potential risks include pancreatitis, hearing and vision impairment, mouth ulcers, and dry skin (Kitoh, 2020).

4.2.2 Senolytic drugs: Dasatinib and quercetin

Approaching FOP as a segmental progeroid syndrome, Wang et al. (2021b) used Dasatinib and quercetin to clear and reduce the senescence-associated secretory phenotype (SASP). The results showed that these senolytic drugs reduced BMP pathway signaling in FOP SHED and ameliorated HO in FOP mouse models (Wang, et al., 2021b).

4.2.3 PI3K inhibitor Alpelisib

PI3K α is critical for bone formation through the regulation of SMAD1 activity (Gamez et al., 2016), and mTOR signaling, downstream of PI3K/AKT, plays a critical role in HO and FOP (Hino, et al., 2018). Valer et al. (2019) found that the PI3K inhibitor Alpelisib (also known as BYL719 and marketed as PiqrayTM) was effective at suppressing HO in the ACVR1^{Q207D} mouse model. Alpelisib was well tolerated and was approved by the FDA for hyperactive PI3K (PIK3CA)-altered solid tumors (Rugo et al., 2020), with the most frequent adverse effects being hyperglycaemia and rash (Ventura, et al., 2021). PI3K α inhibitors could negatively target ACVR1 kinase activity (Qureshi et al., 2017). However, further studies are required to optimize Alpelisib treatment in FOP.

5 Discussion

Despite ongoing efforts, the explicit mechanism and ideal drugs for FOP remain elusive. Several challenges remain, particularly concerning the specificity of ACVR1 inhibitors. Current inhibitors have a limited ability to distinguish between ACVR1 and ACVRL1, as well as between wild-type ACVR1 and mutant ACVR1, and their use would result in side effects.

As research into FOP deepens, new mechanisms of its occurrence and progression are being discovered, providing new therapeutic targets. However, the cell/tissue/stage specificity of new drugs remains a challenge, as evidenced by Palovarotene causing premature maturation and closure of growth plates in some FOP patients. Furthermore, there are concerns regarding the targeting efficacy of FOP therapies that have shown initial clinical promise, necessitating subsequent modifications to enhance targeting and reduce side effects. Combining drugs with novel technologies, such as engineered exosomes, may ensure effective and precise targeting of therapeutic drugs to the tissues affected by FOP.

In addition, while many drugs have proven effective in animal models of FOP, the mechanisms of these drugs need further exploration. For instance, Dasatinib, both a senolytic drug and an Src inhibitor, is effective in

ameliorating HO by clearing and reducing SASP (Wang, et al., 2021b). Given that four Src kinase family members were identified in the EVs of a FOP flare-up group (Hrkac, et al., 2022), Dasatinib may also play a role in inhibiting HO by influencing this aspect.

The development of therapeutic drugs for FOP relies on progress in understanding the mechanisms underlying its occurrence and progression, as well as studies of its identification and confirmation of targets. Emerging technologies, such as genomics, proteomics, and high-throughput screening, can be used to identify potential targets, detect specific molecules or pathways that are dysregulated in the disease, offer new possibilities for old drugs, and validate them *in vitro* and *in vivo* to design new therapeutic drugs. Through a cellular NanoBRET target engagement assay, Fortin et al. (2020) identified a binding interaction between E6201 and ACVR1. E6201, an ATP-competitive MEK1 inhibitor, has been reported to be effective in BRAF V600E-mutated metastatic malignant melanoma (Babiker et al., 2019). It inhibits the activation of the BMP pathway induced by ca-ACVR1 and shows a stronger inhibitory effect on pathway activation induced by mutant ACVR1 compared with wild-type ACVR1 (Fortin, et al., 2020). Moreover, E6201 prolongs survival in mice xenografted with SU-DIPG-XXXVI or HSJD-DIPG-007 tumor cells carrying FOP-related ACVR1 mutations (Fortin, et al., 2020). Although the drug was not developed to target FOP, it could provide some clues for drug development.

CRISPR-Cas9 (clustered regularly interspaced short palindromic repeats-associated protein 9) is also an emerging technology. Using an improved CRISPR-Cas9 system with high specificity, Kawamata et al. (2023) succeeded in editing a mono-allelic 617 G>A (R206H) *Acvr1* mutation in WT/WT mESC, which subsequently causes ectopic ossification in mouse. This method greatly improves the efficiency of editing, but the effect of reverse application of this method to treatment remains to be further studied. However, given the regenerative capacity of progenitors/stem cells and the enhanced immune response in FOP, the cost performance of CRISPR-Cas9 for clinical treatment remains to be determined.

6 Conclusions

The treatment of FOP has long posed significant challenges for medical professionals due to the rarity and variability of the disease. However, years of continuous study on FOP have led to a deeper understanding of its mechanisms, thereby guiding the development of new drug candidates aimed at alleviating patients' suffering. Ongoing research and clinical trials provide hope for improved management strategies and treatment options. Encouragingly, we are gradually progressing toward a potential "cure".

Acknowledgments

This work is supported by the National Natural Science Foundation of China (81920108012, 82270959, 81970903), Jilin University Norman Bethune Program (2023B28) and Department of Finance of Jilin Province (jcsz2023481-37).

Author contributions

Yi-jun ZHOU wrote and edited the manuscript. Ce SHI edited the manuscript. Hong-chen SUN reviewed and edited the manuscript. All authors read and approved the final manuscript and, therefore, had full access to all the data in the study and take responsibility for the integrity and security of the data.

Compliance with ethics guidelines

Yi-jun ZHOU, Ce SHI, and Hong-chen SUN declare that they have no conflict of interest.

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008 (5). Informed consent was obtained from all patients for being included in the study. Additional informed consent was obtained from all patients for whom identifying information is included in this article.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Aykul S, Corpina RA, Goebel EJ, et al., 2020. Activin a forms a non-signaling complex with acvr1 and type ii activin/bmp receptors via its finger 2 tip loop. *eLife*, 9 <https://doi.org/10.7554/eLife.54582>
- Aykul S, Huang L, Wang L, et al., 2022. Anti-acvr1 antibodies exacerbate heterotopic ossification in fibrodysplasia ossificans progressiva (fop) by activating fop-mutant acvr1. *J Clin Invest*, 132(12) <https://doi.org/10.1172/jci153792>
- Babiker HM, Byron SA, Hendricks WPD, et al., 2019. E6201, an intravenous mek1 inhibitor, achieves an exceptional response in braf v600e-mutated metastatic malignant melanoma with brain metastases. *Invest New Drugs*, 37(4):636-645. <https://doi.org/10.1007/s10637-018-0668-8>
- Barruet E, Morales BM, Cain CJ, et al., 2018. Nf-kappab/mapk activation underlies acvr1-mediated inflammation in human heterotopic ossification. *JCI Insight*, 3(22) <https://doi.org/10.1172/jci.insight.122958>
- Brennan TA, Lindborg CM, Bergbauer CR, et al., 2018. Mast cell inhibition as a therapeutic approach in fibrodysplasia ossificans progressiva (fop). *Bone*, 109:259-266. <https://doi.org/10.1016/j.bone.2017.08.023>
- Cai J, Orlova VV, Cai X, et al., 2015. Induced pluripotent stem cells to model human fibrodysplasia ossificans progressiva. *Stem Cell Reports*, 5(6):963-970. <https://doi.org/10.1016/j.stemcr.2015.10.020>
- Capdeville R, Buchdunger E, Zimmermann J, et al., 2002. Glivec (sti571, imatinib), a rationally developed, targeted anticancer drug. *Nat Rev Drug Discov*, 1(7):493-502. <https://doi.org/10.1038/nrd839>
- Chaikuad A, Alfano I, Kerr G, et al., 2012. Structure of the bone morphogenetic protein receptor alk2 and implications for fibrodysplasia ossificans progressiva. *J Biol Chem*, 287(44):36990-36998. <https://doi.org/10.1074/jbc.M112.365932>
- Chakkalakal SA, Uchibe K, Convente MR, et al., 2016. Palovarotene inhibits heterotopic ossification and maintains limb mobility and growth in mice with the human acvr1(r206h) fibrodysplasia ossificans progressiva (fop) mutation. *J Bone Miner Res*, 31(9):1666-1675. <https://doi.org/10.1002/jbmr.2820>
- Chen Y, Stubbs MC, Pusey M, et al. 2020. Characterization of incb00928, a potent and selective alk2 inhibitor for the treatment of anemia. Available from <https://ashpublications.org/blood/article/136/Supplement%201/52/472793/Characterization-of-INCB00928-a-Potent-and>
- Chiu W, Lin TY, Chang YC, et al., 2021. An update on gene therapy for inherited retinal dystrophy: Experience in leber congenital amaurosis clinical trials. *Int J Mol Sci*, 22(9) <https://doi.org/10.3390/ijms22094534>
- Convente MR, Chakkalakal SA, Yang E, et al., 2018. Depletion of mast cells and macrophages impairs heterotopic ossification in an acvr1r206h mouse model of fibrodysplasia ossificans progressiva. *Journal of Bone and Mineral Research*, 33(2):269-282. <https://doi.org/10.1002/jbmr.3304>
- Cuny GD, Yu PB, Laha JK, et al., 2008. Structure-activity relationship study of bone morphogenetic protein (bmp) signaling inhibitors. *Bioorg Med Chem Lett*, 18(15):4388-4392. <https://doi.org/10.1016/j.bmcl.2008.06.052>
- De Ruiter R, Groot T, Schoenmaker T, et al., 2022. Fibroblasts from patients with fibrodysplasia ossificans progressiva exhibit increased activin a production. *Journal of Bone and Mineral Research*, 37:320-320.
- De Ruiter RD, Wisse LE, Schoenmaker T, et al., 2023. Tgf-beta induces activin a production in dermal fibroblasts derived from patients with fibrodysplasia ossificans progressiva. *International Journal of Molecular Sciences*, 24(3):2299. <https://doi.org/10.3390/ijms24032299>
- Del Zotto G, Antonini F, Azzari I, et al., 2017. Peripheral blood mononuclear cell immunophenotyping in fibrodysplasia ossificans progressiva patients: Evidence for monocyte dnam1 up - regulation. *Cytometry Part B: Clinical Cytometry*, 94(4):613-622. <https://doi.org/10.1002/cyto.b.21594>
- Dey D, Bagarova J, Hatsell SJ, et al., 2016. Two tissue-resident progenitor lineages drive distinct phenotypes of heterotopic ossification. *Sci Transl Med*, 8(366):366ra163. <https://doi.org/10.1126/scitranslmed.aaf1090>
- Engers DW, Frist AY, Lindsley CW, et al., 2013. Synthesis and structure-activity relationships of a novel and selective bone morphogenetic protein receptor (bmp) inhibitor derived from the pyrazolo[1.5-a]pyrimidine scaffold of dorsomorphin: The discovery of ml347 as an alk2 versus alk3 selective mlpcn probe. *Bioorg Med Chem Lett*, 23(11):3248-3252. <https://doi.org/10.1016/j.bmcl.2013.03.113>
- Engers DW, Bollinger SR, Felts AS, et al., 2020. Discovery, synthesis and characterization of a series of 7-aryl-imidazo[1,2-a]pyridine-3-ylquinolines as activin-like kinase (alk) inhibitors. *Bioorg Med Chem Lett*, 30(18):127418. <https://doi.org/10.1016/j.bmcl.2020.127418>
- Fortin J, Tian R, Zarrabi I, et al., 2020. Mutant acvr1 arrests glial cell differentiation to drive tumorigenesis in pediatric gliomas. *Cancer Cell*, 37(3):308-323 e312. <https://doi.org/10.1016/j.ccell.2020.02.002>
- Gamez B, Rodriguez-Carballo E, Graupera M, et al., 2016. Class i pi-3-kinase signaling is critical for bone formation through regulation of smad1 activity in osteoblasts. *J Bone Miner Res*, 31(8):1617-1630. <https://doi.org/10.1002/jbmr.2819>
- Hannon RA, Clack G, Rimmer M, et al., 2010. Effects of the src kinase inhibitor saracatinib (azd0530) on bone

- turnover in healthy men: A randomized, double-blind, placebo-controlled, multiple-ascending-dose phase I trial. *J Bone Miner Res*, 25(3):463-471. <https://doi.org/10.1359/jbmr.090830>
- Hao J, Ho JN, Lewis JA, et al., 2010. In vivo structure-activity relationship study of dorsomorphin analogues identifies selective vegf and bmp inhibitors. *ACS Chem Biol*, 5(2):245-253. <https://doi.org/10.1021/cb9002865>
- Hatsell SJ, Idone V, Wolken DM, et al., 2015. Acvr1r206h receptor mutation causes fibrodysplasia ossificans progressiva by imparting responsiveness to activin a. *Sci Transl Med*, 7(303):303ra137. <https://doi.org/10.1126/scitranslmed.aac4358>
- Haviv R, Moshe V, De Benedetti F, et al., 2019. Is fibrodysplasia ossificans progressiva an interleukin-1 driven auto-inflammatory syndrome? *Pediatric Rheumatology*, 17(1) <https://doi.org/10.1186/s12969-019-0386-6>
- Hennequin LF, Allen J, Breed J, et al., 2006. N-(5-chloro-1,3-benzodioxol-4-yl)-7-[2-(4-methylpiperazin-1-yl)ethoxy]-5-(tetrahydro-2h-pyran-4-yloxy)quinazolin-4-amine, a novel, highly selective, orally available, dual-specific c-src/abl kinase inhibitor. *J Med Chem*, 49(22):6465-6488. <https://doi.org/10.1021/jm060434q>
- Herbert KE, Walkley CR, Winkler IG, et al., 2007. Granulocyte colony-stimulating factor and an aralpha specific agonist, vtp195183, synergize to enhance the mobilization of hematopoietic progenitor cells. *Transplantation*, 83(4):375-384. <https://doi.org/10.1097/01.tp.0000251376.75347.b4>
- Hildebrandt S, Kampfrath B, Fischer K, et al., 2021. Activin-induced smad1/5 signaling in an ipsc derived ec model of fibrodysplasia ossificans progressiva (fop) can be rescued by the drug candidate saracatinib. *Stem Cell Reviews and Reports*, 17(3):1039-1052. <https://doi.org/10.1007/s12015-020-10103-9>
- Hino K, Horigome K, Nishio M, et al., 2017. Activin-a enhances mtor signaling to promote aberrant chondrogenesis in fibrodysplasia ossificans progressiva. *J Clin Invest*, 127(9):3339-3352. <https://doi.org/10.1172/JCI93521>
- Hino K, Zhao C, Horigome K, et al., 2018. An mtor signaling modulator suppressed heterotopic ossification of fibrodysplasia ossificans progressiva. *Stem Cell Reports*, 11(5):1106-1119. <https://doi.org/10.1016/j.stemcr.2018.10.007>
- Hochhaus A, Larson RA, Guilhot F, et al., 2017. Long-term outcomes of imatinib treatment for chronic myeloid leukemia. *N Engl J Med*, 376(10):917-927. <https://doi.org/10.1056/NEJMoal609324>
- Hrkac S, Novak R, Salai G, et al., 2022. Heterotopic ossification vs. Fracture healing: Extracellular vesicle cargo proteins shed new light on bone formation. *Bone Reports*, 16:101177. <https://doi.org/10.1016/j.bonr.2022.101177>
- Hu Y, Hao X, Liu C, et al., 2021. Acvr1 deletion in osteoblasts impaired mandibular bone mass through compromised osteoblast differentiation and enhanced srnk1-induced osteoclastogenesis. *J Cell Physiol*, 236(6):4580-4591. <https://doi.org/10.1002/jcp.30183>
- Hwang C, Marini S, Huber AK, et al., 2019. Mesenchymal vegfa induces aberrant differentiation in heterotopic ossification. *Bone Res*, 7:36. <https://doi.org/10.1038/s41413-019-0075-6>
- Hwang CD, Pagani CA, Nunez JH, et al., 2022. Contemporary perspectives on heterotopic ossification. *JCI Insight*, 7(14) <https://doi.org/10.1172/jci.insight.158996>
- Ifopa. 2022. Ds-6016a. Available from https://www.ifopa.org/ds_6016a
- Jacobsen KL, Wiebe V, Davidson AP, et al., 2023. Use of enrofloxacin and hydrotherapy in the management of fibrodysplasia ossificans progressiva (fop) in a savannah cat. *Topics in Companion Animal Medicine*, 52:100757. <https://doi.org/10.1016/j.tcam.2022.100757>
- Jia S, Meng A, 2021. Tgfbeta family signaling and development. *Development*, 148(5) <https://doi.org/10.1242/dev.188490>
- Jiang JK, Huang X, Shamim K, et al., 2018. Discovery of 3-(4-sulfamoylnaphthyl)pyrazolo[1,5-a]pyrimidines as potent and selective alk2 inhibitors. *Bioorg Med Chem Lett*, 28(20):3356-3362. <https://doi.org/10.1016/j.bmcl.2018.09.006>
- Kaplan FS, Andolina JR, Adamson PC, et al., 2018a. Early clinical observations on the use of imatinib mesylate in fop: A report of seven cases. *Bone*, 109:276-280. <https://doi.org/10.1016/j.bone.2017.07.019>
- Kaplan FS, Zeitlin L, Dunn SP, et al., 2018b. Acute and chronic rapamycin use in patients with fibrodysplasia ossificans progressiva: A report of two cases. *Bone*, 109:281-284. <https://doi.org/10.1016/j.bone.2017.12.011>
- Kaplan J, Kaplan FS, Shore EM, 2012. Restoration of normal bmp signaling levels and osteogenic differentiation in fop mesenchymal progenitor cells by mutant allele-specific targeting. *Gene Ther*, 19(7):786-790. <https://doi.org/10.1038/gt.2011.152>
- Katagiri T, Tsukamoto S, Kuratani M, et al., 2023. A blocking monoclonal antibody reveals dimerization of intracellular domains of alk2 associated with genetic disorders. *Nature Communications*, 14(1)

- <https://doi.org/10.1038/s41467-023-38746-5>
- Kawamata M, Suzuki HI, Kimura R, et al., 2023. Optimization of cas9 activity through the addition of cytosine extensions to single-guide rnas. *Nature Biomedical Engineering*, 7(5):672-691. <https://doi.org/10.1038/s41551-023-01011-7>
- Kitoh H, 2020. Clinical aspects and current therapeutic approaches for fop. *Biomedicines*, 8(9) <https://doi.org/10.3390/biomedicines8090325>
- Lees-Shepard JB, Stoessel SJ, Chandler JT, et al., 2022. An anti-acvr1 antibody exacerbates heterotopic ossification by fibro-adipogenic progenitors in fibrodysplasia ossificans progressiva mice. *Journal of Clinical Investigation*, 132(12) <https://doi.org/10.1172/jci153795>
- Lim KH, Han Z, Jeon HY, et al., 2020. Antisense oligonucleotide modulation of non-productive alternative splicing upregulates gene expression. *Nature Communications*, 11(1) <https://doi.org/10.1038/s41467-020-17093-9>
- Luo Y, Alsamarah A, Zhang K, et al., 2016. Development of new therapeutic agents for fibrodysplasia ossificans progressiva. *Curr Mol Med*, 16(1):4-11. <https://doi.org/10.2174/1566524016666151222142446>
- Mao D, Mi J, Pan X, et al., 2022. Galunisertib attenuates progression of trauma-induced heterotopic ossification via blockage of smad2/3 signaling in mice. *Eur J Pharmacol*, 928:175109. <https://doi.org/10.1016/j.ejphar.2022.175109>
- Maruyama R, Yokota T, 2018. Morpholino-mediated exon skipping targeting human acvr1/alk2 for fibrodysplasia ossificans progressiva. 1828:497-502. https://doi.org/10.1007/978-1-4939-8651-4_32
- Maruyama R, Nguyen Q, Roshmi RR, et al., 2022. Allele-selective lna gapmers for the treatment of fibrodysplasia ossificans progressiva knock down the pathogenic acvr1r206h transcript and inhibit osteogenic differentiation. *Nucleic Acid Therapeutics*, 32(3):185-193. <https://doi.org/10.1089/nat.2021.0009>
- National Library of Medicine, 2019. Safety, tolerability, pharmacokinetics, and food effect of blu-782 in healthy adults. Available from <https://classic.clinicaltrials.gov/ct2/show/NCT03858075?term=NCT03858075&draw=2&rank=1>
- National Library of Medicine, 2021. To assess the efficacy, safety, and tolerability of incb000928 in participants with fibrodysplasia ossificans progressiva (progress). Available from <https://classic.clinicaltrials.gov/ct2/show/NCT05090891>
- Meng X, Wang H, Hao J, 2022. Recent progress in drug development for fibrodysplasia ossificans progressiva. *Mol Cell Biochem*, 477(10):2327-2334. <https://doi.org/10.1007/s11010-022-04446-9>
- Mohedas AH, Xing X, Armstrong KA, et al., 2013. Development of an alk2-biased bmp type i receptor kinase inhibitor. *ACS Chem Biol*, 8(6):1291-1302. <https://doi.org/10.1021/cb300655w>
- Nagar G, Mittal P, Gupta SRR, et al., 2023. Multi-omics therapeutic perspective on acvr1 gene: From genetic alterations to potential targeting. *Briefings in Functional Genomics*, 22(2):123-142. <https://doi.org/10.1093/bfpg/elac026>
- Nguyen MH, Atasoylu O, Wu L, et al., 2022. Discovery of novel pyrazolopyrimidines as potent, selective, and orally bioavailable inhibitors of alk2. *ACS Medicinal Chemistry Letters*, 13(7):1159-1164. <https://doi.org/10.1021/acsmchemlett.2c00206>
- Nikishina IP, Arsenyeva SV, Matkava VG, et al., 2023. Successful experience of tofacitinib treatment in patients with fibrodysplasia ossificans progressiva. *Pediatric Rheumatology*, 21(1) <https://doi.org/10.1186/s12969-023-00856-1>
- Ohte S, Yamazaki H, Takahashi O, et al., 2021. Inhibitory effects of sesquiterpene lactones from the indonesian marine sponge lamellodysidea cf. Herbacea on bone morphogenetic protein-induced osteoblastic differentiation. *Bioorganic & Medicinal Chemistry Letters*, 35:127783. <https://doi.org/10.1016/j.bmcl.2021.127783>
- Pacifici M, 2018. Retinoid roles and action in skeletal development and growth provide the rationale for an ongoing heterotopic ossification prevention trial. *Bone*, 109:267-275. <https://doi.org/10.1016/j.bone.2017.08.010>
- Pierce JL, Perrien DS, 2021. Do interactions of vitamin d3 and bmp signaling hold implications in the pathogenesis of fibrodysplasia ossificans progressiva? *Current Osteoporosis Reports*, 19(3):358-367. <https://doi.org/10.1007/s11914-021-00673-z>
- Pignolo RJ, Wang H, Kaplan FS, 2020. Fibrodysplasia ossificans progressiva (fop): A segmental progeroid syndrome. *Frontiers in Endocrinology*, 10 <https://doi.org/10.3389/fendo.2019.00908>
- Pignolo RJ, Pacifici M, 2021. Retinoid agonists in the targeting of heterotopic ossification. *Cells*, 10(11) <https://doi.org/10.3390/cells10113245>
- Pignolo RJ, Bedford-Gay C, Cali A, et al., 2022a. Current challenges and opportunities in the care of patients with fibrodysplasia ossificans progressiva (fop): An international, multi-stakeholder perspective. *Orphanet J Rare Dis*, 17(1):168. <https://doi.org/10.1186/s13023-022-02224-w>

- Pignolo RJ, Mccarrick - Walmsley R, Wang H, et al., 2022b. Plasma - soluble biomarkers for fibrodysplasia ossificans progressiva (fop) reflect acute and chronic inflammatory states. *Journal of Bone and Mineral Research*, 37(3):475-483. <https://doi.org/10.1002/jbmr.4492>
- Pignolo RJ, Hsiao EC, Al Mukaddam M, et al., 2023. Reduction of new heterotopic ossification (ho) in the open - label, phase 3 move trial of palovarotene for fibrodysplasia ossificans progressiva (fop). *Journal of Bone and Mineral Research*, 38(3):381-394. <https://doi.org/10.1002/jbmr.4762>
- Qureshi AT, Dey D, Sanders EM, et al., 2017. Inhibition of mammalian target of rapamycin signaling with rapamycin prevents trauma-induced heterotopic ossification. *Am J Pathol*, 187(11):2536-2545. <https://doi.org/10.1016/j.ajpath.2017.07.010>
- Rugo HS, Andre F, Yamashita T, et al., 2020. Time course and management of key adverse events during the randomized phase iii solar-1 study of pi3k inhibitor alpelisib plus fulvestrant in patients with hr-positive advanced breast cancer. *Ann Oncol*, 31(8):1001-1010. <https://doi.org/10.1016/j.annonc.2020.05.001>
- Sanchez-Duffhues G, Williams E, Benderitter P, et al., 2019. Development of macrocycle kinase inhibitors for alk2 using fibrodysplasia ossificans progressiva-derived endothelial cells. *JBMR Plus*, 3(11):e10230. <https://doi.org/10.1002/jbm4.10230>
- Sanvitale CE, Kerr G, Chaikuad A, et al., 2013. A new class of small molecule inhibitor of bmp signaling. *PLoS One*, 8(4):e62721. <https://doi.org/10.1371/journal.pone.0062721>
- Sato T, Sekimata K, Sakai N, et al., 2020. Structural basis of activin receptor-like kinase 2 (r206h) inhibition by bis-heteroaryl pyrazole-based inhibitors for the treatment of fibrodysplasia ossificans progressiva identified by the integration of ligand-based and structure-based drug design approaches. *ACS Omega*, 5(20):11411-11423. <https://doi.org/10.1021/acsomega.9b04245>
- Schoenmaker T, Zwaak J, Loos BG, et al., 2023. Transcriptomic differences underlying the activin-a induced large osteoclast formation in both healthy control and fibrodysplasia ossificans progressiva osteoclasts. *International Journal of Molecular Sciences*, 24(7):6822. <https://doi.org/10.3390/ijms24076822>
- She D, Zhang K, 2018. Fibrodysplasia ossificans progressiva in china. *Bone*, 109:101-103. <https://doi.org/10.1016/j.bone.2017.11.016>
- Shi W, Shi S, Cai J, et al., 2013. Antisense-oligonucleotide mediated exon skipping in activin-receptor-like kinase 2: Inhibiting the receptor that is overactive in fibrodysplasia ossificans progressiva. *PLoS ONE*, 8(7):e69096. <https://doi.org/10.1371/journal.pone.0069096>
- Shimono K, Morrison TN, Tung WE, et al., 2010. Inhibition of ectopic bone formation by a selective retinoic acid receptor alpha-agonist: A new therapy for heterotopic ossification? *J Orthop Res*, 28(2):271-277. <https://doi.org/10.1002/jor.20985>
- Shimono K, Tung WE, Macolino C, et al., 2011. Potent inhibition of heterotopic ossification by nuclear retinoic acid receptor-gamma agonists. *Nat Med*, 17(4):454-460. <https://doi.org/10.1038/nm.2334>
- Shore EM, Xu M, Feldman GJ, et al., 2006. A recurrent mutation in the bmp type i receptor acvr1 causes inherited and sporadic fibrodysplasia ossificans progressiva. *Nat Genet*, 38(5):525-527. <https://doi.org/10.1038/ng1783>
- Smilde B, Botman E, De Ruiter R, et al., 2022. Monitoring and management of fibrodysplasia ossificans progressiva: Current perspectives. *Orthopedic Research and Reviews*, Volume 14:113-120. <https://doi.org/10.2147/orr.s337491>
- Song G-A, Kim H-J, Woo K-M, et al., 2010. Molecular consequences of the acvr1r206h mutation of fibrodysplasia ossificans progressiva. *Journal of Biological Chemistry*, 285(29):22542-22553. <https://doi.org/10.1074/jbc.M109.094557>
- Stanley A, Tichy ED, Kocan J, et al., 2022. Dynamics of skeletal muscle-resident stem cells during myogenesis in fibrodysplasia ossificans progressiva. *NPJ Regen Med*, 7(1):5. <https://doi.org/10.1038/s41536-021-00201-8>
- Strong AL, Spreadborough PJ, Pagani CA, et al., 2020. Small molecule inhibition of non-canonical (tak1-mediated) bmp signaling results in reduced chondrogenic ossification and heterotopic ossification in a rat model of blast-associated combat-related lower limb trauma. *Bone*, 139:115517. <https://doi.org/10.1016/j.bone.2020.115517>
- Tirone M, Giovenzana A, Vallone A, et al., 2019. Severe heterotopic ossification in the skeletal muscle and endothelial cells recruitment to chondrogenesis are enhanced by monocyte/macrophage depletion. *Frontiers in Immunology*, 10 <https://doi.org/10.3389/fimmu.2019.01640>
- Towler OW, Shore EM, 2022. Bmp signaling and skeletal development in fibrodysplasia ossificans progressiva (fop). *Dev Dyn*, 251(1):164-177. <https://doi.org/10.1002/dvdy.387>
- Ullrich T, Arista L, Weiler S, et al., 2022. Discovery of a novel 2-aminopyrazine-3-carboxamide as a potent and selective inhibitor of activin receptor-like kinase-2 (alk2) for the treatment of fibrodysplasia ossificans

- progressiva. *Bioorg Med Chem Lett*, 64:128667. <https://doi.org/10.1016/j.bmcl.2022.128667>
- Upadhyay J, Xie L, Huang L, et al., 2017. The expansion of heterotopic bone in fibrodysplasia ossificans progressiva is activin a-dependent. *J Bone Miner Res*, 32(12):2489-2499. <https://doi.org/10.1002/jbmr.3235>
- Valer JA, Sanchez-De-Diego C, Gamez B, et al., 2019. Inhibition of phosphatidylinositol 3-kinase alpha (pi3kalpha) prevents heterotopic ossification. *EMBO Mol Med*, 11(9):e10567. <https://doi.org/10.15252/emmm.201910567>
- Ventura F, Williams E, Ikeya M, et al., 2021. Challenges and opportunities for drug repositioning in fibrodysplasia ossificans progressiva. *Biomedicines*, 9(2) <https://doi.org/10.3390/biomedicines9020213>
- Wang H, Lindborg C, Lounev V, et al., 2016. Cellular hypoxia promotes heterotopic ossification by amplifying bmp signaling. *J Bone Miner Res*, 31(9):1652-1665. <https://doi.org/10.1002/jbmr.2848>
- Wang H, De Cunto CL, Pignolo RJ, et al., 2021a. Spatial patterns of heterotopic ossification in fibrodysplasia ossificans progressiva correlate with anatomic temperature gradients. *Bone*, 149:115978. <https://doi.org/10.1016/j.bone.2021.115978>
- Wang H, Zhang Q, Kaplan FS, et al., 2021b. Clearance of senescent cells from injured muscle abrogates heterotopic ossification in mouse models of fibrodysplasia ossificans progressiva. *Journal of Bone and Mineral Research*, 37(1):95-107. <https://doi.org/10.1002/jbmr.4458>
- Ware AD, Brewer N, Meyers C, et al., 2019. Differential vascularity in genetic and nonhereditary heterotopic ossification. *International Journal of Surgical Pathology*, 27(8):859-867. <https://doi.org/10.1177/1066896919857135>
- Wentworth KL, Masharani U, Hsiao EC, 2019. Therapeutic advances for blocking heterotopic ossification in fibrodysplasia ossificans progressiva. *Br J Clin Pharmacol*, 85(6):1180-1187. <https://doi.org/10.1111/bcp.13823>
- Wentworth KL, Lalonde RL, Groppe JC, et al., 2022. Functional testing of bone morphogenetic protein (bmp) pathway variants identified on whole - exome sequencing in a patient with delayed - onset fibrodysplasia ossificans progressiva (fop) using acvr1r206h - specific human cellular and zebrafish models. *Journal of Bone and Mineral Research*, 37(11):2058-2076. <https://doi.org/10.1002/jbmr.4711>
- Weston AD, Hoffman LM, Underhill TM, 2003. Revisiting the role of retinoid signaling in skeletal development. *Birth Defects Res C Embryo Today*, 69(2):156-173. <https://doi.org/10.1002/bdrc.10010>
- Williams E, Bagarova J, Kerr G, et al., 2021. Saracatinib is an efficacious clinical candidate for fibrodysplasia ossificans progressiva. *JCI Insight*, 6(8) <https://doi.org/10.1172/jci.insight.95042>
- Witten MR, Wu L, Lai C-T, et al., 2022. Inhibition of alk2 with bicyclic pyridyllactams. *Bioorganic & Medicinal Chemistry Letters*, 55:128452. <https://doi.org/10.1016/j.bmcl.2021.128452>
- Xiong H, Veedu RN, Diermeier SD, 2021. Recent advances in oligonucleotide therapeutics in oncology. *International Journal of Molecular Sciences*, 22(7):3295. <https://doi.org/10.3390/ijms22073295>
- Yamamoto H, Sakai N, Ohte S, et al., 2021. Novel bicyclic pyrazoles as potent alk2 (r206h) inhibitors for the treatment of fibrodysplasia ossificans progressiva. *Bioorg Med Chem Lett*, 38:127858. <https://doi.org/10.1016/j.bmcl.2021.127858>
- Yamamoto M, Stoessel SJ, Yamamoto S, et al., 2022. Overexpression of wild - type acvr1 in fibrodysplasia ossificans progressiva mice rescues perinatal lethality and inhibits heterotopic ossification. *Journal of Bone and Mineral Research*, 37(11):2077-2093. <https://doi.org/10.1002/jbmr.4617>
- Yang J, Kitami M, Pan H, et al., 2021. Augmented bmp signaling commits cranial neural crest cells to a chondrogenic fate by suppressing autophagic β -catenin degradation. *Science Signaling*, 14(665):eaaz9368. <https://doi.org/doi:10.1126/scisignal.aaz9368>
- Yang Y-S, Kim J-M, Xie J, et al., 2022. Suppression of heterotopic ossification in fibrodysplasia ossificans progressiva using aav gene delivery. *Nature Communications*, 13(1) <https://doi.org/10.1038/s41467-022-33956-9>
- Yang Y-S, Lin C, Ma H, et al., 2023. Aav-mediated targeting of the activin a-acvr1r206h signaling in fibrodysplasia ossificans progressiva. *Biomolecules*, 13(9):1364. <https://doi.org/10.3390/biom13091364>
- Ye Z, Wang S, Shan C, et al., 2023. The serum levels of activin a and bone morphogenetic protein-4 and -6 in patients with fibrodysplasia ossificans progressiva. *Orphanet Journal of Rare Diseases*, 18(1) <https://doi.org/10.1186/s13023-023-02708-3>
- Yu PB, Deng DY, Lai CS, et al., 2008a. Bmp type i receptor inhibition reduces heterotopic ossification. *Nature Medicine*, 14(12):1363-1369. <https://doi.org/10.1038/nm.1888>
- Yu PB, Hong CC, Sachidanandan C, et al., 2008b. Dorsomorphin inhibits bmp signals required for embryogenesis and iron metabolism. *Nat Chem Biol*, 4(1):33-41. <https://doi.org/10.1038/nchembio.2007.54>

Zhang YE, 2017. Non-smad signaling pathways of the tgf-beta family. *Cold Spring Harb Perspect Biol*, 9(2) <https://doi.org/10.1101/cshperspect.a022129>

Zitvogel L, Rusakiewicz S, Routy B, et al., 2016. Immunological off-target effects of imatinib. *Nat Rev Clin Oncol*, 13(7):431-446. <https://doi.org/10.1038/nrclinonc.2016.41>




Fig. 1 Schematic overview of currently known FOP mechanisms (created with BioRender.com) .




Fig. 2 Schematic overview of the drugs and compounds currently used and/or evaluated in the treatment of FOP, and their respective effects (created with BioRender.com)