



Research Article

<https://doi.org/10.1631/jzus.B2500477>



IL-15 aggravates cardiac ischemic injury via impairing macrophage efferocytosis and driving inflammation

Lei GUO^{1,2,3,4*}, Zhehui YIN^{1,2,3*}, Ning ZHANG⁵, Han CHEN^{1,2,3}, Zhuo WANG^{1,2,3}, Yuxue HUANG⁶, Jiniu HUANG^{1,2,3}, Yayu YOU⁷, Chenyun ZHANG^{1,2,3}, Qinyi BAO^{1,2,3}, Shuxin LEI^{1,2,3}, Jun JIANG^{1,2,3}, Xiaojie XIE^{1,2,3}

¹Department of Cardiology, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310009, China

²State Key Laboratory of Transvascular Implantation Devices, Hangzhou 310009, China

³Heart Regeneration and Repair Key Laboratory of Zhejiang Province, Hangzhou 310009, China

⁴Department of General Practice and International Medicine, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310009, China

⁵Department of Cardiology, Hangzhou First Affiliated People's Hospital, Westlake University School of Medicine, Hangzhou 310003, China

⁶Department of Cardiology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China

⁷Department of Cardiology, Hunan Provincial People's Hospital, Changsha 410000, China

Abstract: Acute myocardial infarction (AMI) remains a major global health burden and is characterized by profound cardiac inflammation, apoptotic cell death, and impaired myocardial function. While interleukin-15 (IL-15) has been implicated in immune regulation, its precise role in the pathogenesis of AMI has not been clarified. Therefore, this study sought to delineate the functional role of IL-15 in the progression of AMI, with a particular focus on its influence on macrophage-driven inflammation, efferocytosis, and metabolic reprogramming. IL-15 levels were assessed in AMI patients and murine models. To evaluate the impact of IL-15 on cardiac inflammation, apoptosis, and functional outcomes following AMI, *IL-15* and *IL-15* receptor α (*IL-15R α*) knockout (KO) mouse models were employed. Mechanistic studies were conducted to investigate IL-15-mediated effects on macrophage efferocytosis, polarization, and metabolic remodeling, with an emphasis on nuclear factor- κ B (NF- κ B) signaling and glycolytic flux. Elevated IL-15 levels were detected in both the plasma of AMI patients and the cardiac tissues of murine AMI models, correlating with increased disease severity. The genetic deletion of *IL-15* or *IL-15R α* significantly ameliorated cardiac injury by reducing inflammation and apoptosis while preserving myocardial function. Mechanistic analyses revealed that IL-15 impaired macrophage efferocytosis via Mer tyrosine kinase (MERTK) downregulation and promoted M1 polarization via NF- κ B pathway activation. Furthermore, IL-15 reprogrammed macrophage metabolism by enhancing glycolytic activity. Ultimately, IL-15 restoration exacerbated cardiac ischemic injury following AMI, serving as a critical regulator of macrophage-mediated inflammation in AMI. These findings highlight the role of IL-15 as a potential therapeutic and prognostic target for mitigating cardiac inflammation and improving myocardial recovery in AMI.

Key words: Myocardial infarction; Inflammatory phenotype; Macrophage function; Metabolic reprogramming; Interleukin-15 (IL-15)

1 Introduction

Acute myocardial infarction (AMI), as a leading cause of global morbidity and mortality, is primarily

caused by coronary artery occlusion, leading to myocardial ischemia and infarction (Roth et al., 2020). Despite advancements in AMI management, the relevant global burden remains significant. There is accumulating evidence that inflammation plays a crucial role in the development of AMI, determining the infarct size and contributing to ventricular remodeling and clinical outcomes (Westman et al., 2016). Sterile inflammation secondary to AMI is characterized by the recruitment and activation of innate and adaptive immune cells, which might facilitate the early complication of cardiac rupture (Li et al., 2022). The process by which macrophages eliminate apoptotic cells, known as efferocytosis, plays a crucial role in averting

✉ Xiaojie XIE, xiexj@zju.edu.cn

Jun JIANG, 2106002@zju.edu.cn

* The two authors contributed equally to this work

Xiaojie XIE, <https://orcid.org/0000-0001-8421-2737>

Jun JIANG, <https://orcid.org/0000-0001-6926-9516>

Lei GUO, <https://orcid.org/0000-0001-5934-018X>

Zhehui YIN, <https://orcid.org/0009-0003-8849-7858>

Received Aug. 9, 2025; Revision accepted Nov. 15, 2025;
Crosschecked June 29, 2026; Published online July 3, 2026

© Zhejiang University Press 2026

necrosis and inflammatory responses while simultaneously initiating pro-resolving pathways that promote ongoing efferocytosis (Ngai et al., 2023). However, the precise mechanisms underlying AMI pathogenesis remain unclear, necessitating further investigations to identify novel therapeutic targets that can mitigate AMI-associated mortality and improve clinical prognosis.

Interleukin-15 (IL-15), a member of the IL-2 cytokine family, is a 14–15 kDa protein comprising 114 amino acids, with a 4- α -helical bundle structure (Allard-Chamard et al., 2020). As a pleiotropic cytokine, IL-15 is widely expressed by diverse cell types, including monocytes, macrophages, dendritic cells, fibroblasts, epithelial cells, and skeletal muscle cells (Mortier et al., 2008; Perera et al., 2012; Patidar et al., 2016). Its expression is regulated at three levels: transcriptional, translational, and secretory. IL-15 exerts its biological effects on the inflammatory response and immunoregulation mainly through binding to its specific IL-15 receptor (IL-15R), which is composed of α , β , and γ chains. IL-15 shares the receptor components IL-15R β and IL-15R γ with IL-2, while the unique IL-15R α subunit specifically binds IL-15, conferring ligand specificity. During intracellular signaling, IL-15 binds to the trimer and exerts its activity primarily through the α chain (IL-15R α or cluster of differentiation 215 (CD215)) (Mortier et al., 2004).

Existing reports on the effects of IL-15 and its specific receptor on AMI are limited. Genetic variations of IL-15 and higher serum IL-15 levels have been found in patients with either acute coronary syndrome or ischemic stable coronary heart disease (Gokkusu et al., 2010). It was also suggested that epicardial adipose tissue may be a potential source of both IL-15 and IL-15R α . Furthermore, previous studies have demonstrated that IL-15 is prominently expressed in atherosclerotic lesions of hyperlipidemic mouse aortas and human carotid arteries (Wuttge et al., 2001; Dozio et al., 2014). IL-15 can regulate macrophages in the spinal cord and sciatic nerve after chronic constriction injury, and its expression is also regarded as a signal for the activation of macrophages in the sciatic nerve (Gómez-Nicola et al., 2008). However, the precise roles of IL-15 in the macrophage efferocytosis process during AMI and the underlying mechanisms remain unclear. Based on these previous reports, we hypothesized that IL-15, as a pro-inflammatory cytokine, might aggravate cardiac

ischemic injury by specifically impairing macrophage efferocytosis and driving inflammation.

This study utilized *IL-15* and *IL-15R α* knockout (KO) mice as sophisticated genetic tools to delineate the specific contributions of IL-15 to the pathophysiology of AMI. We investigated the molecular mechanisms through which IL-15 contributes to AMI pathogenesis, with the objective of providing mechanistic insights that could guide the development of innovative therapeutic strategies to enhance AMI outcomes.

2 Materials and methods

The study methods are extensively described in the supplementary materials and methods.

2.1 Human plasma samples

This investigation was performed according to the Declaration of Helsinki. The study protocol was approved by the Ethics Committee of The Second Affiliated Hospital of Zhejiang University School of Medicine (No. 2021-0624), and written informed consent was obtained from all participants prior to inclusion. A total of 243 patients who complained of chest pain or distress and underwent coronary angiography at The Second Affiliated Hospital of Zhejiang University School of Medicine from 2018 to 2022 were enrolled. Among them, 121 patients with 100% coronary artery occlusion were classified as the AMI group, while 122 patients with coronary myocardial bridge (CMB) were assigned to the healthy control (CTL) group. The participants' characteristics are listed in Table S1. No significant differences in age, gender, or smoking history were observed between the two groups. Blood samples were collected during coronary angiography and stored for further tests.

2.2 Animals

Male C57BL/6J mice aged 8–10 weeks and neonatal C57BL/6J mice were purchased from Shanghai Slac Laboratory Animal Technology Corporation (Shanghai, China). Global *IL-15* KO and *IL-15R α* KO mice (all with C57BL/6J background) were purchased from Shanghai Model Organisms (Shanghai, China). Mice were housed at room temperature ((20 $\pm$ 2) °C) with an alternating 12/12 h light/dark cycle and provided with a standard diet. All animal procedures conformed to the

guidelines from Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes, and the study protocol was approved by the Ethical Committee of The Second Affiliated Hospital of Zhejiang University School of Medicine (No. AIRB-2021-1012).

2.3 AMI model

Mice were put under 2.0% (volume fraction) isoflurane (Sigma-Aldrich, USA) anesthesia and mechanical ventilation (Alcott Biotech, Shanghai, China) assisted by tracheotomy, and then subjected to AMI by the ligation of left anterior descending artery. Mice in the sham group underwent thoracotomy without the ligation of coronary arteries. The animals were placed on a heating pad and watched carefully for the next 3 h until they achieved full locomotor behavior. To investigate whether the inhibition of IL-15 protected myocardium from acute ischemic injury, we established AMI in *IL-15* KO and *IL-15Rα* KO mice. To validate the detrimental effect of IL-15 in acute cardiac ischemic injury, *IL-15* KO mice and wild-type (WT) littermates were used and intraperitoneally injected either with recombinant murine IL-15 (rmIL-15, 50 µg/kg, PeproTech, USA) or phosphate-buffered saline (PBS) daily for 14 d after AMI establishment. At the end of the experiment, euthanasia was performed using 2% inhaled isoflurane followed by cervical dislocation.

3 Results

3.1 IL-15 is elevated in AMI patient plasma and murine hearts

To investigate the potential role of IL-15 in AMI, we initially collected plasma samples from patients diagnosed with AMI or CTL subjects. Plasma IL-15 levels were significantly elevated in AMI patients compared to those in CTL subjects (Fig. 1a), and this association remained significant (adjusted odds ratio (OR)=1.320, 95% confidence interval (CI): 1.130–1.542, $P<0.001$; Table S2) after adjustment for body mass index (BMI), hypertension, diabetes, and hyperlipidemia in a multivariable logistic regression model, indicating a potential association between IL-15 and the pathophysiological processes of AMI.

In order to further elucidate the temporal and spatial expression of IL-15, we employed a mouse model of AMI. Tissue lysates from the border zone of infarcted myocardium revealed a gradual increase in IL-15 levels, peaking on the third day post-AMI and subsequently declining (Fig. 1b). Consistently, western blot analysis revealed the same trend in the expression of IL-15 (Figs. 1c and 1d). We sought to identify the specific receptor subtype involved in mediating the interaction with IL-15 in this process. IL-15Rα expression was significantly upregulated in the myocardium following AMI, whereas the levels of IL-15Rβ and IL-15Rγ showed no significant changes (Figs. 1c and 1e–1g). Interestingly, although IL-15Rα expression was elevated in the remote myocardium, IL-15 levels were not significantly changed in this region (Figs. S1a–S1c). Immunofluorescence staining further demonstrated that IL-15 was predominantly localized intracellularly in the border zone of the infarcted myocardium in AMI mice (Figs. 1h and 1i) and confirmed a significant colocalization of IL-15 protein with the macrophage marker CD68 in the cardiac tissues of our AMI model, which suggests that macrophages are prominent producers of IL-15 (Fig. 1j). Collectively, these findings highlight the dynamic regulation of IL-15 and IL-15Rα and suggest a potential role for IL-15 in the pathophysiology of AMI.

3.2 *IL-15/IL-15Rα* deficiency reduces infarct size and improves cardiac function

To elucidate the roles of IL-15 in AMI, *IL-15* KO and *IL-15Rα* KO mice were generated using gene-editing techniques (Fig. S2) and subjected to AMI. Notably, 30% of WT littermates succumbed within one week of AMI, whereas most *IL-15* KO and *IL-15Rα* KO mice survived (Fig. 2a). Echocardiographic analyses revealed that *IL-15* KO and *IL-15Rα* KO mice exhibited significantly higher ejection fraction (EF) and fractional shortening (FS), along with reduced left ventricular internal dimension at end-diastole (LVIDd) compared to WT littermates after AMI, though these differences were not statistically significant at some time points (Figs. 2b and 2c, Table S3). Cardiac mass, as reflected by the heart-to-body weight ratio, was lower in *IL-15* KO and *IL-15Rα* KO mice than in WT littermates (Fig. 2d). Histological examination demonstrated that the infarct size was significantly smaller in *IL-15* KO and *IL-15Rα* KO mice on the third day




Fig. 1 Elevated interleukin-15 (IL-15) in acute myocardial infarction (AMI) patients' plasma and murine hearts. (a) Plasma IL-15 levels in the AMI patients ($n=121$) and the healthy control (CTL) group ($n=122$). (b) IL-15 levels in tissue lysates of the infarct border area of mouse myocardium on different days after AMI ($n=6$). (c–g) Expression of IL-15 and its receptors in the infarct border area of mouse myocardium on different days after AMI via western blot analysis ($n=5$). (h) Representative immunofluorescence staining of IL-15 (red), cardiac troponin I (cTnI) (green), and 4',6-diamidino-2-phenylindole (DAPI) (blue) in the infarct border area of mouse myocardium on the third day after AMI. Positive staining is indicated by white arrows. (i) Fluorescence intensity analysis of IL-15 in the infarct border area of mouse myocardium on the third day after AMI ($n=5$). (j) Representative immunofluorescence staining of IL-15 (red), cluster of differentiation 68 (CD68) (green), and DAPI (blue) in the infarct border area of mouse myocardium on the third day after AMI. Co-localization of IL-15 with CD68 is indicated by white arrows. Data are expressed as mean $\pm$ standard error of the mean (SEM). Statistical analyses were performed using GraphPad Prism v10.1.2. Significant differences between means were analyzed using the unpaired Student's t -test or one-way or two-way analysis of variance (ANOVA) followed by Bonferroni post-test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$. D0, D1, D2, D3, D7, and D14: Day 0, Day 1, Day 2, Day 3, Day 7, and Day 14.




Fig. 2 Effects of interleukin-15 (*IL-15*)/IL-15 receptor α (*IL-15R α*) deficiency on cardiac function and infarct size. (a) Kaplan-Meier survival analysis of *IL-15* knockout (KO) mice with acute myocardial infarction (AMI) ($n=25$) or sham ($n=12$), *IL-15R α* KO mice with AMI ($n=26$) or sham ($n=11$), and wild-type (WT) littermates with AMI ($n=52$) or sham ($n=12$). (b) Representative of M-mode echocardiographic images in each group. (c) Quantitative echocardiographic analysis of ejection fraction (EF), fractional shortening (FS), and left ventricular internal dimension at end-diastole (LVIDd) ($n=5-12$). (d) Quantitative analysis of cardiac mass index ($n=5$ or 6). (e) Representative images of a heart section after triphenyltetrazolium chloride (TTC) staining on the third day after AMI. The infarct size is circled by the dotted line. (f) Quantitative analysis as the percentage of infarct size to the whole heart ($n=6$). Data are expressed as mean $\pm$ standard error of the mean (SEM). * $P<0.05$, ** $P<0.01$, and *** $P<0.001$. D0, D3, D7, and D14: Day 0, Day 3, Day 7, and Day 14.

post-AMI compared to WT mice (Figs. 2e and 2f). Additionally, plasma B-type natriuretic peptide (BNP) levels, which progressively increased after AMI, were significantly lower in *IL-15* KO and *IL-15Ra* KO mice on the fourteenth day post-AMI compared to WT mice (Fig. S3). Taken together, *IL-15* and its receptor *IL-15Ra* are critical mediators in the pathological progression of AMI: the absence of *IL-15* or *IL-15Ra* improved survival, preserved cardiac function, and reduced myocardial damage, suggesting that IL-15 may contribute to cardiac dysfunction.

3.3 Deficiency of *IL-15* and *IL-15Ra* alleviates myocardial apoptosis and inflammation

Apoptosis, or programmed cell death, is a crucial mechanism in AMI pathophysiology. The percentage of terminal-deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL)-positive cells in the heart was markedly lower in *IL-15* KO and *IL-15Ra* KO mice than in WT littermates on the third day post-AMI (Figs. 3a and 3b). Western blot analysis further demonstrated reduced cleaved caspase-3 levels in the border zone of the infarcted myocardium in *IL-15* KO and *IL-15Ra* KO mice (Fig. 3c). To determine whether IL-15 directly influences cardiomyocyte apoptosis, an in vitro hypoxia model was established. Cardiomyocyte apoptosis was strongly induced by hypoxia; however, it was not exacerbated by preconditioning with rmIL-15, as evidenced by TUNEL staining, lactate dehydrogenase activity (Figs. S4a and S4b), and cleaved caspase-3 levels (Fig. 3d). These findings suggest that IL-15 contributes to hypoxia-mediated cardiomyocyte apoptosis, whereas its pro-apoptotic effects may not be directly mediated through cardiomyocytes.

Inflammation also plays a critical role in tissue damage and repair following myocardial infarction (MI). We therefore aimed to explore the contributions of IL-15 and IL-15Ra to inflammation-induced myocardial injury. On the third day post-AMI, the messenger RNA (mRNA) levels of pro-inflammatory cytokines *IL-1 β* and *IL-6* in the border zone of the infarcted myocardium were reduced in *IL-15* KO and *IL-15Ra* KO mice compared to WT littermates. Conversely, the expression of the anti-inflammatory cytokines *IL-4* and *IL-10* was upregulated in these KO mice, indicating anti-inflammatory roles of *IL-15* and *IL-15Ra* deficiency (Fig. 3e). Considering that IL-15 exerts its biological functions through IL-15Ra, we next performed

immunofluorescence staining to localize IL-15Ra in the ischemic myocardium. IL-15Ra was predominantly colocalized with macrophages (e.g., CD68⁺) compared to other cell types (Figs. 3f, 3g, and S4c). Moreover, the number of CD68⁺ macrophages was markedly reduced in the cardiac tissues of *IL-15* KO and *IL-15Ra* KO mice following AMI (Figs. 3h and 3i). These findings suggest that the interaction between IL-15 and IL-15Ra may serve as a critical driver of macrophage-mediated cardiomyocyte apoptosis during AMI.

3.4 IL-15 impairs macrophage efferocytosis by suppressing MERTK expression

Efferocytosis, the process by which phagocytic cells such as macrophages recognize and engulf apoptotic cells, is essential for maintaining tissue homeostasis and resolving inflammation. We hypothesized that IL-15 exacerbates apoptotic cardiomyocyte accumulation by impairing macrophage efferocytosis. Therefore, bone marrow-derived macrophages (BMDMs) were treated with rmIL-15 and their phagocytic activity was assessed. Treatment with rmIL-15 significantly reduced macrophage phagocytosis (Figs. 4a and 4b). To confirm these effects in vitro, CD68-stained BMDMs were co-cultured with PKH26-stained apoptotic mouse cardiomyocytes. Flow cytometry revealed a markedly reduced phagocytic rate in rmIL-15-treated macrophages compared to controls (Figs. 4c and 4d), which was further validated by immunofluorescence imaging (Figs. 4e and 4f). To elucidate the relevant mechanism, we examined the expression of genes associated with macrophage efferocytosis. Mer tyrosine kinase (MERTK) is a receptor with a critical role in recognizing and clearing apoptotic cells. Treatment with rmIL-15 markedly suppressed MERTK expression in macrophages at both the transcriptional and protein expression levels (Figs. 4g and 4h). Our findings strongly suggest that IL-15 suppressed efferocytosis by down-regulating MERTK. To definitively establish this causal relationship, we performed a crucial rescue experiment. As anticipated, IL-15 treatment potently suppressed the endogenous MERTK protein levels in Lenti-Vector cells. In stark contrast, the Lenti-MERTK cells robustly maintained high levels of MERTK expression, successfully overriding the inhibitory signal from IL-15 (Fig. 4i). Most critically, we assessed the functional consequences. In Lenti-Vector cells, IL-15 treatment led to a profound suppression of efferocytosis,




Fig. 3 Effects of interleukin-15 (*IL-15*) and IL-15 receptor α (*IL-15R α*) deficiency on myocardial apoptosis and inflammation. (a) Representative images of heart tissue by terminal-deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) staining on the third day after acute myocardial infarction (AMI). TUNEL-positive cells are indicated by white arrows. (b) Quantitative analysis of TUNEL-positive cells ($n=5$ or 6). (c) Expression of caspase-3 and cleaved caspase-3 in the infarct border area on the third day after AMI. (d) Expression of caspase-3 and cleaved caspase-3 in primary cardiomyocytes under normoxia or hypoxia, preconditioned with recombinant murine IL-15 (rmIL-15) or vehicle. (e) Expression of genes in the infarct border zone on the third day after AMI or sham ($n=3$). (f, g) Representative immunofluorescence staining of IL-15R α , cluster of differentiation 68 (CD68), and 4',6-diamidino-2-phenylindole (DAPI) in the heart section of wild-type (WT) mice (positively stained cells are indicated by white arrows), and quantitative analysis of positively co-stained cells per slice ($n=5$ or 6). (h, i) Representative immunofluorescence staining of CD68, cardiac troponin I (cTnI), and DAPI in the heart section on the third day after AMI, and quantitative analysis of positively co-stained cells per slice (WT: $n=7$; *IL-15* KO: $n=10$; *IL-15R α* KO: $n=6$). Data are expressed as mean $\pm$ standard error of the mean (SEM). * $P<0.05$, ** $P<0.01$, and *** $P<0.001$. KO: knockout; mRNA: messenger RNA; MPO: mouse anti-myeloperoxidase; D0, D3: Day 0, Day 3.




Fig. 4 Interleukin-15 (IL-15)-mediated impairment of macrophage efferocytosis through Mer tyrosine kinase (MERTK) suppression. (a, b) Flow cytometry analysis of phagocytic activity in bone marrow-derived macrophages (BMDMs) treated with recombinant murine IL-15 (rmIL-15) or phosphate-buffered saline (PBS), and quantification of phagocytosis activity, expressed as the percentage of phycoerythrin (PE)-positive macrophages. (c, d) Flow cytometry analysis of efferocytosis, measured as the uptake of PKH26-stained apoptotic cardiomyocytes by cluster of differentiation 68-positive (CD68⁺) macrophages, and quantitative analysis of efferocytosis activity. (e, f) Representative immunofluorescence images of BMDMs co-cultured with apoptotic cardiomyocytes. CD68 (green), PKH26 (red), and 4',6-diamidino-2-phenylindole (DAPI) (blue), and quantitative analysis of positively co-stained cells per slice. (g) Messenger RNA (mRNA) levels of efferocytosis-related genes. (h) Western blot analysis of MERTK protein expression. (i) Representative western blot analysis of MERTK protein expression in Lenti-Vector or Lenti-MERTK RAW 264.7 cells treated with PBS or rmIL-15. (j, k) Flow cytometry analysis of efferocytosis, measured as the uptake of PKH26-stained cardiomyocytes by Lenti-Vector or Lenti-MERTK RAW 264.7 cells treated with PBS or rmIL-15, and quantitative analysis of efferocytosis activity. Data are expressed as mean $\pm$ standard error of the mean (SEM) ($n=3$). * $P<0.05$ and ** $P<0.01$; ns: no significance. M-CSF: macrophage-colony stimulating factor; SSC-A: side scatter-area.

consistent with our findings in primary macrophages. However, Lenti-*MERTK* cells were rendered completely resistant to this IL-15-mediated functional defect. Despite the presence of IL-15, Lenti-*MERTK* cells maintained a high level of efferocytic capacity comparable to that of untreated controls (Figs. 4j and 4k). Collectively, these data unequivocally demonstrate that downregulation of *MERTK* is the direct and rate-limiting mechanism by which IL-15 impairs macrophage efferocytosis.

3.5 IL-15 drives macrophage M1 polarization and activates NF- κ B signaling

Beyond efferocytosis, the polarization of macrophages between the pro-inflammatory M1 and the anti-inflammatory M2 states critically influences the inflammatory microenvironment and overall cardiac outcomes. Thus, we sought to determine whether IL-15 affects macrophage polarization and the signaling pathways involved. We therefore assessed the expression of M1 and M2 markers in *IL-15* KO and *IL-15R α* KO mice. CD206⁺ macrophages (M2 phenotype) were notably increased in both KO models, while inducible nitric oxide synthase-positive (iNOS⁺) macrophages (M1 phenotype) were significantly reduced (Figs. 5a and 5b). These observations were confirmed in ex vivo experiments, where macrophages treated with rmIL-15 exhibited elevated levels of M1-associated cytokines (IL-1 α , IL-1 β , IL-6, interferon- γ (IFN- γ), and tumor necrosis factor- α (TNF- α)) and reduced levels of M2-associated cytokines (IL-4 and IL-10) in the culture supernatants. Other cytokines such as monocyte chemoattractant protein-1 (MCP-1), granulocyte-colony stimulating factor (G-CSF), granulocyte macrophage-colony stimulating factor (GM-CSF), IL-12, and IL-17A showed a significant increase in levels after treatment. There are also some cytokines such as IL-2, IL-3, IL-5, IL-8, and Eotaxin that did not show significant changes after the treatment (Figs. 5c and S5).

Next, we investigated the involvement of the nuclear factor- κ B (NF- κ B) signaling pathway, the classical pathway related to inflammatory responses, in IL-15-induced macrophage polarization. Under baseline conditions, p65, a key component of the NF- κ B pathway, was primarily localized in the cytoplasm of CD68⁺ macrophages. However, following rmIL-15 treatment, p65 translocated from the cytoplasm to the nucleus (Fig. 5d), indicating NF- κ B activation. To confirm the role of this pathway, gossypin, an NF- κ B inhibitor,

was used to treat BMDMs alongside rmIL-15 at varying concentrations (Fig. 5e). Treatment with rmIL-15 significantly increased the transcriptional levels of M1-related genes (*IL-1 α* , *IL-1 β* , and *IL-6*) and decreased the levels of M2-related genes (*IL-4* and *IL-10*). These effects were reversed by gossypin, with the most pronounced inhibition observed at 50 μ mol/L (Fig. 5f). To test the IL-15-induced interaction between macrophages and cardiomyocytes and the effect of the inhibitor on this interaction, primary cardiomyocytes were treated with the supernatant of IL-15-treated macrophages. The supernatant of macrophages treated with rmIL-15 increased the ratio of cleaved caspase-3/caspase-3 in cardiomyocytes under the hypoxic condition, which could be reduced by gossypin (Fig. 5g). These findings suggest that IL-15 promoted the pro-inflammatory M1 polarization of macrophages via the activation of NF- κ B signaling pathway, increasing the secretion of proinflammatory cytokines, which amplified cardiomyocyte apoptosis.

3.6 IL-15 enhances glycolytic activity and metabolic activation in macrophages

Building on the observed effects of IL-15 on macrophage inflammatory responses and efferocytosis, we next investigated whether IL-15 modulates macrophage function via influencing metabolism. Glycolysis, a primary energy source for macrophage activation and inflammatory signaling, was first assessed through intracellular and extracellular lactate measurements. Treatment with rmIL-15 significantly increased lactate concentrations both extracellularly and intracellularly in macrophages (Figs. 6a and 6b). To further explore this metabolic shift, we analyzed glycolysis-related gene expression. IL-15 treatment induced a pronounced up-regulation of glycolytic genes, including phosphoglycerate kinase 1 (*PGK1*) and enolase 1 (*ENO1*), highlighting its role in promoting glycolytic reprogramming (Fig. 6c). *PGK1* and *ENO1* are key glycolytic enzymes that catalyze critical steps in the glycolytic pathway, directly contributing to adenosine triphosphate (ATP) production via substrate-level phosphorylation. Cellular ATP levels were also markedly elevated in IL-15-treated macrophages, indicating enhanced metabolic activity (Fig. 6d). These findings suggest that IL-15 promoted glycolytic reprogramming in macrophages, characterized by increased lactate production and ATP generation, thereby contributing to their activation and inflammatory function.




Fig. 5 Interleukin-15 (IL-15)-driven macrophage M1 polarization and nuclear factor- κ B (NF- κ B) signaling activation. (a) Representative immunofluorescence staining of cluster of differentiation 68 (CD68), inducible nitric oxide synthase (iNOS), CD206, and 4',6-diamidino-2-phenylindole (DAPI) in the heart section of mice on the third day after acute myocardial infarction (AMI). Positively stained cells are indicated by white arrows. (b) Quantitative analysis of positively co-stained cells per slice ($n=5$). (c) Levels of IL-1 α , IL-1 β , IL-6, tumor necrosis factor- α (TNF- α), interferon- γ (IFN- γ), monocyte chemoattractant protein-1 (MCP-1), granulocyte colony-stimulating factor (G-CSF), IL-4, and IL-10 in the supernatant of macrophages by protein microarray assays, incubated with either recombinant murine IL-15 (rmIL-15) (50 ng/mL) or phosphate-buffered saline (PBS) ($n=4$). (d) Representative immunofluorescence staining of CD68, NF- κ B p65, and DAPI in cultured bone marrow-derived macrophages (BMDMs). (e) Expression of NF- κ B p65, phosphorylated p65 (p-p65), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and histone H3 by western blot in BMDMs incubated with rmIL-15 (50 ng/mL) or/and gossypin (10 and 50 μ mol/L). (f) Expression of *IL-1 α* , *IL-1 β* , *IL-6*, *IL-4*, and *IL-10* messenger RNAs (mRNAs) by quantitative polymerase chain reaction (qPCR) in cultured BMDMs, incubated with rmIL-15 (50 ng/mL) and/or gossypin (10 and 50 μ mol/L) ($n=3$). (g) Expression of caspase-3, cleaved caspase-3, and β -tubulin in primary cardiomyocytes after treatment with the supernatant of macrophages exposed to rmIL-15 (50 ng/mL) and gossypin (10 and 50 μ mol/L). Data are expressed as mean $\pm$ standard error of the mean (SEM). * $P<0.05$, ** $P<0.01$, and *** $P<0.001$. WT: wild-type; KO: knockout; M-CSF: macrophage-colony stimulating factor; DMSO: dimethyl sulfoxide; D0, D3: Day 0, Day 3.




Fig. 6 Interleukin-15 (IL-15) enhancement of glycolytic activity and metabolic activation in macrophages. (a, b) Levels of extracellular and intracellular lactates in macrophages measured by enzyme-linked immunosorbent assay (ELISA) kit, incubated with recombinant murine IL-15 (rmIL-15) (50 ng/mL) or phosphate-buffered saline (PBS). (c) Messenger RNA (mRNA) levels of glycolysis-related genes. (d) Levels of cellular adenosine triphosphate (ATP) in macrophages measured by ELISA kit. Data are expressed as mean±standard error of the mean (SEM) ($n=3$). * $P<0.05$, ** $P<0.01$, *** $P<0.001$, and **** $P<0.0001$.

3.7 Restoring IL-15 exacerbates cardiac ischemia injury after AMI

To confirm and validate the adverse effects of IL-15, exogenous rmIL-15 or vehicle was administered intraperitoneally to *IL-15* KO and WT mice once daily for 14 d following AMI. The administration of rmIL-15 significantly increased mortality in both WT and *IL-15* KO mice (Fig. 7a). Furthermore, rmIL-15 treatment aggravated cardiac dysfunction in WT mice, as evidenced by reduced EF and FS, along with LVIDd expansion post-AMI. Similarly, reconstitution of rmIL-15 in *IL-15* KO mice resulted in impaired EF and FS, although it did not significantly affect LVIDd (Figs. 7b and 7c, Table S4). Moreover, rmIL-15 administration increased the cardiac mass index and resulted in larger infarct sizes in both WT and *IL-15* KO mice (Figs. 7d–7f). These findings indicate that restoring IL-15 exacerbates cardiac dysfunction and myocardial injury following AMI, further supporting its detrimental role in AMI pathogenesis.

4 Discussion

This study identified IL-15 as a pivotal regulator in the early-phase pathogenesis of AMI. IL-15 was

significantly upregulated in the plasma of AMI patients and the ischemic myocardium of AMI mice, as an independent risk factor for the occurrence of AMI. IL-15R α , as the receptor for IL-15, was shown to be a key mediator in driving macrophage polarization toward the pro-inflammatory M1 phenotype, exacerbating inflammation, impairing efferocytosis, and increasing myocardial apoptosis, which ultimately contributed to adverse cardiac remodeling and heart failure. These effects were mediated through the activation of NF- κ B signaling pathway, which underscores the role of IL-15 as a driver of macrophage pro-inflammatory responses. Beyond these effects, IL-15 was found to modulate macrophage glycolysis, enhancing metabolic activation to support inflammation and other energy-intensive processes. Hence, these new findings allowed us to reach the potential mechanisms discussed below.

Notably, our findings underscore the critical role of IL-15 in the early pathogenesis of AMI following coronary occlusion, establishing IL-15 as a compelling therapeutic target for intervention. AMI is fundamentally an inflammatory condition, as supported by evidence showing that targeting pro-inflammatory cytokines reduces recurrent cardiovascular events in AMI patients (Ridker et al., 2017). A variety of immune cells




Fig. 7 Interleukin-15 (IL-15)-mediated aggravation of cardiac ischemic injury after acute myocardial infarction (AMI). (a) Kaplan-Meier survival analysis after AMI of *IL-15* knockout (KO) mice administrated rmIL-15 (50 μ g/kg, $n=21$) or vehicle (phosphate-buffered saline (PBS), $n=21$) and wild-type (WT) littermates administrated rmIL-15 (50 μ g/kg, $n=26$) or vehicle (PBS, $n=16$). (b) Representative images of M-mode echocardiography in each group. (c) Quantitative echocardiographic analysis of ejection fraction (EF), fractional shortening (FS), and left ventricular internal dimension at end-diastole (LVIDd) ($n=5-8$). (d) Quantitative echocardiographic analysis of cardiac mass index ($n=5$ or 6). (e) Representative images of heart section by triphenyltetrazolium chloride (TTC) staining on the third day after AMI. The infarct size is circled by the dotted line. (f) Quantitative analysis as the percentage of infarct size to the whole heart ($n=6$). Data are expressed as mean $\pm$ standard error of the mean (SEM). * $P<0.05$, ** $P<0.01$, and *** $P<0.001$. D0, D3, D7, and D14: Day 0, Day 3, Day 7, and Day 14.

constitutively express cytokines in the acute phase of AMI, exerting multifaceted biological roles and contributing to inflammation status in the ischemic microenvironment. If effective therapies are not promptly applied after coronary occlusion, severe complications may occur, such as malignant arrhythmia, heart failure, mitral valve insufficiency, and even cardiac rupture. Even though the incidence of cardiac rupture is relatively rare at 1% to 3%, it is almost always fatal as the mortality is nearly up to 100% once it occurs (Hao et al., 2020). Therefore, it has very high clinical significance to investigate the pathogenic mechanisms in the early phase of AMI incidence and figure out the potential targets. Our research demonstrated that the content of human plasma IL-15 still has an independent association with AMI incidence after adjusting for risk factors such as BMI, hypertension, diabetes, and hyperlipidemia.

It is becoming increasingly well recognized that interleukins are involved in the development and early-phase remodeling of AMI. IL-4 and IL-10 have been proved to improve early-phase remodeling by modulating the balance between anti-inflammation and pro-inflammation, whereas IL-17A and IL-32 play pathogenic roles in the inflammation cascade (Zhou et al., 2014; Jung et al., 2017; Xuan et al., 2017; Daseke et al., 2020). IL-33 exhibits biphasic regulatory effects, facilitating significant inflammatory resolution by the fourth day post-AMI while suppressing prolonged inflammation between Days 4 and 7. However, these effects ultimately result in exacerbated cardiac remodeling and an increased risk of cardiac rupture (Ghali et al., 2020). Different from sharing the $\beta\gamma$ signaling complex with other interleukins (Patidar et al., 2016), IL-15R α binds IL-15 and anchors it to the cell surface, facilitating its presentation and stability in the extracellular microenvironment (Wuttge et al., 2001). This supports our findings that the expression of IL-15R α in the infarct border area was upregulated, but no significant change occurred in the expression of either IL-15R β or IL-15R γ . A previous study has demonstrated that IL-15R α signaling exerts opposing regulatory effects on IL-17-producing $\gamma\delta$ T cells (Colpitts et al., 2015). In the present study, we constructed both *IL-15* KO and its specific receptor *IL-15R α* KO mice and investigated their roles in the early phase after AMI, which might have greater clinical translational significance for reducing the risk of heart failure after AMI.

Accumulating evidence underscores the diverse and critical roles of immune cells, including neutrophils, macrophages, eosinophils, mast cells, and dendritic cells, in modulating the ischemic and hypoxic microenvironment (Daseke et al., 2020; Ghali et al., 2020; Wei et al., 2020). It has been well-documented that macrophages are involved in ventricular remodeling after AMI (Lambert et al., 2008; Hamada et al., 2014; Jakubzick et al., 2017). The infiltration of macrophages begins on the first day and peaks on the 3rd–5th day post-AMI (Yan et al., 2013). The early intervention of inflammation after ischemia may protect cardiomyocytes against ischemic injury and improve cardiomyocytes survival in the border zone of the infarcted myocardium. In the present study, we noticed that the expression of IL-15 in the border area of infarcted myocardium increased gradually, peaked on the third day post-AMI and then decreased thereafter.

An increasing body of evidence underscores that controlling the macrophage M1/M2 phenotype may represent a novel therapeutic strategy for treating AMI (van den Bossche et al., 2016). It has been indicated that promoting macrophage M2 polarization through the exogenous administration of IL-10 leads to the activation of cardiac fibroblasts, a reduction in microenvironmental inflammation, and accelerated wound healing following AMI (Jung et al., 2017). In the current study, the peak of IL-15 expression was highly consistent with the presence of M1 phenotype macrophages. Also, exogenous IL-15 administration promoted the M1 polarization of macrophages, whereas the absence of either IL-15 or IL-15R α inhibited macrophage infiltration and M1 polarization during hypoxia. These results imply the importance of IL-15 signaling in macrophage polarization after AMI.

Targeting the IL-15 signaling pathway could be beneficial in addressing many clinical problems. NF- κ B is a key regulator in the inflammatory response. It has been reported that IL-15 triggers NF- κ B activation in human neutrophils (McDonald et al., 1998). Furthermore, NF- κ B might be responsible for IL-15 actions on brain microvessel endothelial cells, thus affecting cellular permeability, endocytosis, and intracellular trafficking at the level of the blood–brain barrier (Stone et al., 2011). Consistent with our study, some studies have demonstrated that the downregulation of phosphorylated NF- κ B p65 expression led to a decrease in Ly6C^{low} macrophages in the heart (indicated as M1

macrophage), ameliorating isoprenaline-induced cardiac hypertrophy and fibrosis (Li et al., 2020). Wang et al. (2023) also showed that the high mobility group box 1 (HMGB1)/NF- κ B pathway regulates macrophage M2 polarization and alleviates AMI by impeding inflammation, oxidation, apoptosis, and autophagy. In the present work, we further revealed that IL-15 activated the NF- κ B signaling pathway in macrophages during ischemic injury by enhancing intracellular NF- κ B translocation and phosphorylation.

Cellular energy metabolism and ferroptosis were particularly important during the process of cardiomyocyte death (Luan et al., 2024). Through the regulation of myocardial energetic metabolism, cinnaroside exhibits considerable promise in attenuating doxorubicin (DOX)-triggered oxidative stress, inflammatory cell demise, and nucleotide-binding oligomerization domain-like receptor protein 3 (NLRP3)-dependent cardiomyocyte pyroptosis (Zou et al., 2024). Existing literature suggests that immune-metabolic reprogramming is critically involved in shaping macrophage phenotypic transitions, providing a mechanistic link between cellular metabolism and immune function (van den Bossche et al., 2016). This underscores the potential of targeting metabolic pathways to modulate macrophage polarization and develop novel therapeutic strategies for AMI. Our findings demonstrated that IL-15 disrupted macrophage metabolic homeostasis by significantly upregulating the expression of key glycolytic enzymes, such as PGK1 and ENO1, and enhancing the production of downstream metabolites, including lactate and ATP. These results highlight IL-15 as a potential target for therapeutic intervention aimed at restoring metabolic balance and mitigating inflammation. In summary, our findings demonstrate that IL-15 treatment markedly amplified macrophage-mediated inflammation and glycolytic metabolism while impairing efferocytosis, underscoring the multifaceted role of IL-15 in orchestrating macrophage activation and functional reprogramming. These effects collectively exacerbated infarct expansion, heightened myocardial apoptosis, and impaired cardiac function, highlighting their detrimental impact on post-AMI cardiac pathology. Our study focused on the acute inflammatory phase following MI. In contrast, Ameri et al. (2020) elegantly demonstrated a beneficial role of IL-15 in the late-phase remodeling stage. In their study, IL-15 treatment significantly increased vessel area density

and reduced fibrosis after MI. IL-15 is not the only cytokine that shows a rapid increase in expression in early-stage MI and exacerbates myocardial damage. However, when administered exogenously, it can actually reduce the infarcted area. In mice, TNF- α levels rose significantly on the first day post-AMI, and the deletion of TNF- α significantly improved myocardial function on the third day post-MI but not on the 7th day (Zhang et al., 2013). In male rats with left anterior descending artery ligation, TNF- α mRNA and protein production increased in the myocardium on the first day, which were detectable until the 35th day post-AMI (Irwin et al., 1999). In young rat hearts, the injection of TNF- α significantly reduced the extent of myocardial injury after coronary ligation. This protective effect may be related to its activation of the downstream platelet-derived growth factor-B (PDGF-B) pathway (Cai et al., 2003). These studies, combined with our own research, further suggest the complexity of the role of cytokines in the early and late stages of MI.

Despite the promising findings, our study has certain limitations. To investigate the exact roles of IL-15 in myocardial ischemia, we constructed *IL-15* KO and *IL-15Ra* KO mice and established an AMI model. The mice we used were global KO mice, not conditional KO mice, and these animals were not able to demonstrate the effect of IL-15 on macrophages well. Our future research will employ macrophage-specific *IL-15* KO mice using *Lyz2-Cre* to directly verify this hypothesis, which will be regarded as the core direction of our subsequent studies. Furthermore, we only reported IL-15 interactions between macrophages and cardiomyocytes after AMI. Future studies must assess the long-term effects of IL-15 inhibition on repair, metabolism, and immune homeostasis. Last but not least, follow-up research examining the molecular pathways linking IL-15 signaling, glycolysis, and efferocytosis could provide deeper insights into how metabolic reprogramming facilitates macrophage functions in inflammatory diseases.

5 Conclusions

Our study identified IL-15 as a pivotal mediator in the early-stage pathogenesis of AMI, driving inflammatory responses, disrupting macrophage efferocytosis,

and promoting glycolytic reprogramming in macrophages. These mechanisms synergistically contribute to infarct expansion, heighten myocardial apoptosis, and facilitate progressive cardiac dysfunction. Collectively, our findings underscore IL-15 as a promising therapeutic target that has the potential to attenuate inflammation, limit cardiac injury, and improve clinical outcomes in AMI patients.

Data availability statement

The data supporting this research can be requested from the corresponding author upon reasonable request.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 82270498 and 81570322). We thank Prof. Wei ZHU (Department of Cardiology, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China) for his instructional advice and technical assistance.

Author contributions

Lei GUO and Zhehui YIN conducted the experiments, analyzed the data, and wrote the manuscript draft. Ning ZHANG and Han CHEN provided conceptual advice. Zhuo WANG, Yuxue HUANG, Jiniu HUANG, Yayou YOU, Chenyun ZHANG, Qinyi BAO, and Shuxin LEI participated in specific experiments and analyzed the results. Jun JIANG and Xiaojie XIE designed the study and revised the manuscript. All authors have read and approved the final manuscript, and therefore, have full access to all the data in the study and take responsibility for the integrity and security of the data.

Compliance with ethics guidelines

Lei GUO, Zhehui YIN, Ning ZHANG, Han CHEN, Zhuo WANG, Yuxue HUANG, Jiniu HUANG, Yayu YOU, Chenyun ZHANG, Qinyi BAO, Shuxin LEI, Jun JIANG, and Xiaojie XIE declare that they have no conflicts of interest.

This investigation was performed according to the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards, and written informed consent was obtained from all study participants prior to inclusion in the study with approval by the Ethics Committee of The Second Affiliated Hospital of Zhejiang University School of Medicine (No. 2021-0624). All animal procedures conformed to the guidelines from Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes. The animal's study protocol was approved by the Ethics Committee of The Second Affiliated Hospital of Zhejiang University School of Medicine (No. AIRB-2021-1012). All institutional and national guidelines for the care and use of laboratory animals were followed.

Declaration on the use of generative AI tools

During the preparation of this work, the authors used Grammarly to improve language and readability. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

- Allard-Chamard H, Mishra HK, Nandi M, et al., 2020. Interleukin-15 in autoimmunity. *Cytokine*, 136:155258. <https://doi.org/10.1016/j.cyto.2020.155258>
- Ameri K, Bayardorj D, Samurkashian R, et al., 2020. Administration of interleukin-15 peptide improves cardiac function in a mouse model of myocardial infarction. *J Cardiovasc Pharmacol*, 75(1):98-102. <https://doi.org/10.1097/FJC.0000000000000764>
- Cai DQ, Xaymardan M, Holm JM, et al., 2003. Age-associated impairment in TNF- α cardioprotection from myocardial infarction. *Am J Physiol Heart Circ Physiol*, 285(2):H463-H469. <https://doi.org/10.1152/ajpheart.00144.2003>
- Colpitts SL, Puddington L, Lefrançois L, 2015. IL-15 receptor α signaling constrains the development of IL-17-producing $\gamma\delta$ T cells. *Proc Natl Acad Sci USA*, 112(31):9692-9697. <https://doi.org/10.1073/pnas.1420741112>
- Daseke MJ II, Tenkorang-Impraim MAA, Ma YG, et al., 2020. Exogenous IL-4 shuts off pro-inflammation in neutrophils while stimulating anti-inflammation in macrophages to induce neutrophil phagocytosis following myocardial infarction. *J Mol Cell Cardiol*, 145:112-121. <https://doi.org/10.1016/j.yjmcc.2020.06.006>
- Dozio E, Malavazos AE, Vianello E, et al., 2014. Interleukin-15 and soluble interleukin-15 receptor α in coronary artery disease patients: association with epicardial fat and indices of adipose tissue distribution. *PLoS ONE*, 9(3):e90960. <https://doi.org/10.1371/journal.pone.0090960>
- Ghali R, Habeichi NJ, Kaplan A, et al., 2020. IL-33 induces type-2-cytokine phenotype but exacerbates cardiac remodeling post-myocardial infarction with eosinophil recruitment, worsened systolic dysfunction, and ventricular wall rupture. *Clin Sci*, 134(11):1191-1218. <https://doi.org/10.1042/CS20200402>
- Gokkusu C, Aydin M, Ozkok E, et al., 2010. Influences of genetic variants in interleukin-15 gene and serum interleukin-15 levels on coronary heart disease. *Cytokine*, 49(1):58-63. <https://doi.org/10.1016/j.cyto.2009.09.004>
- Gómez-Nicola D, Valle-Argos B, Suardíaz M, et al., 2008. Role of IL-15 in spinal cord and sciatic nerve after chronic constriction injury: regulation of macrophage and T-cell infiltration. *J Neurochem*, 107(6):1741-1752. <https://doi.org/10.1111/j.1471-4159.2008.05746.x>
- Hamada M, Nakamura M, Tran MTN, et al., 2014. MafB promotes atherosclerosis by inhibiting foam-cell apoptosis. *Nat Commun*, 5:3147. <https://doi.org/10.1038/ncomms4147>
- Hao ZY, Ma J, Dai JJ, et al., 2020. A real-world analysis of cardiac rupture on incidence, risk factors and in-hospital outcomes in 4190 ST-elevation myocardial infarction patients from 2004 to 2015. *Coron Artery Dis*, 31(5):424-429. <https://doi.org/10.1097/MCA.0000000000000877>

- Irwin MW, Mak S, Mann DL, et al., 1999. Tissue expression and immunolocalization of tumor necrosis factor- α in postinfarction dysfunctional myocardium. *Circulation*, 99(11):1492-1498.
<https://doi.org/10.1161/01.cir.99.11.1492>
- Jakubzick CV, Randolph GJ, Henson PM, 2017. Monocyte differentiation and antigen-presenting functions. *Nat Rev Immunol*, 17(6):349-362.
<https://doi.org/10.1038/nri.2017.28>
- Jung M, Ma YG, Iyer RP, et al., 2017. IL-10 improves cardiac remodeling after myocardial infarction by stimulating M2 macrophage polarization and fibroblast activation. *Basic Res Cardiol*, 112(3):33.
<https://doi.org/10.1007/s00395-017-0622-5>
- Lambert JM, Lopez EF, Lindsey ML, 2008. Macrophage roles following myocardial infarction. *Int J Cardiol*, 130(2):147-158.
<https://doi.org/10.1016/j.ijcard.2008.04.059>
- Li YF, Feng LF, Li GR, et al., 2020. Resveratrol prevents ISO-induced myocardial remodeling associated with regulating polarization of macrophages through VEGF-B/AMPK/NF- κ B pathway. *Int Immunopharmacol*, 84:106508.
<https://doi.org/10.1016/j.intimp.2020.106508>
- Li YQ, Sun XQ, Liu XN, et al., 2022. P2X7R-NEK7-NLRP3 inflammasome activation: a novel therapeutic pathway of Qishen granule in the treatment of acute myocardial ischemia. *J Inflamm Res*, 15:5309-5326.
<https://doi.org/10.2147/JIR.S373962>
- Luan Y, Yang Y, Luan Y, et al., 2024. Targeting ferroptosis and ferritinophagy: new targets for cardiovascular diseases. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)*, 25(1):1-22.
<https://doi.org/10.1631/jzus.B2300097>
- McDonald PP, Russo MP, Ferrini S, et al., 1998. Interleukin-15 (IL-15) induces NF- κ B activation and IL-8 production in human neutrophils. *Blood*, 92(12):4828-4835.
<https://doi.org/10.1182/blood.V92.12.4828>
- Mortier E, Bernard J, Plet A, et al., 2004. Natural, proteolytic release of a soluble form of human IL-15 receptor α -chain that behaves as a specific, high affinity IL-15 antagonist. *J Immunol*, 173(3):1681-1688.
<https://doi.org/10.4049/jimmunol.173.3.1681>
- Mortier E, Woo T, Advincula R, et al., 2008. IL-15R α chaperones IL-15 to stable dendritic cell membrane complexes that activate NK cells via trans presentation. *J Exp Med*, 205(5):1213-1225.
<https://doi.org/10.1084/jem.20071913>
- Ngai D, Schilperoort M, Tabas I, 2023. Efferocytosis-induced lactate enables the proliferation of pro-resolving macrophages to mediate tissue repair. *Nat Metab*, 5(12):2206-2219.
<https://doi.org/10.1038/s42255-023-00921-9>
- Patidar M, Yadav N, Dalai SK, 2016. Interleukin 15: a key cytokine for immunotherapy. *Cytokine Growth Factor Rev*, 31:49-59.
<https://doi.org/10.1016/j.cytogfr.2016.06.001>
- Perera PY, Lichy JH, Waldmann TA, et al., 2012. The role of interleukin-15 in inflammation and immune responses to infection: implications for its therapeutic use. *Microbes Infect*, 14(3):247-261.
<https://doi.org/10.1016/j.micinf.2011.10.006>
- Ridker PM, Everett BM, Thuren T, et al., 2017. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med*, 377(12):1119-1131.
<https://doi.org/10.1056/NEJMoa1707914>
- Roth GA, Mensah GA, Johnson CO, et al., 2020. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. *J Am Coll Cardiol*, 76(25):2982-3021.
<https://doi.org/10.1016/j.jacc.2020.11.010>
- Stone KP, Kastin AJ, Pan WH, 2011. NF- κ B is an unexpected major mediator of interleukin-15 signaling in cerebral endothelium. *Cell Physiol Biochem*, 28(1):115-124.
<https://doi.org/10.1159/000331720>
- van den Bossche J, Baardman J, Otto NA, et al., 2016. Mitochondrial dysfunction prevents repolarization of inflammatory macrophages. *Cell Rep*, 17(3):684-696.
<https://doi.org/10.1016/j.celrep.2016.09.008>
- Wang YC, Zheng Y, Qi BC, et al., 2023. α -Lipoic acid alleviates myocardial injury and induces M2b macrophage polarization after myocardial infarction via HMGB1/NF- κ B signaling pathway. *Int Immunopharmacol*, 121:110435.
<https://doi.org/10.1016/j.intimp.2023.110435>
- Wei YZ, Lan Y, Zhong YC, et al., 2020. Interleukin-38 alleviates cardiac remodelling after myocardial infarction. *J Cell Mol Med*, 24(1):371-384.
<https://doi.org/10.1111/jcmm.14741>
- Westman PC, Lipinski MJ, Luger D, et al., 2016. Inflammation as a driver of adverse left ventricular remodeling after acute myocardial infarction. *J Am Coll Cardiol*, 67(17):2050-2060.
<https://doi.org/10.1016/j.jacc.2016.01.073>
- Wuttge DM, Eriksson P, Sirsjö A, et al., 2001. Expression of interleukin-15 in mouse and human atherosclerotic lesions. *Am J Pathol*, 159(2):417-423.
[https://doi.org/10.1016/S0002-9440\(10\)61712-9](https://doi.org/10.1016/S0002-9440(10)61712-9)
- Xuan WL, Huang WX, Wang RJ, et al., 2017. Elevated circulating IL-32 presents a poor prognostic outcome in patients with heart failure after myocardial infarction. *Int J Cardiol*, 243:367-373.
<https://doi.org/10.1016/j.ijcard.2017.03.065>
- Yan XX, Anzai A, Katsumata Y, et al., 2013. Temporal dynamics of cardiac immune cell accumulation following acute myocardial infarction. *J Mol Cell Cardiol*, 62:24-35.
<https://doi.org/10.1016/j.yjmcc.2013.04.023>
- Zhang YQ, Zhao JL, Lau WB, et al., 2013. Tumor necrosis factor- α and lymphotoxin- α mediate myocardial ischemic injury via TNF receptor 1, but are cardioprotective when activating TNF receptor 2. *PLoS ONE*, 8(5):e60227.
<https://doi.org/10.1371/journal.pone.0060227>
- Zhou SF, Yuan J, Liao MY, et al., 2014. IL-17A promotes ventricular remodeling after myocardial infarction. *J Mol Med*, 92(10):1105-1116.
<https://doi.org/10.1007/s00109-014-1176-8>
- Zou H, Zhang MY, Yang X, et al., 2024. Cynaroside regulates the AMPK/SIRT3/Nrf2 pathway to inhibit doxorubicin-induced cardiomyocyte pyroptosis. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)*, 25(9):756-772.
<https://doi.org/10.1631/jzus.B2300691>

Supplementary information

Materials and methods; Tables S1–S4; Figs. S1–S5