

Rietveld quantification of γ -C₂S conversion rate supported by synchrotron X-ray diffraction images

基于同步回旋加速器X射线衍射图像的 γ -C₂S转化率的Rietveld量化法

Citation: Pi-qi Zhao, Xian-ping Liu, Jian-guo Wu, Pei-ming Wang, 2013. Rietveld quantification of γ -C₂S conversion rate supported by synchrotron X-ray diffraction image. *Journal of Zhejiang University- SCIENCE A (Applied Physics & Engineering)*, 14(11):815-821. [doi:10.1631/jzus.A1300215]

Main goal

The Rietveld method and synchrotron X-ray diffraction images were used to accurately quantify γ -C₂S conversion rate.

Results and discussion




Fig. 2 LXRD pattern of β -C₂S air-cooled at the first time




Fig. 3 LXRD pattern of β -C₂S water-cooled at the second time




Fig. 5 Synchrotron X-ray diffraction images of C_2S sample with (a) and without (b) polyimide tape (Fig. 5b is obtained through subtracting the transmission X-ray diffraction images of polyimide tape from Fig. 5a)

Integrated diffraction pattern




Fig. 7 Rietveld QXRD pattern of β - C_2S air-cooled at the first time

Fig. 8 Rietveld QXRD pattern of β - C_2S water-cooled at the second time





Fig. 10 Integrated diffraction pattern of C_2S and $\alpha-Al_2O_3$ mixture converted from synchrotron X-ray diffraction images



\ominus : $\beta-C_2S$; Δ : $\gamma-C_2S$; Π : $\alpha-Al_2O_3$

Fig. 11 Rietveld QXRD pattern for C_2S and $\alpha-Al_2O_3$ mixture

Conclusions

The relative error between the weighted fraction and the Rietveld weight fraction of the phases (sample of C_2S and $\alpha-Al_2O_3$) is 2.6%. The absolute error of the $\gamma-C_2S$ conversion rate between the two LXRD Rietveld refinements is 3.6%. The Rietveld QXDA analysis is an appropriate and accurate method to quantify the $\gamma-C_2S$ phase conversion.