

Cite this as: Shuang LIANG, Xue-lai WANG, Ming-yang ZOU, Han WANG, Xue ZHOU, Cai-hong SUN, Wei XIA, Li-jie WU, Takashi X. FUJISAWA, Akemi TOMODA, 2014. Family-based association study of *ZNF533*, *DOCK4* and *IMMP2L* gene polymorphisms linked to autism in a northeastern Chinese Han population. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, 15(3):264-271. [doi:10.1631/jzus.B1300133]

Family-based association study of *ZNF533*, *DOCK4* and *IMMP2L* gene polymorphisms linked to autism in a northeastern Chinese Han population

ZNF533、*DOCK4*和*IMMP2L*基因多态性与 中国东北汉族孤独症的关联研究

Key words: Autism, *ZNF533*, *DOCK4*, *IMMP2L*, Northeastern Chinese Han population, Single-nucleotide polymorphism

关键词: 孤独症; *ZNF533*; *DOCK4*; *IMMP2L*; 中国东北汉族; 单核苷酸多态性

- *The prevalence of autism has increased to one per 50 children aged under 8 years
- *A study in a Caucasian population has identified two single-nucleotide polymorphisms (SNPs) in *ZNF533*, one in *DOCK4*, and two in *IMMP2L*, which were all significantly associated with autism.
- *They are located in *AUTS1* and *AUTS5*, which have been identified as autism susceptibility loci in several genome-wide screens.

- *Autism is clinically and genetically heterogeneous.
- *Meaningful study needs to replicate it in other population especially in Chinese Han
- *The present study aimed to investigate whether the *ZNF533*, *DOCK4*, and *IMMP2L* genes are also associated with autism in a northeastern Chinese Han population.
- *Significant associations were found between autism and the SNPs of *ZNF533* and *DOCK4*