

Cite this as: Jian Shen, Yu Zhu, Hai Yu, Zuo-xu Fan, Feng Xiao, Pan Wu, Qi-hui Zhang, Xiao-xing Xiong, Jian-wei Pan, Ren-ya Zhan, 2014. Buyang Huanwu decoction increases angiotensin-1 expression and promotes angiogenesis and functional outcome after focal cerebral ischemia. *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)*, **15**(3):272-280. [doi:10.1631/jzus.B1300166]

Buyang Huanwu Decoction increases angiotensin-1 expression and promotes angiogenesis and functional outcome after focal cerebral ischemia

Key words: Buyang Huanwu decoction, Angiotensin-1, Angiogenesis, Neurorestoration, Chronic brain injury

- BYHWD significantly improves neurological deficit, reduces the infarct and atrophy volume in rats, indicating its protective effect against cerebral ischemia.
- BYHWD significantly increased the microvessels density in the ischemic area, indicating its angiogenic effect against cerebral ischemia.
- BYHWD modulates angiogenesis in the boundary ischemic region and increases the expressions of Ang-1, which may be the mechanism by which BYHWD ameliorates chronic brain injury after focal cerebral ischemia in rats
- .