

Cite this as: Chang-ying LIU, Rui-hua LÜ, Jun LI, Ai-chun ZHAO, Xi-ling WANG, Umuhoza DIANE, Xiao-hong WANG, Chuan-hong WANG, Ya-sheng YU, Shu-mei HAN, Cheng LU, Mao-de YU, 2014. Characterization and expression profiles of MaACS and MaACO genes from mulberry (*Morus alba* L.). *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)*, 15(7):611-623. [doi:10.1631/jzus.B1300320]



Characterization and expression profiles of *MaACS* and *MaACO* genes from mulberry (*Morus alba* L.)

Key words: Mulberry fruit, MaACS, MaACO, Gene expression, ABA, Ethephon

Introduction

- Ethylene is involved in many aspects of the plant life cycle and one of the most important factors regulating softening, senescence and abscission of fruit, and the concentration of ethylene increases gradually during the development of climacteric fruit.
- ACS and ACO are the key enzymes producing ethylene, and are typically encoded by multiple *ACS* genes and *ACO* genes. ACS and ACO act as accelerators in fruit ripening by catalyzing the production of ethylene at the late fruit ripening stage, leading to irreversible senescence. It has been shown that once the expression levels of *ACS* and *ACO* genes were suppressed or mutated, the production of ethylene was down-regulated and the shelf life of fruit was also prolonged.
- Mulberry fruit is one of the most popular fruits worldwide with short maturity stages and shelf-life. There are no reports on *ACS* and *ACO* genes in mulberry, and never know which how to regulate the development and senescence of mulberry fruit.




Result



MaACS1	413	VCIANMDDHTMEVALKRI ⁺ RSFV-LQKKEPVVVP--NKKKCWQDH-NLRLS ⁺ SARRFD---
MaACS2	412	VCFANME ⁺ EETVNVALERI ⁺ RSFVGKEAKQAAVPPVAKAKRNWRNNGNLRLS ⁺ FNSRRHEQEG
MaACS3	408	VCFANMSEETLELAMQRLK ⁺ SFVTEYNRI ⁺ NNSNKESSAKNSRRRSLTKWVFRIS ⁺ FDGADRQ
MaACS4	406	MCFANMSEETLHVALQRLKAF ⁺ AETLIK ⁺ SKQ-QQQFGLS ⁺ FSRTRSLTRWVFQLASDHDHRE
MaACS5	415	VCFANMSEK ⁺ TLEVALKRI ⁺ HNFT ⁺ EKRETISSRN-----

MaACS1	466	DFL----MASPH ⁺ SPIPH ⁺ SPLVQART
MaACS2	472	GFMSAAHMLS ⁺ PHSPMPQSPLLRART
MaACS3	468	PEER-----
MaACS4	465	TDR-----
MaACS5	446	-----

Result



Conclusion

- We are the first to report *ACS* and *ACO* genes from mulberry and demonstrate their expression profiles.
- Tissue-specific-expression analysis revealed that *MaACS* and *MaACO* genes showed different patterns in various mulberry tissues, and two patterns throughout the development of mulberry fruit. *MaACS* genes also tend to be expressed in fruit rather than in other tissues.
- Further, the transcription levels of *MaACS* and *MaACO* genes could be up-regulated by ABA and ethephon