

Cite this as: Xi-ming LI, Ming-tao SHAO, Jian-juan WANG, Yue-lan WANG, 2014. Relationship between post-operative cognitive dysfunction and regional cerebral oxygen saturation and β -amyloid protein. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, 15(10):870-878. [doi:10.1631/jzus.B1400130]

Relationship between post-operative cognitive dysfunction and regional cerebral oxygen saturation and β -amyloid protein

Key words:

Laparoscopic pancreaticoduodenectomy, Post-operative cognitive dysfunction, Regional cerebral oxygen saturation, β -amyloid protein

Summary

This article mainly focused on the correlation of post-operative cognitive dysfunction (POCD) and rSO_2 and $A\beta$ in patients undergoing laparoscopic pancreaticoduodenectomy.

- The development of POCD is associated with rSO_2 and $A\beta$.
- Monitoring of rSO_2 might be useful in the prediction of POCD.
- $A\beta$ might be used as a sensitive biochemical marker to predict the occurrence of POCD.



Innovation points

- ◆ The $rSO_{2\%max}$ value of over 10.2% might be a potential predictor of neurocognitive injury for those patients.
- ◆ plasma $A\beta$ might be a potential biomarker for POCD.



Innovation points

◆ Multivariate logistic regression analysis has shown that basal body temperature, PaCO₂, rSO_{2%max} and Aβ seem to play more important role for the interpretation of POCD.

◆ Routine blood gas analysis, monitoring of rSO₂ and detection of plasma Aβ might be useful tools for predicting the occurrence of POCD.

