

Cite this as: Zhao-ji LIU, Gregg L. SEMENZA, Hua-feng ZHANG, 2015. Hypoxia-inducible factor 1 and breast cancer metastasis. *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)*, **16**(1):32-43. [doi:10.1631/jzus.B1400221]

Hypoxia-inducible factor 1 and breast cancer metastasis


Key words: breast cancer, hypoxia-inducible factor 1(HIF-1),
metastasis

Research Summary

This review mainly focused on the roles of HIF-1 in regulating breast cancer cell metastasis, specifically its effects on multiple key steps of metastasis:

METASTATIC BREAST CANCER

The cancer has started to spread to other parts of the body.



- EMT
- Invasion
- Extravasation
- Metastatic niche formation



HIF-1-regulated non-coding RNAs in breast cancer metastasis was discussed as well.

Innovation points

- Introduction of Basic Biology of HIF-1 and the roles of hypoxia environment in breast cancer progression.

- Summary of the most updated research progress about HIF-1 in breast cancer metastasis and non-coding RNAs involved.

- Emphasis of the therapeutic opportunities for breast cancer through targeting the HIF-1 pathway





Figure 2

Innovation points

A series of comprehensive tables and figures were generated to summarize the latest knowledge about Hypoxia-inducible factor-1 mediates breast cancer metastasis.

Table 1 | Examples of the hypoxia/HIF-1 regulated ncRNAs in cancer metastasis.

Table 2 | Inhibitors of HIF-1 pathway.

Figure 1. HIF-1 is regulated by oxygen tension.

Figure 2. Cancer metastasis regulated by HIF-1.

