




Mutation breeding of high 4-androstene-3,17-dione-producing *Mycobacterium neoaurum* ZADF-4 by atmospheric and room temperature plasma treatment

Key words: *Mycobacterium neoaurum*, Atmospheric and room temperature plasma (ARTP), Mutation breeding, 4-Androstene-3,17-dione (AD), 1,4-Androstadiene-3,17-dione (ADD)

Cite this as: Chao LIU, Xian ZHANG, Zhi-ming RAO, Ming-long SHAO, Le-le ZHANG, Dan WU, Zheng-hong XU, Hui LI, 2015. Mutation breeding of high 4-androstene-3,17-dione-producing *Mycobacterium neoaurum* ZADF-4 by atmospheric and room temperature plasma treatment . *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, **16**(4):286-295. [doi:10.1631/jzus.B1400274]

This paper mainly focused on increasing the production ratio of AD in the mixture of AD/ADD in *Mycobacterium neoaurum* mutant through the deficiency of key enzyme, 3-ketosteroid- Δ 1-dehydrogenases(KSDD)



- .Mutagenesis with the ARTP biological breeding system
- .Mutant selection
- . Enzyme activity assay of KSDD and product analysis
- . Sequencing and comparison of the mutant *ksdd* gene with original strain



Innovation points



This work provides a promising candidate of *M. neoaurum* for industrial AD production as well as an extension of the H⁺ dependent dehydrogenase deficient screening method.