


Cite this as: Ya-qian LI, Kai SONG, Ya-chai LI, Jie CHEN, 2016. Statistical culture-based strategies to enhance chlamydospore production by *Trichoderma harzianum* SH2303 in liquid fermentation. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*. **17**(8):619-627.
<http://dx.doi.org/10.1631/jzus.B1500226>

Statistical culture-based strategies to enhance chlamydospore production by *Trichoderma harzianum* SH2303 in liquid fermentation


Key words: *Trichoderma harzianum* SH2303, Chlamydospore; Plackett-Burman screening, Box-Behnken design, Liquid fermentation optimization

Research Summary

This study mainly focused on the optimization of a cost effective medium and cultural conditions for the production of chlamydospores in *T. harzianum* SH2303, and thus to lay a solid foundation for the development of *Trichoderma*-based active chlamydospore formulations in agriculture.



***T. harzianum* SH2303**





chlamydospore



Trichoderma formulations

Innovation points


- **Introduction** of the three factors cornmeal, pH, and glycerol play pivotal roles for enhancing the production of chlamydospores in strain SH2303.
- **Summary** of the optimal chlamydospores (4.5×10^8 spores/mL) obtained under the following condition: corn flour 62.86 g/L, glycerol 7.54 ml/L, pH 4.17 and 6 days incubation in liquid fermentation.
- **Emphasis** of the possibility for development of novel biocontrol agents, of which chlamydospore as the main composition of liquid fermentation in a cost-efficient mode .



Open strategy

Closed strategy

**Medium and cultural conditions
optimization design process**



A integrated fermentation optimization design for the maximize production of chlamydopores in *T. harzianum*.