


**Cite this as:** Shi-cheng YU, Hai-yang WU, Wei WANG, Li-wei XU, Guo-qing DING, Zhi-gen ZHANG, Gong-hui LI, 2016. High-pressure balloon dilation for male anterior urethral stricture: single-center experience. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*. 17(9):722-727. <http://dx.doi.org/10.1631/jzus.B1600096>


# High-pressure balloon dilation for male anterior urethral stricture: single-center experience

**Key words:** Urethral stricture, High-pressure balloon dilation, Urethrotomy

From January 2009 to December 2012, a total of 31 patients with anterior urethral strictures underwent high-pressure balloon dilation (HPBD), while another 25 cases were treated by direct vision internal urethrotomy (DVIU).





A



B

**(a)** The appearance of urethral stricture at the beginning of the HPBD procedure. **(b)** The high pressure balloon catheter was inserted through the stricture segment over the guide wire.

**A****B**

Kaplan-Meier curves predicting stricture-free rate after HPBD and DVIU: **(A)** Survival analysis demonstrated no significant difference at 36 months ( $p=0.21$ ). **(B)** Survival analysis showed significantly higher stricture-free rate in the HPBD group at 12 months ( $p=0.02$ ).

# Conclusion

HPBD might be an alternative for the treatment modality of anterior urethral stricture disease.