


Cite this as: Lan-lan WAN, Da-qi ZHANG, Jin-nan ZHANG, Li-qun REN, 2017. Anti-hepatocarcinoma activity of TT-1, an analog of melittin, combined with interferon- α via promoting the interaction of NKG2D and MICA. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, 18(6):522-531.
<http://dx.doi.org/10.1631/jzus.B1600369>

Anti-hepatocarcinoma activity of TT-1, an analog of melittin, combined with interferon- α via promoting the interaction of NKG2D and MICA


Key words: TT-1; IFN- α ; NK cells; Hepatocarcinoma; Immunotherapy

Research Summary

This article mainly focused on the anti-hepatocellular carcinoma effect of the combined therapy of TT-1 and IFN- α *in vitro* and *in vivo*, and the key effect and mechanisms were summarized as following:



- **TT-1 inhibited the proliferation of liver cancer cells**
- **TT-1 + IFN- α suppressed the tumor growth *in vivo* though stimulating NK cells**
- **TT-1 increased the expression of MICA and inhibited sMICA secretion**
- **TT-1 + IFN- α enhanced the cytotoxicity of NK92 cells**



Innovation points

A series of novelties about this research was shown as below:

- **This study was firstly to investigate the effect of TT-1 combined IFN- α**
- **TT-1 combined IFN- α suppressed the tumor growth *in vivo***
- **TT-1 not only increased the membrane MICA expression but also prevented membrane MICA from shedding as inhibitory soluble MICA**
- **The activation of NK cytotoxicity was the target in TT-1 + IFN- α - induced anti-tumor effect**