


***Cite this as:*** Meng-qi YAO, Qiu-xian ZHENG, Jia XU, Jing-wen DENG, Tian-tian GE, Hai-bo ZHOU, Feng-tian WU, Xin-yu GU, Qin YANG, Yan-li REN, Gang WANG, Zhi CHEN, 2020. Factors associated with a SARS-CoV-2 recurrence after hospital discharge among patients with COVID-19: systematic review and meta-analysis. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, **21**(12):940-947.  
<http://doi.org/10.1631/jzus.B2000304>

# **Factors associated with a SARS-CoV-2 recurrence after hospital discharge among patients with COVID-19: systematic review and meta-analysis**



**Key words:** Recurrence cases, SARS-CoV-2, Risk factors, Meta-analysis

# Research Summary

This meta-analysis summarized the demographical, clinical, and laboratory characteristics of non-recurrence and recurrence groups.



<https://curehht.org/covid19/>



# *Research summary*

- In this analysis the proportion of SARS-CoV-2 recurrence after hospital discharge patients with COVID-19 is 17.65%.
- The older patients are more likely to have SARS-CoV-2 recurrence.
- Patients combined with clinical symptoms cough, sputum, and dizziness are at higher risk of SARS-CoV-2 recurrence.
- Patients with bilateral lung infiltrates and decreased white blood cell, platelet, and CD4 + T counts are more likely have SARS-CoV-2 recurrence.

# ***Innovation points***

- These factors can be regarded as early warning indicators of SARS-CoV-2 recurrence
- These factors may help the clinical development of individualized management strategies.