


Cite this as: Wei-ling XIANG, Jing-jing CHENG, Lian-peng WU, Bing-yu CHEN, Wen-xin LI, Dan-ying QIU, Wei ZHANG, Fei-hang GE, Dong CHEN, Zhen WANG, 2020. Clinical characteristics and plasma antibody titer of patients with COVID-19 in Zhejiang, China. *Journal of Zhejiang University-Science B (Biomedicine & Biotechnology)*, **21**(12):955-960. <http://doi.org/10.1631/jzus.B2000593>

Clinical characteristics and plasma antibody titer of patients with COVID-19 in Zhejiang, China

Key words: Coronavirus disease 2019 (COVID-19), Convalescent plasma, Clinical characteristics, Antibody titer

Research Summary

This study mainly focused on antibodies profile, laboratory indicators and clinical feature, whether there is correlation between appearance of IgG/IgM and clinical characteristics.




Innovation points

- **Our study use COVID-19 IgG/IgM Rapid Test Cassette to detect the plasma antibody titers of convalescent plasma , search for antibody profile.**

- **Analysis and summarized correlation between clinical characteristics of severe and critical patients and laboratory indicators.**



Innovation points



- 4-6 weeks after the onset of symptoms, the level of plasma IgG titers in patients with COVID-19 during the recovery period can reach 1:320-1:640 or higher.

- Coagulation dysfunction and inflammation in the early stage of COVID-19 may influence the level of IgG and IgM antibody titers in the convalescent plasma.

