Cite this as: Yiying YANG, Qingqing SUN, Yang LIU, Hanzhi YIN, Wenping YANG, Yang WANG, Ying LIU, Yuxian LI, Shen PANG, Wenxi LIU, Qian ZHANG, Fang YUAN, Shiwen QIU, Jiong LI, Xuefeng WANG, Keqiang FAN, Weishan WANG, Zilong LI, Shouliang YIN. Development of a *pyrF*-based counterselectable system for targeted gene deletion in *Streptomyces rimosus*[J]. *Journal of Zhejiang University Science B*, 2021, 22(5): 383-396. https://doi.org/10.1631/jzus.B2000606

Development of a *pyrF*-based counterselectable system for targeted gene deletion in *Streptomyces rimosus*

JZUSB

Key words: **Counterselectable system, *pyrF*, 5-fluoroorotic acid, Gene deletion, *Streptomyces rimosus***

Research Summary

This paper mainly focused on the core components of development of a *pyrF*-based counterselectable system for targeted gene deletion in the following aspects:

- Identification of orotidine-5`-phosphate decarboxylase gene *pyrF in Streptomyces.*
- Constitution and principle of the *pyrF*-based screening system
- Targeted gene deletion in *S. rimosus* with *pyrF*-based gene knockout systems

JZUS-B

Innovation points

● Alignment analysis revealed that only one copy of the putative orotidine 5’-phosphate decarboxylase gene *pyrF* exists on the genome of *Streptomyces*.


●Deletion of *pyrF* conferred 5-fluoroorotic acid (5-FOA) resistance in the mutant lacking *pyrF* (Δ*pyrF*).

● This *pyrF*-based counterselectable system was very effective and enabled us to obtain the target gene knockout mutants by analyzing a small number of clones, and the time to obtain the desired mutants was greatly shortened.

● The advantage of this system is that the *pyrF* marker does not remain in the genome of Δ*pyrF* , so additional deletion and/or insertion using the same vector system is possible.

IZUSB

Innovation points



Schematic diagram depicting the steps for *otcR* deletion using the *pyrF*-based counterselectable system