
Cite this as: Xiaoyan TANG, Dezheng CHEN, Ling ZHANG, Ping FU, Yanxia CHEN, Zhou XIAO, Xiangcheng XIAO, Weisheng PENG, Li CHENG, Yanmin ZHANG, Hongbo LI, Kehui LI, Bizhen GOU, Xin WU, Qian YU, Lijun JIAN, Zaizhi ZHU, Yu WEN, Cheng LIU, Hen XUE, Hongyu ZHANG, Xin HE, Bin YAN, Liping ZHONG, Bin HUANG, Mingying MAO. Application of regional citrate anticoagulation in patients at high risk of bleeding during intermittent hemodialysis: a prospective multicenter randomized controlled trial[J]. Journal of Zhejiang University Science B, 2022, 23(11): 931-942. https://doi.org/10.1631/jzus.B2200082

Application of regional citrate anticoagulation in patients at high risk of bleeding during intermittent hemodialysis: a prospective multicenter randomized controlled trial

Key words: Citrate anticoagulation; intermittent hemodialysis; calcium-containing dialysate; saline flushing; anticoagulation


Research Summary

The purpose of this trial is to evaluate the effectiveness and safety of two-stage regional citrate anticoagulation (RCA) combined with sequential anticoagulation and standard calcium-containing dialysate in intermittent hemodialysis (IHD) treatment



Research methods

two-stage regional citrate anticoagulation with ordinary calcium containing dialysate





Dialysate flow 300ml/min
Ca2+ 1.25 ~ 1.5 mmol/L
HCO3- 22 ~ 26 mmol/L
Na+ 138 mmol/L
K+ 2.0 ~ 2.5 mmol/L
Mg2+ 0.5 mmol/L

Research results

RCA vs 0.9%NS



