Cite this as: Yunlong CHENG, Shu LIAN, Shuhui LI, Yusheng LU, Jie WANG, Xiaoxiao DENG, Shengyi ZHAI, Lee JIA. Combination of Se-methylselenocysteine, D-α-tocopheryl succinate, β-carotene, and L-lysine can prevent cancer metastases using as an adjuvant therapy[J]. Journal of Zhejiang University Science B, 2022, 23(11): 943-956. https://doi.org/10.1631/jzus.B2200232

Combination of Se-methylselenocysteine, D-α-tocopheryl succinate, β-carotene, and L-lysine can prevent cancer metastases using as an adjuvant therapy


Key words: Cancer prevention; Cancer metastases; Se; Combination drugs; Nutrition

Lee Jia
E-mail: cmapcjia1234@163.com or jiali@fzu.edu.cn.
ORICD: 0000-0001-6839-5545.

Nutrition for cancer therapy

Dietary modifications to improve cancer- drug efficacy

- **Nutrients nourished**
- **Enhancement of drug-based therapy**
- **Anticancer immune response activation**
- **Cancer-specific toxicity**




Pictures adapted from: Naama Kanarek, Boryana Petrova, David M. Sabatini 2020. Dietary modifications for enhanced cancer therapy. *Nature*, 579(7800):507-517. [doi:10.1038/s41586-020-2124-0]

Research Hightlights

A functional health product consists of four compounds(as follows)


- **improve drug efficacy**
- **enhance immunity**
- **antioxidant capacity**
- **improve life quality and prolong survival time**

A




Alpha Tocopheryl Acid Succinat

B




Se-methylselenocysteine

C



Beta Carotene

D



L-Lysine

JZUS-B