

Cite this as: Yunlong CHENG, Shu LIAN, Shuhui LI, Yusheng LU, Jie WANG, Xiaoxiao DENG, Shengyi ZHAI, Lee JIA. Combination of Se-methylselenocysteine, D- α -tocopheryl succinate, β -carotene, and L-lysine can prevent cancer metastases using as an adjuvant therapy[J]. Journal of Zhejiang University Science B, 2022, 23(11): 943-956.
<https://doi.org/10.1631/jzus.B2200232>

Combination of Se-methylselenocysteine, D- α -tocopheryl succinate, β -carotene, and L-lysine can prevent cancer metastases using as an adjuvant therapy

Key words: Cancer prevention; Cancer metastases; Se;
Combination drugs; Nutrition

Lee Jia


E-mail: cmapcjia1234@163.com or jjiali@fzu.edu.cn.

ORCID: 0000-0001-6839-5545.

Nutrition for cancer therapy

Dietary modifications to improve cancer- drug efficacy

- Nutrients nourished
- Enhancement of drug-based therapy
- Anticancer immune response activation
- Cancer-specific toxicity



Pictures adapted from: Naama Kanarek, Boryana Petrova, David M. Sabatini 2020.
Dietary modifications for enhanced cancer therapy. *Nature*, 579(7800):507-517.
[doi:10.1038/s41586-020-2124-0]

Research Highlights

A functional health product consists of four compounds(as follows)

- improve drug efficacy
- enhance immunity
- antioxidant capacity
- improve life quality and prolong survival time

