Cite this as: Natalia V. DEMENTIEVA, Yuri S. SHCHERBAKOV, Olga I. STANISHEVSKAYA, Anatoly B. VAKHRAMEEV, Tatiana A. LARKINA, Artem P. DYSIN, Olga A. NIKOLAEVA, Anna E. RYABOVA, Anastasiia I. AZOVTSEVA, Olga V. MITROFANOVA, Grigoriy K. PEGLIVANYAN, Natalia R. REINBACH, Darren K. GRIFFIN, Michael N. ROMANOV. Large-scale genome-wide SNP analysis reveals the rugged (and ragged) landscape of global ancestry, phylogeny, and demographic history in chicken breeds[J]. Journal of Zhejiang University Science B, 2024, 25(4): 324-340. http://doi.org/10.1631/jzus.B2300443

Large-scale genome-wide SNP analysis reveals the rugged (and ragged) landscape of global ancestry, phylogeny, and demographic history in chicken breeds

Key words: Chicken genome diversity; SNP analysis; Gene pool; Global ancestry; Phylogeny; Demographic history

JZUS-B

Research Summary

This research aimed to perform a large-scale genome-wide analysis of the landscape of the complex molecular architecture, genetic variability and detailed structure among 49 populations.



- Genetic diversity and genetic variability in small chicken populations
- Effective population size
- ADMIXTURE analysis
- Overall 49-population phylogeny analysis

Innovation points

- **Introduction of the genetic diversity landscape assessment in poultry for:**
  - preserving genetic resources;
  - performing the genetic identification of breeds;
  - studying the history of their genetic divergence;
  - improving the efficiency of breeding in farm animal populations



- **Summary** of large-scale genome-wide analysis of the landscape of the complex molecular architecture, genetic variability and detailed structure among 49 chicken populations

- **Emphasis** a refined evolutionary model of chicken breed formation proposed, which included egg, meat, dual dual-purpose types, and ambiguous breeds.

JZUSB

Innovation points

A series of tables and figures summarizing the genetic architecture data of 49 bird populations was compiled

Table 1 | Characteristics of runs of homozygosity (ROHs) on average for the populations of the main chicken breeds studied.

Figure 1 | Genetic identification and clustering of populations by multivariate scaling (MDS) in all 49 breeds and subpopulations distributed among four clusters.

Figure 2 | Effective population size in all 49 chicken populations.

Figure 3 | The ADMIXTURE-assisted rugged (and ragged) landscape of global ancestry, with the optimal number of ancestral populations being K = 29, in the gene pool breeds/subpopulations.

Figure 4 | A phylogenetic tree based on SNP genotypes and visualizing relationships among 49 chicken breeds and (sub)populations

JZUS-B