Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology) in press
www.jzus.zju.edu.cn; www.springer.com/journal/11585
E-mail: jzus_b@zju.edu.cn

Review

https://doi.org/10.1631/jzus.B2300607

The role of lncRNA in the crosstalk between osteogenesis and angiogenesis in the bone microenvironment

Shihua ZHANG[1*], Jianmin GUO[2*], Yuting HE[1], Zhiang SU[1], Yao FENG[1], Lan ZHANG[3], Jun ZOU[4], Xiquang WENG[1]✉, Yu YUAN[1]✉

[1]*School of Exercise and Health, Guangzhou Sport University, Guangzhou 510500, China*
[2]*School of Life Sciences,South University of Science and Technology, Shenzhen 518055, China*
[3]*College of Sports and Health, Shandong Sport University, Jinan 250102, China*
[4]*School of Exercise and Health, Shanghai University of Sport, Shanghai, 200438, China*

Abstract: Bone is a highly calcified and vascularized tissue. The vascular system plays a vital role in supporting bone growth and repair, such as the provision of nutrients, growth factors, and metabolic waste transfer. Moreover, the additional functions of the bone vasculature, such as the secretion of various factors and the regulation of bone-related signaling pathways, are essential for maintaining bone health. In the bone microenvironment, bone tissue cells play a critical role in regulating angiogenesis, including osteoblasts, bone marrow mesenchymal stem cells (BMSCs), osteoclasts, and others. Osteogenesis and bone angiogenesis are closely linked. The decrease in osteogenesis and bone angiogenesis caused by aging leads to osteoporosis. Long noncoding RNAs (lncRNAs) are involved in various physiological processes, including osteogenesis and angiogenesis. Recent studies have shown that lncRNAs could mediate the crosstalk between angiogenesis and osteogenesis. However, the mechanism by which lncRNAs regulate angiogenesis–osteogenesis crosstalk remains unclear. In this review, we describe in detail the ways in which lncRNAs regulate the crosstalk between osteogenesis and angiogenesis to promote bone health, aiming to provide new directions for the study of the mechanism by which lncRNAs regulate bone metabolism.

Key words: lncRNAs; Osteogenesis; Bone angiogenesis; Osteoporosis; Bone microenvironment

Unedited

1 Introduction

Long noncoding RNAs (lncRNAs) are involved in a wide range of aspects of gene expression regulation, including epigenetic, transcriptional, and post-transcriptional regulation. In addition to this, they also participate in the biological regulation of cell growth, migration, differentiation, reprogramming, and stress responses (Herman et al., 2022; Nojima and Proudfoot, 2022). LncRNA abnormalities lead to deficiencies in protein-coding capacity. At the epigenetic level, lncRNAs regulate DNA methylation and histone modification (Allas et al., 2019). Recently, many researchers have shown interest in studying bone diseases at the genetic level and discovering new bone disease-associated lncRNAs using high-throughput sequencing or microarray studies. LncRNAs are able to regulate osteogenesis, adipogenesis, and osteoclast differentiation through epigenetic modifications and miRNA adsorption, which can affect bone metabolism (Ghafouri-Fard

✉ Yu YUAN, yuanyumail@126.com
✉ Xiquan WENG, wengxq@gzsport.edu.cn
* The two authors contributed equally to this work
Yu YUAN, https://orcid.org/0000-0003-1032-6330
Shihua ZHANG, https://orcid.org/0000-0001-5429-0443

Received Aug. 29, 2023; Revision accepted Jan. 16, 2024;
Crosschecked xxx. xx, 20xx; Published online xxx. xx, 20xx

© Zhejiang University Press 2024

et al., 2021; Wang et al., 2022b). The vasculature, another essential factor that directly influences osteogenesis in the bone microenvironment, plays a crucial role in bone development, remodeling, and the maintenance of bone homeostasis (Chim et al., 2013; Dirckx et al., 2013; He et al., 2013). The continuing process of bone resorption and formation is cyclic during the bone life cycle, and neo-angiogenesis is an integral part of this process. Additionally, angiogenesis occurs prior to osteogenesis, and new blood vessels ensure the transit of osteoblast precursors and osteoclasts to specific sites (Zhang et al., 2021; Zheng et al., 2022), suggesting that osteogenesis is a vascular-dependent process. It has been reported that angiogenesis and osteogenesis are closely related spatially and temporally, and have activation and interactions between signaling molecules and signaling pathways, known as the "crosstalk between angiogenesis and osteogenesis" (Schipani et al., 2009; Lafage-Proust et al., 2010; Maes, 2013; Saran et al., 2014; Grosso et al., 2017). The crosstalk between angiogenesis and osteogenesis is a necessary pathway for the co-growth of the vascular and bone phases in the bone microenvironment and a catalyst for both. Many scholars have devoted themselves to the study of mechanisms for the treatment of osteoporosis and other related diseases. Among them, the crosstalk between angiogenesis and osteogenesis is essential. For example, it has been shown that mechanical stimulation can couple osteogenesis and angiogenesis and promote the interactive effects of vascular and osteogenic factors (Ma et al., 2023). In addition, the exogenous supplementation of angiogenic factors can be achieved through hydrogel delivery systems to promote bone disease recovery (Tsai et al., 2023). In recent years, lncRNAs have been shown to play crucial roles as endogenous regulators in the crosstalk between angiogenesis and osteogenesis, including regulating various cell proliferation and differentiation processes, regulating the expression of osteogenic and angiogenic factors, and so on. However, the mechanism by which lncRNAs regulate the crosstalk between angiogenesis and osteogenesis is still unclear. Therefore, we summarize the latest studies on the involvement of lncRNAs in the regulation of angiogenesis and osteogenesis, as well as those on the pathways by which lncRNAs regulate the crosstalk between angiogenesis and osteogenesis, to provide support for the treatment of pathologic bone-related disorders, and to provide new ideas for the study of the mechanism of lncRNA-regulated bone metabolism.

Unedited

2 Overview of LncRNA

More than 90% of the DNA sequences in the human genome are involved in transcription, and only 2% of the human DNA sequences code for proteins. RNAs that do not encode any proteins are referred to as noncoding RNAs (Dangwal et al., 2017). LncRNAs, widely distributed in mammalian cells, are more than 200 nucleotides long and are known as lncRNAs (Wilusz et al., 2009). Most of the lncRNAs identified so far are produced by the transcription of RNA polymerase II (pol-II) or RNA polymerase III (pol-III) (Kung et al., 2013). Initially, lncRNAs were considered to lack protein-coding capacity and to not have biological functions. However, increasing studies have shown that lncRNAs, an important class of regulatory molecules in the human genome, play a crucial role in various biological functions, such as cell proliferation and differentiation, cellular autophagy, embryonic development, substance metabolism, and tumorigenesis (Duret et al., 2006; Wang et al., 2021a). Although lncRNAs have no protein-coding ability, they can regulate gene expression levels at various levels, including epigenetic regulation, transcriptional level regulation, and post-transcriptional level regulation. Thus, there are various mechanisms by which lncRNAs perform their biological functions.

Currently, there is no uniform standard for classifying lncRNAs, mainly based on the indicators of genomic localization, mediated phenotype, mechanism of action, and subcellular localization. LncRNAs can be classified into positive lncRNAs, antisense lncRNAs, intronic lncRNAs, bi-directional lncRNAs, and intergenic lncRNAs according to the positional relationship of their neighboring protein-coding genes. Meanwhile, depending on their functions, lncRNAs can also be categorized as backbone molecules, signaling molecules, decoy molecules, and primers (St Laurent et al., 2015; Mattick, 2018). LncRNAs are closely associated with their subcellular localization. A few lncRNAs are localized in the cytoplasm and regulate biological functions

by affecting the stability of mRNAs, participating in translational regulation and competing for binding microRNAs (miRNAs) (Han et al., 2014; Kim et al., 2015) to regulate cellular biological functions. The diverse subcellular localization of lncRNAs makes their functions even more complex.

Numerous studies have confirmed that lncRNAs participate in the regulation of gene expression in three main ways. Firstly, at the transcriptional level, lncRNAs can control the transcription of other genes by regulating the binding and assembly of transcription factors, forming triple-stranded complexes with regulatory sequences, or interfering with the transcription process through binding with pol-II. Secondly, lncRNAs are more complex in epigenetic regulation because lncRNAs can control chromatin remodeling by recruiting chromatin modifiers to DNA targets or by acting as carriers of histone modification complexes by inducing their DNA methylation (Zhu et al., 2019). Finally, the function of lncRNAs relies on interactions with RNA-binding proteins (RBPs), which are highly expressed in the brain in terms of regulation at the post-transcriptional level (Smart et al., 2007). LncRNAs bind to RBPs to form a large number of lncRNA/RBP complexes, which can recruit various protein complexes and activate downstream molecules. For example, lncRNA regulates the expression of osteogenic genes during post-transcriptional mRNA modification as a competing endogenous RNA (ceRNA) (Chen et al., 2021a).

It has been shown that the subcellular equilibrium between ceRNAs, miRNAs, and mRNA targets constitutes a complex network that can fine-tune the regulation of gene expression during adaptation, stress response, and development (Mattick, 2018). Here, we emphasize the significance of lncRNAs and miRNA interactions in this network. They regulate each other by competing for binding to the corresponding MREs, effectively controlling the post-transcriptional regulation of miRNAs (Karreth et al., 2011; Tay et al., 2011). In a cancer biology study, the lncRNA SNHG1 was found to regulate cancer cell metabolism by way of the sponge adsorption of multiple miRNAs (Yan et al., 2017; Li et al., 2018; Tian et al., 2018; Xu et al., 2018; Zhang et al., 2018b). MiRNAs are endogenous noncoding RNAs consisting of 21–25 nucleotides, which regulate the stability, translation, and degradation of mRNAs, leading to the modulation of physiological processes or signaling pathways. The function of miRNAs is related to tissue and organ formation, apoptosis, cell proliferation, differentiation, embryonic development, and various diseases. It can regulate the expression of more than 30% of the genes in the body (Bartel, 2004). MiRNAs have been demonstrated to regulate osteogenesis and angiogenesis in a variety of studies and are strongly correlated with the differentiation of BMSCs, osteoblasts, and osteoclasts (Bellavia et al., 2019). For example, miR-188 is significantly elevated in human mesenchymal stem/stromal cells (hMSCs) from senescent mice, leading to defective differentiation of hMSCs (Li et al., 2015), and miR-214 is significantly up-regulated in osteoblasts from older adults, inhibiting bone formation by suppressing the transcription factor 4(TCF4)(Wang et al., 2013). Notably, the specific expression of miRNAs in blood vessels is a crucial regulator of angiogenesis. LncRNAs have been shown to activate the relevant pathways by targeting miRNAs to promote angiogenesis and osteogenic differentiation (Ding et al., 2021), suggesting that lncRNAs can regulate osteogenic/angiogenic coupling by adsorbing miRNAs. It has also been proved that lncRNAs can improve the process of crosstalk between osteogenesis and angiogenesis by enhancing the expression of vascular endothelial growth factor (VEGF) during the osteogenic differentiation of BMSCs (Zhang et al., 2020). It is worth noting that the structure and function of lncRNAs define the way they exert their biological regulatory roles. Much research indicates that lncRNAs, together with miRNAs, have participated in the crosstalk between osteogenesis and angiogenesis.

Unedited

3 Angiogenesis-coupled osteogenesis in the bone microenvironment

There is a close relationship between bone blood flow and bone density; people with osteoporosis tend to have a lower bone blood supply than healthy individuals (Alagiakrishnan et al., 2003). Furthermore, animal studies have revealed that a reduction in bone mass in osteoporotic mice is accompanied by a corresponding decrease in the amount of blood supplied to the bones (Ding et al., 2011). Bone angiogenesis and osteogenesis

are mutually coupled processes; on the one hand, bone angiogenesis promotes osteogenesis, and a decline in bone angiogenesis capacity decreases osteogenesis ability (Xiao et al., 2018). On the other hand, in the bone microenvironment, osteoblasts can secrete angiogenic factors to regulate the proliferation and vascularization of vascular endothelial cells, thereby promoting bone angiogenesis (Maes et al., 2010).

3.1 Influence of angiogenesis on osteogenesis in the bone microenvironment

Bone vasculature plays an important regulatory role in the occurrence, development, and rehabilitation of various bone-related diseases. At the time of fracture, the blood vessels in the bone will rapidly gather and grow toward the injury site, delivering various growth factors and cells that promote osteogenesis and participate in the repair of the fracture (Baker et al., 2018; Bahney et al., 2019). Osteoporosis can be caused by various factors, including vascular anomalies that disrupt blood flow to the bone, reduce the number of blood vessels in the bone, and lead to abnormal blood vessel structure. Fortunately, several strategies can help alleviate osteoporosis symptoms, including improving blood flow to the bone (Peng et al., 2014; Jing et al., 2017). The type, spatial structure, and function of blood vessels in the bone microenvironment underlie the vascular effects on osteogenesis. Bone is a highly vascularized tissue, similar to other organs, but with some inconsistencies with the vasculature of other organs, such as growth plates and articular cartilage. Simultaneously, the skeletal system functions as an adaptable feedback regulatory mechanism that continuously integrates mechanical, biochemical, and neural signals (Harada and Rodan, 2003). The vascular system in bone exhibits a typical hierarchical structure, with branches of the arterial trunk flowing into an extensive network of capillaries before converging into large veins in the center of the backbone (Kusumbe et al., 2014). Although bone is the primary site for bone marrow storage, a sophisticated vascular system consisting of arteries, veins, and capillaries also exists. The capillaries are particularly specialized and have been shown to play a crucial role in maintaining bone health.

In 2014, Kusumbe et al. identified new microvessels in the mouse skeletal system with distinct morphological, molecular, and functional properties (Kusumbe, et al., 2014). They distinguished them based on the expression levels of platelet endothelial cell adhesion molecule-1 (CD31) (known as PECAM1) and endomucin (EMCN) as a distinguishing criterion, referring to the CD31hiEMCNhi subpopulation as H-type blood vessels, and the CD31loEMCNlo sinusoidal capillaries as L-type blood vessels (Kusumbe, et al., 2014). H-type capillaries are located in the vicinity of the metaphyseal growth plate and are interconnected with one another as vascular columns. High levels of CD31 and EMCN are expressed. The L-type connects to the central vein and is surrounded by dense hematopoietic cells expressing low levels of CD31 and EMCN (Ramasamy et al., 2014). Recently, the capillary network was first observed in the long bones of mice. These bone marrow-derived capillaries, which run vertically across the cortical bone and connect the bone marrow to the periosteal circulatory system, are known as trans-cortical vessels (TCVs) and play a crucial role in the endoskeletal circulatory system of mice. In fact, TCVs account for more than 80% of arterial blood flow and 59% of venous blood flow in the mouse tibia. In addition, this study also found that capillaries similar in structure to TCVs also exist in human long bones (Gruneboom et al., 2019). The complex and dense network of vascular tissue is the basis of osteogenesis in the bone microenvironment and is also a facilitator of it.

3.2 LncRNA regulates angiogenesis and neovascularization in the bone microenvironment

Bone growth is accompanied by the formation of new blood vessels, otherwise known as neovascularization, which occurs in two ways: angiogenesis and neovascularization. Angiogenesis is the process of growing new blood vessels from the existing vascular system by recruiting endothelial cells (ECs), while neovascularization is the process of the re-formation of blood vessels from bone marrow-derived endothelial progenitor cells (EPCs) (Carmeliet and Jain, 2011; Laschke et al., 2011). EPCs are precursor cells of ECs and belong to the class of pluripotent stem cells (Chopra et al., 2018). The main area of vascular endothelial cell proliferation in the bone microenvironment is the metaphysis of long bones (Ramasamy, et al., 2014). Both angiogenesis and neovascularization participate in embryonic vascular development and formation and play important

Unedited

roles in blood vessel regeneration, repair, and reconstruction (Patschan et al., 2018). Angiogenesis is a complex process including vasodilation, basement membrane degradation, endothelial cell migration, chemotaxis, increased vascular permeability, endothelial cell proliferation, and blood vessel formation (Kruger-Genge et al., 2019).

The initiation of angiogenesis is primarily triggered by VEGF. VEGF recruits and mobilizes EPCs, increases endothelial cell differentiation and proliferation, and promotes vascular tube formation (Asahara et al., 1999). Numerous studies have found that VEGF promotes the proliferation, migration, and chemotaxis of vascular endothelial cells in a variety of tissues and organs (Ferrara et al., 2003; Coultas et al., 2005; Apte et al., 2019). It has three isoforms, i.e., A, B, and C, which form homo and heterodimers and bind to the dimeric complexes of the two receptors, VEGFR-1 (Flt-1) and VEGFR-2 (Flk-1)(Ferrara, 1999; Shineh et al., 2023). Endothelial progenitor cell recruitment induces endothelial cell differentiation into stem tip cells by binding to VEGFR-2 (Carmeliet and Jain, 2011). ECs are the most sensitive to VEGF, which is determined by their higher expression of VEGFR-2 than other neighboring cells (Shimizu et al., 2023).

In recent years, numerous studies have proved that lncRNA is able to participate in the regulation of angiogenesis and neovascularization in the bone microenvironment. Among all lncRNA-regulated angiogenic signaling pathways, VEGF plays an important role and is the pivot of most signaling pathways. LncRNA TUG1 functions as a ceRNA for miR-6321, and miR-6321 inhibits EPC migration and differentiation through its target, activation transcription factor 2 (ATF2) (Yu et al., 2020). LncRNAs play important roles during angiogenesis, and the lncRNA-miRNA-mRNA regulatory pathway is a common way in which they function. Similarly, lncRNA SNHG1 is a competitive endogenous RNA for miR-140-3p, while Hypoxia induicible factor-1 alpha (HIF-1α) has been identified as a target of miR-140-3p. LncRNA SNHG1 promotes cell proliferation, tube formation, and the expression of VEGF, vascular endothelial cell calreticulin, and matrix metalloproteinase 2(MMP2) expression, while SNHG1 knockdown showed the opposite effect (Liang et al., 2020).

In addition, lncRNAs can directly regulate the expression of angiogenic factors to affect angiogenesis. Ac103746.1 is one of the highly expressed lncRNAs among genes detected in human embryonic stem cell-derived mesenchymal stem cells (HES-MSCs) and human bone marrow-derived MSCs (HBM-MSCs), and is named SCDAL (stem cell-derived angiogenic lncRNA) (Wu et al., 2021). Knockdown of SCDAL significantly reduces the angiogenic potential and repair of HES-MSCs after myocardial infarction. In contrast, SCDAL overexpression in HES-MSCs and HBM-MSCs promotes angiogenesis and cardiac function recovery. Mechanistically, SCDAL induces growth differentiation factor 6 (GDF6) expression through direct interaction with SNF5 on the GDF6 promoter. Secreted GDF6 promotes vascular endothelial cell production by activating non-canonical VEGFR2 (Wu, et al., 2021). Furthermore, SCDAL-GDF6 is expressed in human ECs and directly promotes endothelial cell angiogenesis (Wu, et al., 2021). The studies above indicate that lncRNA is essential in regulating angiogenesis and neovascularization in the bone microenvironment.

Unedited



Fig. 1 The crosstalk between osteogenesis and angiogenesis in the bone microenvironment.
A complex vascular network exists in the microenvironment of bone. LncRNAs play a crucial role in regulating interactions between ECs, EPCs, osteoclasts, osteoblasts, and BMSCs in the bone microenvironment.

Unedited

4 LncRNAs regulate the crosstalk between angiogenesis and osteogenesis in the bone microenvironment

4.1 LncRNAs affect angiogenesis by regulating osteogenic differentiation in the bone microenvironment

4.1.1 LncRNA-regulated osteogenic differentiation underlies the crosstalk between angiogenesis and osteogenesis

BMSCs are tissue repair cells that play an essential role in tissue engineering and bone development strategies. The development of osteoporosis is associated with a balance of the multidirectional differentiation of BMSCs, which are mesoderm-derived pluripotent stem cells that can differentiate into osteoblasts, chondrocytes, adipocytes, and even myoblasts. In the bone microenvironment, osteoblasts secrete angiogenic factors to regulate the proliferation and vascularization of vascular endothelial cells (Maes, et al., 2010). Osteoblasts and their precursor cells are mainly distributed around H-type vascular endothelial cells and are involved in the regulation of bone angiogenesis through the secretion of angiogenic factors such as VEGF, fibroblast growth factor (FGF), and epidermal growth factor-like members (EGFL). Osteoblasts can secrete the receptor for VEGF (VEGFR) to promote the proliferation and vascularization of vascular endothelial cells and bone angiogenesis (Potente et al., 2011). The FGF signaling pathway can also be involved in the regulation of angiogenesis and bone formation. FGF has been shown to induce the expression of vascular endothelial growth factor-A (VEGF-A) and VEGFR2 to stimulate bone arteriolar vasodilatation, and the knockdown of FGF2 in mice leads to a decrease in the rate of bone formation and the volume of bone trabeculae (Byun et al., 2014). Knockdown of FGF9 and FGF18 has been shown to reduce VEGF-A expression and decrease angiogenesis (Hung et al., 2016). Osteoblasts also regulate bone angiogenesis

through the EGFL signaling pathway. For example, osteoblasts secrete EGFL6 to promote endothelial cell proliferation and vascularization by activating ERK1/2 (Chim et al., 2011). It is widely accepted that osteoblasts are essential for the promotion of bone angiogenesis, and they primarily originate from BMSCs. The osteogenic differentiation of BMSCs is an essential process of bone formation, and lncRNA is an important regulator in this process.

Meanwhile, bone angiogenesis is highly correlated with osteogenesis. From a microscopic point of view, lncRNAs can affect angiogenesis by regulating the osteogenic differentiation of BMSCs. Recently, several studies have indicated that lncRNAs are able to modulate the proliferation and differentiation of BMSCs and regulate the balance of the multidirectional differentiation of BMSCs (Guo et al., 2020; Yu et al., 2022; Kieu Nguyen et al., 2023). The effects of lncRNAs on the osteogenic differentiation of BMSCs primarily occur in two respects, by promoting and inhibiting differentiation. Additionally, the regulatory role of specific lncRNAs in osteogenic differentiation is inconsistent across different studies.

LncRNAs regulate the osteogenic differentiation of BMSCs by targeting miRNA, the target genes of which are osteogenic factors. LncRNA MEG3 is a tumor suppressor (Zhou et al., 2012; Mitra et al., 2017) that promotes the osteogenic differentiation of MSCs in multiple myeloma patients (Zhuang et al., 2015; Li et al., 2017). LncRNA MEG3 is shown to inhibit the osteogenesis of BMSCs in postmenopausal osteoporosis by binding to miRNA-133a-3p to up-regulate miRNA-133a-3p expression, thereby down-regulating SLC39A1 expression (Wang et al., 2017). It has also been suggested that MEG3 works in conjunction with DEPTOR. DEPTOR is highly expressed in the cancellous bone and BMSCs of osteoporotic mice, and continuously decreases during the osteogenic differentiation of BMSCs. The down-regulation of DEPTOR by lentiviral transfection can promote the osteogenic differentiation of BMSCs, as shown from the results of in vivo and in vitro experiments (Chen et al., 2018). Furthermore, lncRNA and mRNA sequencing indicated that the reduction of DEPTOR up-regulated the expression of MEG3, which subsequently activated bone morphogenetic protein 4 (BMP4)/SMAD1/5/8 signaling. Chen et al. discovered that DEPTOR could bind to a specific region (−1000 bp ~ 0) of the MEG3 promoter to regulate its transcription, and the inhibition of MEG3 reduced BMP4 activation triggered by DEPTOR knockdown. The suppression of MEG3 decreases the promotion of osteogenic differentiation by DEPTOR knockdown, showing that DEPTOR can inhibit the differentiation of BMSCs toward osteogenesis through MEG3 (Chen, et al., 2018). SNHG5 promotes osteogenesis and attenuates the apoptosis of hBMSCs via regulating RUNX3 expression through the sponge adsorption of miR-582-5p (Zheng et al., 2020). The expression of lncRNA GAS5 was reduced in the hMSCs of osteoporosis patients, and the overexpression of lncRNA GAS5 was able to up-regulate RUNX2 expression by regulating miR-498 and thus promoted the osteogenic differentiation of hMSCs, which attenuates the development of osteoporosis (Feng et al., 2019). Some studies have demonstrated that lncRNA MALAT1 regulates the osteogenic differentiation of BMSCs by raising the expression of Osterix through the sponge adsorption of miR-143 and miR-96, respectively (Gao et al., 2018; Zang et al., 2022).

In addition to MEG3, SNHG5, GAS5, and MALAT1, many lncRNAs can regulate the osteogenic differentiation of BMSCs, such as HOTTIP, DANCR, and SNHG1, etc. LncRNA HOTTIP enhances osteogenic differentiation by interacting with WD40 repeat domain protein 5 (WDR5), activating the Wnt/β-catenin signaling pathway and up-regulating β-catenin gene expression (Liu et al., 2020). In contrast, lncRNA DANCR and lncRNA SNHG1 play different roles in osteogenesis. Yu et al. (Yu et al., 2021) showed in their latest findings that lncRNA-SNHG1 can bind to miRNA-181c-5p as a sponge and down-regulate the expression of miRNA-181c-5p, which in turn regulates the SFRP1/Wnt/β-catenin signaling pathway, inhibiting the osteogenic differentiation of BMSCs and promoting the pathological development of osteoporosis. In addition, in an experimental study in a postmenopausal osteoporosis mouse model, Wang et al. discovered that lncRNA-DANCR was capable of inhibiting the Wnt/β-catenin signaling pathway during the differentiation of osteoporotic BMSCs by down-regulating the expression of CTNNB1 and eliminating the up-regulatory effect of miRNA-320a on CTNNB1. This negative regulation of the osteogenic process has significant implications for understanding the role of lncRNAs in regulating bone metabolism in vivo (Wang

Unedited

et al., 2020).

Some lncRNAs are shown to regulate osteogenesis by signaling pathways such as the BMP2/Smad signaling pathway, the PI3K/AKT signaling pathway, and the p38/MAPK signaling pathway. For example, lncRNA-MSC-AS1 could act as a miRNA-140e-5p sponge and bind to it to up-regulate the expression of BMP2, which promotes the osteogenic differentiation of BMSCs and delays the progression of osteoporosis through the miRNA-140e-5p/BMP2/Smad signaling pathway (Zhang et al., 2019a). LNC_000052 inhibits apoptosis and promotes the proliferation and osteogenic differentiation of BMSCs through the miR-96-5p/PI3K/AKT signaling pathway axis (Li et al., 2020). In ovariectomized mice, lncRNA-SNHG1 negatively regulates the Nedd4-mediated p38/MAPK signaling pathway, thereby inhibiting the osteogenic differentiation of BMSCs (Jiang et al., 2019). LncRNA-DANCR is considered to reverse the osteogenic differentiation fate of BMSCs in osteoporosis and accelerate the development of osteoporosis. It could inactivate the p38/MAPK pathway, thus significantly inhibiting the proliferation and osteogenic differentiation of BMSCs (Zhang et al., 2018a).

Additionally, there are also individual lncRNAs whose role in regulating the osteogenic differentiation of BMSCs is controversial, such as lncRNA-TUG1 and lncRNA-MEG3. Some researchers analyzed plasma from osteoporosis patients by assay and found that lncRNA-TUG1 expression was decreased and miRNA-23b expression was increased. They indicated that the down-regulation of lncRNA-TUG1 could inhibit the differentiation of BMSCs by targeting the miRNA-23b/RUNX2 pathway (Teng et al., 2021). However, another study suggested that lncRNA-TUG1 could promote the differentiation of BMSCs and inhibit cell apoptosis by targeting the miRNA-34a/FGFR1 pathway, providing a potential therapeutic target for osteoporosis (Wang et al., 2021b). As mentioned earlier, lncRNA-MEG3 can inhibit the transformation of BMSCs into osteoblasts in postmenopausal osteoporosis. However, Yang et al. recently conducted an animal model study and came to the opposite conclusion (Yang et al., 2021). They found that lncRNA MEG3 can act as ceRNAs of miRNA-214 to down-regulate its expression level and thus increase the expression of TXNIP (MEG3/miRNA-214/TXNIP axis). For these controversial lncRNAs, although earlier studies have been conducted to investigate their role in BMSCs transformation, more and more studies have been undertaken in recent years with opposite conclusions.

We concluded that the main reason for the effects of the opposed regulation of the same lncRNAs is that they exert their regulatory effects through different signaling pathways. LncRNA MALAT1 in exosomes derived from BMSCs in ovariectomized osteoporotic mice can be transferred to osteoblasts, which in turn act as an miRNA-34c sponge for and up-regulate the expression of SATB2 in osteoblasts, and enhance the activity of osteoblasts, thus alleviating osteoporosis (Yang et al., 2019). However, some scholars have come to a contrary conclusion. Zheng et al. (Zheng et al., 2019) found that lncRNA-MALAT1 could enhance the MAPK signaling pathway to reduce the osteogenic differentiation of BMSCs, thus accelerating the progression of osteoporosis in the osteoporosis model mice. In addition, there are some other lncRNAs that we do not discuss in detail regarding their functions, but rather organize them in Table 1.

Table 1 Some lncRNAs in osteogenic differentiation of BMSCs

LncRNAs	Targets	Function	Reference
MEG3	miR-133a-3p/ SLC39A1	Inhibition	(Wang, et al., 2017)
SNHG5	miR-582-5p/RUNX3	Promotion	(Zheng, et al., 2020)
GAS5	miR-498/RUNX2	Promotion	(Feng, et al., 2019)
MALAT1	miR-96/Osterix	Promotion	(Zang, et al., 2022)
MALAT1	miR-143/Osterix	Promotion	(Gao, et al., 2018)
MEG3	BMP4/ Smad	Inhibition	(Chen, et al., 2018) (Tsuji et al., 2008) (Wu et al., 2016) (Liu, et al., 2020)
HOTTIP	WDR5/ Wnt/β-catenin signaling pathway	Promotion	

Unedited

SNHG1	miRNA-181c-5p, SFRP1/Wnt/β-catenin signaling pathway	Inhibition	(Yu, et al., 2021)
DANCR	CTNNB1/ Wnt/β-catenin signaling pathway	Inhibition	(Wang, et al., 2020)
MSC-AS1	miRNA-140e-5p/ BMP2/Smad signaling pathway	Promotion	(Zhang, et al., 2019a)
000052	miR-96-5p/PI3K/AKT signaling pathway	Inhibition	(Li, et al., 2020)
SNHG1	Nedd4/ p38/MAPK signaling pathway	Inhibition	(Jiang, et al., 2019)
DANCR	p38/MAPK signaling pathway	Inhibition	(Zhang, et al., 2018a)
TUG1	miRNA-23b/ RUNX2	Inhibition	(Teng, et al., 2021)
TUG1	miRNA-34a/FGFR1	Promotion	(Wang, et al., 2021b)
MEG3	miRNA-214/TXNIP	Promotion	(Yang, et al., 2021)
MALAT1	miR-34c/ SATB2	Promotion	(Yang, et al., 2019)
MALAT1	MAPK signaling pathway	Inhibition	(Zheng, et al., 2019)
ANCR	RUNX2	Inhibition	(Cai, et al., 2019)
DANCR	FOXO1	Inhibition	(Tang, et al., 2018)

4.2 LncRNAs-mediated angiogenesis regulates osteogenesis in the bone microenvironment

EPCs are thought to promote bone repair by stimulating neovascularization and osteogenesis (Atesok et al., 2010; Jules et al., 2012; Kamei et al., 2017). Pro-angiogenic factors secreted by osteoblasts, such as VEGF, can trigger signaling responses in various cells, including ECs and osteoblasts that express VEGF receptors (Novack and Teitelbaum, 2008). EPCs contribute to the formation of new blood vessels and indirectly promote new bone formation during bone repair, while VEGF has a strong chemotactic effect on osteoblasts and osteoclasts (Mayr-Wohlfart et al., 2002; Henriksen et al., 2003; Bates et al., 2017).

EPC-derived exosomes are involved in the communication between EPCs and BMSCs and promote osteoblast differentiation by suppressing the expression of osteogenic genes and promoting proliferation in vitro (Qin and Zhang, 2017). It has been shown that lncRNA-MALAT1 from the exosomes of EPCs directly binds to miR-124 and negatively regulates miR-124 expression, while integrin ITGB1 is a target of miR-124. LncRNA-MALAT1 in EPCs-derived exosomes promotes bone repair by facilitating the recruitment and differentiation of osteoclast precursors through the lnc-MALAT1/miR-124 pathway (Cui et al., 2019). Human umbilical vein endothelial cell (HUVEC)-derived exosomes can induce macrophage M2 polarization via the DDX3X/NLRP3 regulatory axis, which delivers NEAT1 to attenuate inflammation and ultimately promotes osteogenesis in vivo with the help of alginate/GelMA IPN hydrogels (Chen et al., 2023). These studies demonstrate that exosomes are one of the ways in which lncRNAs regulate the crosstalk between osteogenesis and angiogenesis in the bone microenvironment.

Unedited



Fig. 2 The role of lncRNAs in regulating the crosstalk between angiogenesis and osteogenesis in the bone microenvironment.

LncRNAs regulate the crosstalk between angiogenesis and osteogenesis coupling in the bone microenvironment through various modes of action, which are described in detail in this figure. LncRNAs can regulate ECs and EPC-dominated angiogenesis directly or indirectly by regulating the osteogenic differentiation of BMSCs. These processes are inseparable from angiogenic factors, e.g., VEGF, VEGFR, VEGFA, EGFL, and bFGF.

Unedited

4.3 LncRNAs-mediated crosstalk between angiogenesis and osteogenesis in bone diseases

4.3.1 Osteosarcoma

Osteosarcoma (OS) is extremely prevalent, especially malignant tumors occurring near the growth plate of the metaphysis of long bones or around the knee joint (Luetke et al., 2014; Huang et al., 2022). Over the last few years, an increasingly large number of studies have been focusing on the role of angiogenesis in OS progression, including the imbalance between anti-angiogenic and pro-angiogenic effects, including proliferation, migration, and metastasis (Chang and Yu, 2016). Vasculogenic mimicry is a new form of angiogenesis in tumor tissues that promotes angiogenesis, increases the supply of oxygen and nutrients to tumor tissues, and provides the basis for tumor cell metastasis (Qiao et al., 2015; Wei et al., 2017b). LncRNAs participate in vasculogenic mimicry formation in a variety of tumors, including gastric, lung, and liver cancers, among others (Luo et al., 2020). It is an unfavorable prognostic factor in cancer patients. LncRNAs are aberrantly expressed in OS and are involved in vasculogenic mimicry in OS (Ren et al., 2019). The suppression of n340532 exhibited potent anti-vasculogenic mimicry and anti-metastasis effects in vivo, suggesting its potential role in vasculogenic mimicry and metastasis in OS (Ren et al., 2019).

VEGF-A is considered a key factor in tumor-induced angiogenesis (Claesson-Welsh and Welsh, 2013). Zhang (Zhang et al., 2017) et al. found that high expression of lncRNA MALAT1 in MNNG/HOS cells resulted in the up-regulation of angiogenic factors, including VEGF-A. The LOC100129620/miR-335-3p/CDK6 signaling axis contributes to osteosarcoma metastasis by regulating osteosarcoma cell proliferation, macrophage polarization, and angiogenesis (Chen et al., 2021b). MiR-335-3p mediates the regulation of CDK6 expression by LOC100129620 and achieves the indirect promotion of angiogenesis and macrophage

polarization by LOC100129620. LINC00265 may promote the metastatic effects of LOC100129620 on angiogenesis and macrophage polarization in osteosarcoma by targeting miR-382-5p/SAT1 (spermine N1-acetyltransferase-1) and miR-382-5p/VAV3 (vav guanine nucleotide exchange factor 3) to promote proliferation, migration, invasion, and angiogenesis in osteosarcoma (Xiao et al., 2020). Exosomes derived from serum transport lncRNA MIAT (EV-MIAT) into osteosarcoma cells to compete for binding to miR-613, thus up-regulating GPR158 and thereby promoting osteosarcoma cell proliferation and angiogenesis (Wang et al., 2022a). Exosome-derived lncRNA NORAD, from BMSCs, regulates CREBBP through miR-877-3p to promote OS cell proliferation, invasion, migration, and angiogenesis (Feng et al., 2022). EWSAT1 regulates OS-induced angiogenesis via two pathways, called the “double stacking effect”, which is a combination of the increase in the sensitivity of vascular endothelial cells induced by exosomes carrying EWSAT1 and the EWSAT1-induced increase in angiogenic factor secretion from OS cells (Tao et al., 2020). EWSAT1 regulates Akt and Erk signaling pathways through the EWSAT1/miR-326/KRAS ceRNA pathway, significantly promoting OS proliferation, migration, colony formation, and survival (Tao et al., 2020). The above-mentioned studies provide new ideas for the study of the pathogenesis of angiogenesis in osteosarcoma.

4.3.2 Osteoarthritis

Angiogenesis is involved in almost every process of osteoarthritis, including synovitis caused by the appearance of neovascularization in the synovium, cartilage destruction caused by the appearance of neovascularization at the osteochondral joint, and the formation of bone capillaries, all of which are closely related to angiogenesis (Ashraf et al., 2011). VEGF is considered a key factor in angiogenesis. Inflammatory factors (IL-1β, TNF-α), hypoxia, and mechanical stress all increase the expression of VEGF in OA joints through multiple signaling pathways (Ashraf and Walsh, 2008). VEGF expression in the surface, middle, and deep layers of OA cartilage have been shown to be up-regulated, whereas angiogenesis occurs predominantly in the deeper cartilage (Mapp and Walsh, 2012). Angiogenesis is an important part of the cartilage's osteogenesis process, leading to subchondral bone remodeling, synovial hyperplasia, and osteophytes (He et al., 2021). MEG3 is significantly down-regulated in the articular cartilage of osteoarthritis patients, while VEGF expression is significantly increased (Su et al., 2015). MEG3 promotes the transcription of p53, which inhibits VEGF expression by binding to Sp1, a transcription site on the VEGF promoter (Kim et al., 2011; Zhang et al., 2019b). The down-regulation of MEG3 may indirectly promote VEGF production and endochondral vascularization through this pathway. LncRNA H19 is a gene that regulates chondrocyte and osteoblast growth as well as cellular anabolic processes (He et al., 2016). Some studies have shown that lncRNA H19 is highly expressed in the peripheral blood of osteoarthritis patients, which is expected to be a new indicator for the diagnosis of osteoarthritis (Zhou et al., 2020). Exosomal H19 acts as a “sponge” to absorb miR-106, thereby up-regulating the expression of Angpt1, an angiogenic factor that activates H19/Tie2-NO signaling in mesenchymal and ECs (Behera et al., 2021). In hypertrophic chondrocytes, lncRNA HCAR promotes endochondral bone repair by competitively binding to miR-15b-5p to increase the expression of matrix metallopeptidase 13 (MMP13) and VEGFA (Bai et al., 2022). In a bone repair model, lncRNA HCAR knockdown in hypertrophied chondrocytes inhibited cartilage matrix remodeling and reduced the number of CD31hiEmcnhi vessels (Bai, et al., 2022).

4.3.3 Fractures

As a clinical diagnosis, fracture nonunion is characterized by patients with fractures exhibiting clinical signs of pain at the fracture site. At the same time, the radiologic findings are consistent with bone nonunion (Hankenson et al., 2011). New blood vessels bring oxygen and nutrients to the highly metabolically active regenerating bone scab, providing a pathway for inflammatory cells, cartilage, and bone precursor cells to reach the injury site (Hankenson et al., 2011). The normal development of blood vessels in the fracture scab is an important part of fracture rehabilitation. LncRNA STEEL expression and vascular density in bone scab tissues of fracture model mice were significantly lower than those of controls; meanwhile, it was verified with

Unedited

in vitro experiments that the proliferative capacity of HUVEC cells was significantly reduced after the low expression of STEEL in HUVECs (Zhang et al., 2018c). This is due to the fact that STEEL can promote angiogenesis by interacting with PARP-1 and up-regulating the expression of VEGF (Zhang et al., 2018c). In a study on osteodysplasia, the expression of lncRNA ENST00000563492 was down-regulated in osteodysplasia tissues, which could promote the differentiation of BMSCs towards osteogenesis by up-regulating the expression of CDH11 and could also improve the process of osteogenic–angiogenic coupling by enhancing the expression of VEGF during the process of the osteogenic differentiation of BMSCs. CDH11 and VEGF are the target genes of miR-205-5p, so LncRNA ENST00000563492 indirectly regulates the expression of both of them as the ceRNA of miR-205-5p, and can promote the osteogenesis of BMSCs in vivo, so it is said that lncRNA ENST00000563492 may become a new target of osteodysplasia (Ouyang et al., 2020). The application of normal-source BMSCs exosomal-lncRNA H19 competitively binding to miR-467 regulated the expression of Hoxa10 and inhibited poor fracture healing caused by a high-fat diet (Wang et al., 2021c). Wei et al. (Wei et al., 2017a) demonstrated, in a study related to non-traumatic femoral head necrosis, that lncRNA HOTAIR was found to regulate the expression of miR-17-5p and its target gene Smad7 through the BMPs/TGF-β pathway to inhibit osteogenic differentiation.

5 Summary and Prospects

Angiogenesis is a continuous process that occurs throughout the life cycle of bone from generation to growth. Angiogenesis and osteogenesis in the bone microenvironment are not independent parts but interact with each other. LncRNAs regulate the crosstalk between angiogenesis and osteogenesis through multiple miRNAs, mRNAs, and signaling pathways, mainly in osteogenesis, osteoclastogenesis, and angiogenesis. Meanwhile, the roles of some lncRNAs in osteogenesis and angiogenesis are subject to controversy due to conflicting findings from various studies, resulting from the progressively deeper research into lncRNAs. However, numerous queries necessitate further scientific and comprehensive investigations. For instance, can the findings in this field substantiate the clinical intervention for osteoporosis? It is suggested that more researchers concentrate on simulating the bone microenvironment in vitro to achieve multi-cell co-culture. It is even possible to incorporate the latest reactive biomaterials into encapsulating lncRNAs and stimulate their release at distinct stages of osteogenesis or angiogenesis, aiming to treat various bone-related diseases. Therefore, in the future, the advancement toward the modulation of bone-related illnesses through the use of lncRNAs promises a vibrant field of research.

Unedited

Acknowledgments

This work is supported by the National Natural Science Foundation of China(No. 81901430), Guangdong Basic and Applied Basic Research Foundation (No. 2022A1515010379), the Science and Technology Projects in Guangzhou(No. 2023A04J0555), the Guangdong Provincial Key Laboratory of Physical Activity and Health Promotion (2021B1212040014).

Author contributions

Shihua Zhang performed conceptualization, formal analysis, writing—original draft preparation, review and editing; Jianmin Guo performed writing—original draft preparation, review and editing;Yuting He and Zhiang Su performed validation, review and editing; Yao Feng performed formal analysis, review and editing; Lan Zhang and Jun Zou performed funding acquisition, supervision, writing-original draft, writing-review and editing; Xiquan Weng and Yu Yuan performed funding acquisition, supervision, writing-review and editing, project administration. All authors read and approved the final manuscript and, therefore, had full access to all the data in the study and took responsibility for the integrity and security of the data.

Compliance with ethics guidelines

Shihua ZHANG, Jianmin Guo, Yuting He, Zhiang Su, Yao Feng, Lan Zhang, Jun Zou, Xiquan Weng and Yu Yuan declare that they have no conflicts of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

Alagiakrishnan K, Juby A, Hanley D, et al., 2003. Role of vascular factors in osteoporosis. *J Gerontol A Biol Sci Med Sci*, 58(4):362-366. https://doi.org/10.1093/gerona/58.4.m362

Allas L, Boumediene K, Bauge C, 2019. Epigenetic dynamic during endochondral ossification and articular cartilage development. *Bone*, 120:523-532. https://doi.org/10.1016/j.bone.2018.10.004

Apte RS, Chen DS, Ferrara N, 2019. Vegf in signaling and disease: Beyond discovery and development. *Cell*, 176(6):1248-1264. https://doi.org/10.1016/j.cell.2019.01.021

Asahara T, Takahashi T, Masuda H, et al., 1999. Vegf contributes to postnatal neovascularization by mobilizing bone marrow-derived endothelial progenitor cells. *EMBO J*, 18(14):3964-3972. https://doi.org/10.1093/emboj/18.14.3964

Ashraf S, Walsh DA, 2008. Angiogenesis in osteoarthritis. *Curr Opin Rheumatol*, 20(5):573-580. https://doi.org/10.1097/BOR.0b013e3283103d12

Ashraf S, Mapp PI, Walsh DA, 2011. Contributions of angiogenesis to inflammation, joint damage, and pain in a rat model of osteoarthritis. *Arthritis Rheum*, 63(9):2700-2710. https://doi.org/10.1002/art.30422

Atesok K, Li R, Stewart DJ, et al., 2010. Endothelial progenitor cells promote fracture healing in a segmental bone defect model. *J Orthop Res*, 28(8):1007-1014. https://doi.org/10.1002/jor.21083

Bahney CS, Zondervan RL, Allison P, et al., 2019. Cellular biology of fracture healing. *J Orthop Res*, 37(1):35-50. https://doi.org/10.1002/jor.24170

Bai Y, Gong X, Dong R, et al., 2022. Long non-coding rna hcar promotes endochondral bone repair by upregulating vegf and mmp13 in hypertrophic chondrocyte through sponging mir-15b-5p. *Genes Dis*, 9(2):456-465. https://doi.org/10.1016/j.gendis.2020.07.013

Baker CE, Moore-Lotridge SN, Hysong AA, et al., 2018. Bone fracture acute phase response-a unifying theory of fracture repair: Clinical and scientific implications. *Clin Rev Bone Miner Metab*, 16(4):142-158. https://doi.org/10.1007/s12018-018-9256-x

Bartel DP, 2004. Micrornas: Genomics, biogenesis, mechanism, and function. *Cell*, 116(2):281-297. https://doi.org/10.1016/s0092-8674(04)00045-5

Bates BD, Godbout C, Ramnaraign DJ, et al., 2017. Delayed endothelial progenitor cell therapy promotes bone defect repair in a clinically relevant rat model. *Stem Cells Int*, 2017:7923826. https://doi.org/10.1155/2017/7923826

Behera J, Kumar A, Voor MJ, et al., 2021. Exosomal lncrna-h19 promotes osteogenesis and angiogenesis through mediating angpt1/tie2-no signaling in cbs-heterozygous mice. *Theranostics*, 11(16):7715-7734. https://doi.org/10.7150/thno.58410

Bellavia D, De Luca A, Carina V, et al., 2019. Deregulated mirnas in bone health: Epigenetic roles in osteoporosis. *Bone*, 122:52-75. https://doi.org/10.1016/j.bone.2019.02.013

Byun MR, Kim AR, Hwang JH, et al., 2014. Fgf2 stimulates osteogenic differentiation through erk induced taz expression. *Bone*, 58:72-80. https://doi.org/10.1016/j.bone.2013.09.024

Cai N, Li C, Wang F, 2019. Silencing of lncrna-ancr promotes the osteogenesis of osteoblast cells in postmenopausal osteoporosis via targeting ezh2 and runx2. *Yonsei Med J*, 60(8):751-759. https://doi.org/10.3349/ymj.2019.60.8.751

Carmeliet P, Jain RK, 2011. Molecular mechanisms and clinical applications of angiogenesis. *Nature*, 473(7347):298-307. https://doi.org/10.1038/nature10144

Chang LC, Yu YL, 2016. Dietary components as epigenetic-regulating agents against cancer. *Biomedicine (Taipei)*, 6(1):2. https://doi.org/10.7603/s40681-016-0002-8

Chen K, Zhu H, Zheng MQ, et al., 2021a. Lncrna meg3 inhibits the degradation of the extracellular matrix of chondrocytes in osteoarthritis via targeting mir-93/tgfbr2 axis. *Cartilage*, 13(2_suppl):1274S-1284S. https://doi.org/10.1177/1947603519855759

Chen S, Jia L, Zhang S, et al., 2018. Deptor regulates osteogenic differentiation via inhibiting meg3-mediated activation of bmp4 signaling and is involved in osteoporosis. *Stem Cell Res Ther*, 9(1):185. https://doi.org/10.1186/s13287-018-0935-9

Chen Y, Tang G, Qian H, et al., 2021b. Lncrna loc100129620 promotes osteosarcoma progression through regulating cdk6 expression, tumor angiogenesis, and macrophage polarization. *Aging (Albany NY)*, 13(10):14258-14276. https://doi.org/10.18632/aging.203042

Chen Y, Wu Y, Guo L, et al., 2023. Exosomal lnc neat1 from endothelial cells promote bone regeneration by regulating macrophage polarization via ddx3x/nlrp3 axis. *J Nanobiotechnology*, 21(1):98. https://doi.org/10.1186/s12951-023-01855-w

Chim SM, Qin A, Tickner J, et al., 2011. Egfl6 promotes endothelial cell migration and angiogenesis through the activation of extracellular signal-regulated kinase. *J Biol Chem*, 286(25):22035-22046. https://doi.org/10.1074/jbc.M110.187633

Chim SM, Tickner J, Chow ST, et al., 2013. Angiogenic factors in bone local environment. *Cytokine Growth Factor Rev*, 24(3):297-310. https://doi.org/10.1016/j.cytogfr.2013.03.008

Chopra H, Hung MK, Kwong DL, et al., 2018. Insights into endothelial progenitor cells: Origin, classification, potentials, and prospects. *Stem Cells Int*, 2018:9847015. https://doi.org/10.1155/2018/9847015

Claesson-Welsh L, Welsh M, 2013. Vegfa and tumour angiogenesis. *J Intern Med*, 273(2):114-127. https://doi.org/10.1111/joim.12019

Coultas L, Chawengsaksophak K, Rossant J, 2005. Endothelial cells and vegf in vascular development. *Nature*, 438(7070):937-

Unedited

945. https://doi.org/10.1038/nature04479
Cui Y, Fu S, Sun D, et al., 2019. Epc-derived exosomes promote osteoclastogenesis through lncrna-malat1. *J Cell Mol Med*, 23(6):3843-3854. https://doi.org/10.1111/jcmm.14228
Dangwal S, Schimmel K, Foinquinos A, et al., 2017. Noncoding rnas in heart failure. *Handb Exp Pharmacol*, 243:423-445. https://doi.org/10.1007/164_2016_99
Ding A, Li CH, Yu CY, et al., 2021. Long non-coding rna malat1 enhances angiogenesis during bone regeneration by regulating the mir-494/sp1 axis. *Lab Invest*, 101(11):1458-1466. https://doi.org/10.1038/s41374-021-00649-8
Ding WG, Wei ZX, Liu JB, 2011. Reduced local blood supply to the tibial metaphysis is associated with ovariectomy-induced osteoporosis in mice. *Connect Tissue Res*, 52(1):25-29. https://doi.org/10.3109/03008201003783011
Dirckx N, Van Hul M, Maes C, 2013. Osteoblast recruitment to sites of bone formation in skeletal development, homeostasis, and regeneration. *Birth Defects Res C Embryo Today*, 99(3):170-191. https://doi.org/10.1002/bdrc.21047
Duret L, Chureau C, Samain S, et al., 2006. The xist rna gene evolved in eutherians by pseudogenization of a protein-coding gene. *Science*, 312(5780):1653-1655. https://doi.org/10.1126/science.1126316
Feng D, Li Z, Yang L, et al., 2022. Bmsc-ev-derived lncrna norad facilitates migration, invasion, and angiogenesis in osteosarcoma cells by regulating crebbp via delivery of mir-877-3p. *Oxid Med Cell Longev*, 2022:8825784. https://doi.org/10.1155/2022/8825784
Feng J, Wang JX, Li CH, 2019. Lncrna gas5 overexpression alleviates the development of osteoporosis through promoting osteogenic differentiation of mscs via targeting microrna-498 to regulate runx2. *Eur Rev Med Pharmacol Sci*, 23(18):7757-7765. https://doi.org/10.26355/eurrev_201909_18985
Ferrara N, 1999. Molecular and biological properties of vascular endothelial growth factor. *J Mol Med (Berl)*, 77(7):527-543. https://doi.org/10.1007/s001099900019
Ferrara N, Gerber HP, Lecouter J, 2003. The biology of vegf and its receptors. *Nat Med*, 9(6):669-676. https://doi.org/10.1038/nm0603-669
Gao Y, Xiao F, Wang C, et al., 2018. Long noncoding rna malat1 promotes osterix expression to regulate osteogenic differentiation by targeting mirna-143 in human bone marrow-derived mesenchymal stem cells. *J Cell Biochem*, 119(8):6986-6996. https://doi.org/10.1002/jcb.26907
Ghafouri-Fard S, Abak A, Tavakkoli Avval S, et al., 2021. Contribution of mirnas and lncrnas in osteogenesis and related disorders. *Biomed Pharmacother*, 142:111942. https://doi.org/10.1016/j.biopha.2021.111942
Grosso A, Burger MG, Lunger A, et al., 2017. It takes two to tango: Coupling of angiogenesis and osteogenesis for bone regeneration. *Front Bioeng Biotechnol*, 5:68. https://doi.org/10.3389/fbioe.2017.00068
Gruneboom A, Hawwari I, Weidner D, et al., 2019. A network of trans-cortical capillaries as mainstay for blood circulation in long bones. *Nat Metab*, 1(2):236-250. https://doi.org/10.1038/s42255-018-0016-5
Guo Q, Guo Q, Xiao Y, et al., 2020. Regulation of bone marrow mesenchymal stem cell fate by long non-coding rna. *Bone*, 141:115617. https://doi.org/10.1016/j.bone.2020.115617
Han P, Li W, Lin CH, et al., 2014. A long noncoding rna protects the heart from pathological hypertrophy. *Nature*, 514(7520):102-106. https://doi.org/10.1038/nature13596
Hankenson KD, Dishowitz M, Gray C, et al., 2011. Angiogenesis in bone regeneration. *Injury*, 42(6):556-561. https://doi.org/10.1016/j.injury.2011.03.035
Harada S, Rodan GA, 2003. Control of osteoblast function and regulation of bone mass. *Nature*, 423(6937):349-355. https://doi.org/10.1038/nature01660
He CP, Jiang XC, Chen C, et al., 2021. The function of lncrnas in the pathogenesis of osteoarthritis. *Bone Joint Res*, 10(2):122-133. https://doi.org/10.1302/2046-3758.102.BJR-2020-0228.R1
He P, Zhang Z, Huang G, et al., 2016. Mir-141 modulates osteoblastic cell proliferation by regulating the target gene of lncrna h19 and lncrna h19-derived mir-675. *Am J Transl Res*, 8(4):1780-1788.
He X, Dziak R, Yuan X, et al., 2013. Bmp2 genetically engineered mscs and epcs promote vascularized bone regeneration in rat critical-sized calvarial bone defects. *PLoS One*, 8(4):e60473. https://doi.org/10.1371/journal.pone.0060473
Henriksen K, Karsdal M, Delaisse JM, et al., 2003. Rankl and vascular endothelial growth factor (vegf) induce osteoclast chemotaxis through an erk1/2-dependent mechanism. *J Biol Chem*, 278(49):48745-48753. https://doi.org/10.1074/jbc.M309193200
Herman AB, Tsitsipatis D, Gorospe M, 2022. Integrated lncrna function upon genomic and epigenomic regulation. *Mol Cell*, 82(12):2252-2266. https://doi.org/10.1016/j.molcel.2022.05.027
Huang X, Wu W, Jing D, et al., 2022. Engineered exosome as targeted lncrna meg3 delivery vehicles for osteosarcoma therapy. *J Control Release*, 343:107-117. https://doi.org/10.1016/j.jconrel.2022.01.026
Hung IH, Schoenwolf GC, Lewandoski M, et al., 2016. A combined series of fgf9 and fgf18 mutant alleles identifies unique and redundant roles in skeletal development. *Dev Biol*, 411(1):72-84. https://doi.org/10.1016/j.ydbio.2016.01.008
Jiang Y, Wu W, Jiao G, et al., 2019. Lncrna snhg1 modulates p38 mapk pathway through nedd4 and thus inhibits osteogenic differentiation of bone marrow mesenchymal stem cells. *Life Sci*, 228:208-214. https://doi.org/10.1016/j.lfs.2019.05.002
Jing H, Liao L, Su X, et al., 2017. Declining histone acetyltransferase gcn5 represses bmsc-mediated angiogenesis during osteoporosis. *FASEB J*, 31(10):4422-4433. https://doi.org/10.1096/fj.201700118R
Jules J, Zhang P, Ashley JW, et al., 2012. Molecular basis of requirement of receptor activator of nuclear factor kappab signaling for interleukin 1-mediated osteoclastogenesis. *J Biol Chem*, 287(19):15728-15738.

Unedited

https://doi.org/10.1074/jbc.M111.296228
Kamei N, Atesok K, Ochi M, 2017. The use of endothelial progenitor cells for the regeneration of musculoskeletal and neural tissues. *Stem Cells Int*, 2017:1960804. https://doi.org/10.1155/2017/1960804
Karreth FA, Tay Y, Perna D, et al., 2011. In vivo identification of tumor- suppressive pten cernas in an oncogenic braf-induced mouse model of melanoma. *Cell*, 147(2):382-395. https://doi.org/10.1016/j.cell.2011.09.032
Kieu Nguyen NT, Tu Y, Lee HS, et al., 2023. Split dcas12a activator for lncrna h19 activation to enhance bmsc differentiation and promote calvarial bone healing. *Biomaterials*, 297:122106. https://doi.org/10.1016/j.biomaterials.2023.122106
Kim DH, Marinov GK, Pepke S, et al., 2015. Single-cell transcriptome analysis reveals dynamic changes in lncrna expression during reprogramming. *Cell Stem Cell*, 16(1):88-101. https://doi.org/10.1016/j.stem.2014.11.005
Kim HA, Seo KH, Kang YR, et al., 2011. Mechanisms of platelet-activating factor-induced enhancement of vegf expression. *Cell Physiol Biochem*, 27(1):55-62. https://doi.org/10.1159/000325205
Kruger-Genge A, Blocki A, Franke RP, et al., 2019. Vascular endothelial cell biology: An update. *Int J Mol Sci*, 20(18) https://doi.org/10.3390/ijms20184411
Kung JT, Colognori D, Lee JT, 2013. Long noncoding rnas: Past, present, and future. *Genetics*, 193(3):651-669. https://doi.org/10.1534/genetics.112.146704
Kusumbe AP, Ramasamy SK, Adams RH, 2014. Coupling of angiogenesis and osteogenesis by a specific vessel subtype in bone. *Nature*, 507(7492):323-328. https://doi.org/10.1038/nature13145
Lafage-Proust MH, Prisby R, Roche B, et al., 2010. Bone vascularization and remodeling. *Joint Bone Spine*, 77(6):521-524. https://doi.org/10.1016/j.jbspin.2010.09.009
Laschke MW, Giebels C, Menger MD, 2011. Vasculogenesis: A new piece of the endometriosis puzzle. *Hum Reprod Update*, 17(5):628-636. https://doi.org/10.1093/humupd/dmr023
Li CJ, Cheng P, Liang MK, et al., 2015. Microrna-188 regulates age-related switch between osteoblast and adipocyte differentiation. *J Clin Invest*, 125(4):1509-1522. https://doi.org/10.1172/JCI77716
Li M, Cong R, Yang L, et al., 2020. A novel lncrna lnc_000052 leads to the dysfunction of osteoporotic bmscs via the mir-96-5p-pik3r1 axis. *Cell Death Dis*, 11(9):795. https://doi.org/10.1038/s41419-020-03006-7
Li Z, Jin C, Chen S, et al., 2017. Long non-coding rna meg3 inhibits adipogenesis and promotes osteogenesis of human adipose-derived mesenchymal stem cells via mir-140-5p. *Mol Cell Biochem*, 433(1-2):51-60. https://doi.org/10.1007/s11010-017-3015-z
Li Z, Lu Q, Zhu D, et al., 2018. Lnc-snhg1 may promote the progression of non-small cell lung cancer by acting as a sponge of mir-497. *Biochem Biophys Res Commun*, 506(3):632-640. https://doi.org/10.1016/j.bbrc.2018.10.086
Liang S, Ren K, Li B, et al., 2020. Lncrna snhg1 alleviates hypoxia-reoxygenation-induced vascular endothelial cell injury as a competing endogenous rna through the hif-1alpha/vegf signal pathway. *Mol Cell Biochem*, 465(1-2):1-11. https://doi.org/10.1007/s11010-019-03662-0
Liu R, Li Z, Song E, et al., 2020. Lncrna hottip enhances human osteogenic bmscs differentiation via interaction with wdr5 and activation of wnt/beta-catenin signalling pathway. *Biochem Biophys Res Commun*, 524(4):1037-1043. https://doi.org/10.1016/j.bbrc.2020.02.034
Luetke A, Meyers PA, Lewis I, et al., 2014. Osteosarcoma treatment - where do we stand? A state of the art review. *Cancer Treat Rev*, 40(4):523-532. https://doi.org/10.1016/j.ctrv.2013.11.006
Luo Q, Wang J, Zhao W, et al., 2020. Vasculogenic mimicry in carcinogenesis and clinical applications. *J Hematol Oncol*, 13(1):19. https://doi.org/10.1186/s13045-020-00858-6
Ma Q, Miri Z, Haugen HJ, et al., 2023. Significance of mechanical loading in bone fracture healing, bone regeneration, and vascularization. *J Tissue Eng*, 14:20417314231172573. https://doi.org/10.1177/20417314231172573
Maes C, Kobayashi T, Selig MK, et al., 2010. Osteoblast precursors, but not mature osteoblasts, move into developing and fractured bones along with invading blood vessels. *Dev Cell*, 19(2):329-344. https://doi.org/10.1016/j.devcel.2010.07.010
Maes C, 2013. Role and regulation of vascularization processes in endochondral bones. *Calcif Tissue Int*, 92(4):307-323. https://doi.org/10.1007/s00223-012-9689-z
Mapp PI, Walsh DA, 2012. Mechanisms and targets of angiogenesis and nerve growth in osteoarthritis. *Nat Rev Rheumatol*, 8(7):390-398. https://doi.org/10.1038/nrrheum.2012.80
Mattick JS, 2018. The state of long non-coding rna biology. *Noncoding RNA*, 4(3) https://doi.org/10.3390/ncrna4030017
Mayr-Wohlfart U, Waltenberger J, Hausser H, et al., 2002. Vascular endothelial growth factor stimulates chemotactic migration of primary human osteoblasts. *Bone*, 30(3):472-477. https://doi.org/10.1016/s8756-3282(01)00690-1
Mitra R, Chen X, Greenawalt EJ, et al., 2017. Decoding critical long non-coding rna in ovarian cancer epithelial-to-mesenchymal transition. *Nat Commun*, 8(1):1604. https://doi.org/10.1038/s41467-017-01781-0
Nojima T, Proudfoot NJ, 2022. Mechanisms of lncrna biogenesis as revealed by nascent transcriptomics. *Nat Rev Mol Cell Biol*, 23(6):389-406. https://doi.org/10.1038/s41580-021-00447-6
Novack DV, Teitelbaum SL, 2008. The osteoclast: Friend or foe? *Annu Rev Pathol*, 3:457-484. https://doi.org/10.1146/annurev.pathmechdis.3.121806.151431
Ouyang Z, Tan T, Zhang X, et al., 2020. Lncrna enst00000563492 promoting the osteogenesis-angiogenesis coupling process in bone mesenchymal stem cells (bmscs) by functions as a cerna for mir-205-5p. *Cell Death Dis*, 11(6):486. https://doi.org/10.1038/s41419-020-2689-4
Patschan D, Sugiarto N, Henze E, et al., 2018. Early endothelial progenitor cells and vascular stiffness in psoriasis and psoriatic

Unedited

arthritis. *Eur J Med Res*, 23(1):56. https://doi.org/10.1186/s40001-018-0352-7
Peng J, Lai ZG, Fang ZL, et al., 2014. Dimethyloxalylglycine prevents bone loss in ovariectomized c57bl/6j mice through enhanced angiogenesis and osteogenesis. *PLoS One*, 9(11):e112744. https://doi.org/10.1371/journal.pone.0112744
Potente M, Gerhardt H, Carmeliet P, 2011. Basic and therapeutic aspects of angiogenesis. *Cell*, 146(6):873-887. https://doi.org/10.1016/j.cell.2011.08.039
Qiao L, Liang N, Zhang J, et al., 2015. Advanced research on vasculogenic mimicry in cancer. *J Cell Mol Med*, 19(2):315-326. https://doi.org/10.1111/jcmm.12496
Qin Y, Zhang C, 2017. Endothelial progenitor cell-derived extracellular vesicle-meditated cell-to-cell communication regulates the proliferation and osteoblastic differentiation of bone mesenchymal stromal cells. *Mol Med Rep*, 16(5):7018-7024. https://doi.org/10.3892/mmr.2017.7403
Ramasamy SK, Kusumbe AP, Wang L, et al., 2014. Endothelial notch activity promotes angiogenesis and osteogenesis in bone. *Nature*, 507(7492):376-380. https://doi.org/10.1038/nature13146
Ren K, Ni Y, Li X, et al., 2019. Expression profiling of long noncoding rnas associated with vasculogenic mimicry in osteosarcoma. *J Cell Biochem*, 120(8):12473-12488. https://doi.org/10.1002/jcb.28514
Saran U, Gemini Piperni S, Chatterjee S, 2014. Role of angiogenesis in bone repair. *Arch Biochem Biophys*, 561:109-117. https://doi.org/10.1016/j.abb.2014.07.006
Schipani E, Maes C, Carmeliet G, et al., 2009. Regulation of osteogenesis-angiogenesis coupling by hifs and vegf. *J Bone Miner Res*, 24(8):1347-1353. https://doi.org/10.1359/jbmr.090602
Shimizu S, Iba T, Naito H, et al., 2023. Aging impairs the ability of vascular endothelial stem cells to generate endothelial cells in mice. *Angiogenesis*, https://doi.org/10.1007/s10456-023-09891-8
Shineh G, Patel K, Mobaraki M, et al., 2023. Functional approaches in promoting vascularization and angiogenesis in bone critical-sized defects via delivery of cells, growth factors, drugs, and particles. *J Funct Biomater*, 14(2) https://doi.org/10.3390/jfb14020099
Smart F, Aschrafi A, Atkins A, et al., 2007. Two isoforms of the cold-inducible mrna-binding protein rbm3 localize to dendrites and promote translation. *J Neurochem*, 101(5):1367-1379. https://doi.org/10.1111/j.1471-4159.2007.04521.x
St Laurent G, Wahlestedt C, Kapranov P, 2015. The landscape of long noncoding rna classification. *Trends Genet*, 31(5):239-251. https://doi.org/10.1016/j.tig.2015.03.007
Su W, Xie W, Shang Q, et al., 2015. The long noncoding rna meg3 is downregulated and inversely associated with vegf levels in osteoarthritis. *Biomed Res Int*, 2015:356893. https://doi.org/10.1155/2015/356893
Tang Z, Gong Z, Sun X, 2018. Lncrna dancr involved osteolysis after total hip arthroplasty by regulating foxo1 expression to inhibit osteoblast differentiation. *J Biomed Sci*, 25(1):4. https://doi.org/10.1186/s12929-018-0406-8
Tao SC, Huang JY, Wei ZY, et al., 2020. Ewsat1 acts in concert with exosomes in osteosarcoma progression and tumor-induced angiogenesis: The "double stacking effect". *Adv Biosyst*, 4(9):e2000152. https://doi.org/10.1002/adbi.202000152
Tay Y, Kats L, Salmena L, et al., 2011. Coding-independent regulation of the tumor suppressor pten by competing endogenous mrnas. *Cell*, 147(2):344-357. https://doi.org/10.1016/j.cell.2011.09.029
Teng Z, Zhu Y, Hao Q, et al., 2021. Long non-coding rna taurine upregulated gene 1 is downregulated in osteoporosis and influences the osteogenic differentiation of bone marrow mesenchymal stem cells. *PeerJ*, 9:e11251. https://doi.org/10.7717/peerj.11251
Tian T, Qiu R, Qiu X, 2018. Snhg1 promotes cell proliferation by acting as a sponge of mir-145 in colorectal cancer. *Oncotarget*, 9(2):2128-2139. https://doi.org/10.18632/oncotarget.23255
Tsai MH, Megat Abdul Wahab R, Zainal Ariffin SH, et al., 2023. Enhanced osteogenesis potential of mg-63 cells through sustained delivery of vegf via liposomal hydrogel. *Gels*, 9(7) https://doi.org/10.3390/gels9070562
Tsuji K, Cox K, Bandyopadhyay A, et al., 2008. Bmp4 is dispensable for skeletogenesis and fracture-healing in the limb. *J Bone Joint Surg Am*, 90 Suppl 1:14-18. https://doi.org/10.2106/JBJS.G.01109
Wang BD, Yu XJ, Hou JC, et al., 2022a. Bevacizumab attenuates osteosarcoma angiogenesis by suppressing miat encapsulated by serum-derived extracellular vesicles and facilitating mir-613-mediated gpr158 inhibition. *Cell Death Dis*, 13(3):272. https://doi.org/10.1038/s41419-022-04620-3
Wang CG, Hu YH, Su SL, et al., 2020. Lncrna dancr and mir-320a suppressed osteogenic differentiation in osteoporosis by directly inhibiting the wnt/beta-catenin signaling pathway. *Exp Mol Med*, 52(8):1310-1325. https://doi.org/10.1038/s12276-020-0475-0
Wang MQ, Zhu WJ, Gao P, 2021a. New insights into long non-coding rnas in breast cancer: Biological functions and therapeutic prospects. *Exp Mol Pathol*, 120:104640. https://doi.org/10.1016/j.yexmp.2021.104640
Wang Q, Li Y, Zhang Y, et al., 2017. Lncrna meg3 inhibited osteogenic differentiation of bone marrow mesenchymal stem cells from postmenopausal osteoporosis by targeting mir-133a-3p. *Biomed Pharmacother*, 89:1178-1186. https://doi.org/10.1016/j.biopha.2017.02.090
Wang S, Xiong G, Ning R, et al., 2022b. Lncrna meg3 promotes osteogenesis of hbmscs by regulating mir-21-5p / sod3 axis. *Acta Biochim Pol*, 69(1):71-77. https://doi.org/10.18388/abp.2020_5661
Wang X, Guo B, Li Q, et al., 2013. Mir-214 targets atf4 to inhibit bone formation. *Nat Med*, 19(1):93-100. https://doi.org/10.1038/nm.3026
Wang X, He J, Wang H, et al., 2021b. Fluid shear stress regulates osteoblast proliferation and apoptosis via the lncrna tug1/mir-34a/fgfr1 axis. *J Cell Mol Med*, 25(18):8734-8747. https://doi.org/10.1111/jcmm.16829
Wang Y, Chen W, Zhao L, et al., 2021c. Obesity regulates mir-467/hoxa10 axis on osteogenic differentiation and fracture healing

Unedited

by bmsc-derived exosome lncrna h19. *J Cell Mol Med*, 25(3):1712-1724. https://doi.org/10.1111/jcmm.16273

Wei B, Wei W, Zhao B, et al., 2017a. Long non-coding rna hotair inhibits mir-17-5p to regulate osteogenic differentiation and proliferation in non-traumatic osteonecrosis of femoral head. *PLoS One*, 12(2):e0169097. https://doi.org/10.1371/journal.pone.0169097

Wei H, Wang F, Wang Y, et al., 2017b. Verteporfin suppresses cell survival, angiogenesis and vasculogenic mimicry of pancreatic ductal adenocarcinoma via disrupting the yap-tead complex. *Cancer Sci*, 108(3):478-487. https://doi.org/10.1111/cas.13138

Wilusz JE, Sunwoo H, Spector DL, 2009. Long noncoding rnas: Functional surprises from the rna world. *Genes Dev*, 23(13):1494-1504. https://doi.org/10.1101/gad.1800909

Wu M, Chen G, Li YP, 2016. Tgf-beta and bmp signaling in osteoblast, skeletal development, and bone formation, homeostasis and disease. *Bone Res*, 4:16009. https://doi.org/10.1038/boneres.2016.9

Wu R, Hu W, Chen H, et al., 2021. A novel human long noncoding rna scdal promotes angiogenesis through snf5-mediated gdf6 expression. *Adv Sci (Weinh)*, 8(18):e2004629. https://doi.org/10.1002/advs.202004629

Xiao F, Wang C, Wang C, et al., 2018. Bmper enhances bone formation by promoting the osteogenesis-angiogenesis coupling process in mesenchymal stem cells. *Cell Physiol Biochem*, 45(5):1927-1939. https://doi.org/10.1159/000487969

Xiao Y, Li C, Wang H, et al., 2020. Linc00265 targets mir-382-5p to regulate sat1, vav3 and angiogenesis in osteosarcoma. *Aging (Albany NY)*, 12(20):20212-20225. https://doi.org/10.18632/aging.103762

Xu M, Chen X, Lin K, et al., 2018. The long noncoding rna snhg1 regulates colorectal cancer cell growth through interactions with ezh2 and mir-154-5p. *Mol Cancer*, 17(1):141. https://doi.org/10.1186/s12943-018-0894-x

Yan Y, Fan Q, Wang L, et al., 2017. Lncrna snhg1, a non-degradable sponge for mir-338, promotes expression of proto-oncogene cst3 in primary esophageal cancer cells. *Oncotarget*, 8(22):35750-35760. https://doi.org/10.18632/oncotarget.16189

Yang C, Gu Z, Ding R, et al., 2021. Long non-coding rna meg3 silencing and microrna-214 restoration elevate osteoprotegerin expression to ameliorate osteoporosis by limiting txnip. *J Cell Mol Med*, 25(4):2025-2039. https://doi.org/10.1111/jcmm.16096

Yang X, Yang J, Lei P, et al., 2019. Lncrna malat1 shuttled by bone marrow-derived mesenchymal stem cells-secreted exosomes alleviates osteoporosis through mediating microrna-34c/satb2 axis. *Aging (Albany NY)*, 11(20):8777-8791. https://doi.org/10.18632/aging.102264

Yu G, Li S, Liu P, et al., 2020. Lncrna tug1 functions as a cerna for mir-6321 to promote endothelial progenitor cell migration and differentiation. *Exp Cell Res*, 388(1):111839. https://doi.org/10.1016/j.yexcr.2020.111839

Yu X, Rong PZ, Song MS, et al., 2021. Lncrna snhg1 induced by sp1 regulates bone remodeling and angiogenesis via sponging mir-181c-5p and modulating sfrp1/wnt signaling pathway. *Mol Med*, 27(1):141. https://doi.org/10.1186/s10020-021-00392-2

Yu X, Song MS, Rong PZ, et al., 2022. Lncrna snhg1 modulates adipogenic differentiation of bmscs by promoting dnmt1 mediated opg hypermethylation via interacting with ptbp1. *J Cell Mol Med*, 26(1):60-74. https://doi.org/10.1111/jcmm.16982

Zang LY, Yang XL, Li WJ, et al., 2022. Long noncoding rna metastasis-associated lung adenocarcinoma transcript 1 promotes the osteoblast differentiation of human bone marrow-derived mesenchymal stem cells by targeting the microrna-96/osterix axis. *J Craniofac Surg*, 33(3):956-961. https://doi.org/10.1097/SCS.0000000000008092

Zhang J, Tao Z, Wang Y, 2018a. Long non-coding rna dancr regulates the proliferation and osteogenic differentiation of human bone-derived marrow mesenchymal stem cells via the p38 mapk pathway. *Int J Mol Med*, 41(1):213-219. https://doi.org/10.3892/ijmm.2017.3215

Zhang N, Meng X, Mei L, et al., 2018b. The long non-coding rna snhg1 attenuates cell apoptosis by regulating mir-195 and bcl2-like protein 2 in human cardiomyocytes. *Cell Physiol Biochem*, 50(3):1029-1040. https://doi.org/10.1159/000494514

Zhang N, Hu X, He S, et al., 2019a. Lncrna msc-as1 promotes osteogenic differentiation and alleviates osteoporosis through sponging microrna-140-5p to upregulate bmp2. *Biochem Biophys Res Commun*, 519(4):790-796. https://doi.org/10.1016/j.bbrc.2019.09.058

Zhang SZ, Lu ZF, Xu YJ, et al., 2018c. Steel participates in fracture healing through upregulating angiogenesis-related genes by recruiting parp 1. *Eur Rev Med Pharmacol Sci*, 22(12):3669-3675. https://doi.org/10.26355/eurrev_201806_15245

Zhang X, Huang P, Jiang G, et al., 2021. A novel magnesium ion-incorporating dual-crosslinked hydrogel to improve bone scaffold-mediated osteogenesis and angiogenesis. *Mater Sci Eng C Mater Biol Appl*, 121:111868. https://doi.org/10.1016/j.msec.2021.111868

Zhang Y, Wu J, Jing H, et al., 2019b. Long noncoding rna meg3 inhibits breast cancer growth via upregulating endoplasmic reticulum stress and activating nf-kappab and p53. *J Cell Biochem*, 120(4):6789-6797. https://doi.org/10.1002/jcb.27982

Zhang Y, Cao X, Li P, et al., 2020. Lncrna nkila integrates rxfp1/akt and nf-kappab signalling to regulate osteogenesis of mesenchymal stem cells. *J Cell Mol Med*, 24(1):521-529. https://doi.org/10.1111/jcmm.14759

Zhang ZC, Tang C, Dong Y, et al., 2017. Targeting the long noncoding rna malat1 blocks the pro-angiogenic effects of osteosarcoma and suppresses tumour growth. *Int J Biol Sci*, 13(11):1398-1408. https://doi.org/10.7150/ijbs.22249

Zheng J, Guo H, Qin Y, et al., 2020. Snhg5/mir-582-5p/runx3 feedback loop regulates osteogenic differentiation and apoptosis of bone marrow mesenchymal stem cells. *J Cell Physiol*, https://doi.org/10.1002/jcp.29527

Zheng S, Wang YB, Yang YL, et al., 2019. Lncrna malat1 inhibits osteogenic differentiation of mesenchymal stem cells in

Unedited

osteoporosis rats through mapk signaling pathway. *Eur Rev Med Pharmacol Sci*, 23(11):4609-4617. https://doi.org/10.26355/eurrev_201906_18038

Zheng S, Zhou C, Yang H, et al., 2022. Melatonin accelerates osteoporotic bone defect repair by promoting osteogenesis-angiogenesis coupling. *Front Endocrinol (Lausanne)*, 13:826660. https://doi.org/10.3389/fendo.2022.826660

Zhou L, Wan Y, Cheng Q, et al., 2020. The expression and diagnostic value of lncrna h19 in the blood of patients with osteoarthritis. *Iran J Public Health*, 49(8):1494-1501. https://doi.org/10.18502/ijph.v49i8.3893

Zhou Y, Zhang X, Klibanski A, 2012. Meg3 noncoding rna: A tumor suppressor. *J Mol Endocrinol*, 48(3):R45-53. https://doi.org/10.1530/JME-12-0008

Zhu J, Wang Y, Yu W, et al., 2019. Long noncoding rna: Function and mechanism on differentiation of mesenchymal stem cells and embryonic stem cells. *Curr Stem Cell Res Ther*, 14(3):259-267. https://doi.org/10.2174/1574888X14666181127145809

Zhuang W, Ge X, Yang S, et al., 2015. Upregulation of lncrna meg3 promotes osteogenic differentiation of mesenchymal stem cells from multiple myeloma patients by targeting bmp4 transcription. *Stem Cells*, 33(6):1985-1997. https://doi.org/10.1002/stem.1989

Unedited