Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology) in press
www.jzus.zju.edu.cn; www.springer.com/journal/11585
E-mail: jzus_b@zju.edu.cn

Review

https://doi.org/10.1631/jzus.B2300743

Prefrontal cortical circuits in social behaviors: an overview

Wei CAO[1,3,4], Huiyi LI[1,3], Jianhong LUO[2,5,6]✉

[1] *Institute of Brain Science and Department of Physiology, School of Basic Medical Sciences, Hangzhou Normal University, Hangzhou 311121, China*
[2] *Affiliated Mental Health Center & Hangzhou Seventh People's Hospital, School of Brain Science and Brain Medicine, Zhejiang University School of Medicine, Hangzhou 310013, China*
[3] *Key Laboratory of Aging and Cancer Biology of Zhejiang Province, School of Basic Medical Sciences, Hangzhou Normal University, Hangzhou 311121, China*
[4] *Zhejiang Philosophy and Social Science Laboratory for Research in Early Development and Childcare, Hangzhou Normal University, Hangzhou 311121, China*
[5] *Key Laboratory of Novel Targets and Drug Study for Neural Repair of Zhejiang Province, School of Medicine, Hangzhou City University, Hangzhou 310015, China*
[6] *NHC and CAMS Key Laboratory of Medical Neurobiology, Zhejiang University, Hangzhou 310058, China*

Abstract: Social behaviors are fundamental and intricate functions in both humans and animals, governed by the interplay of social cognition and emotions. A noteworthy feature of several neuropsychiatric disorders, including autism spectrum disorders (ASD) and schizophrenia (SCZ), is a pronounced deficit in social functioning. Despite a burgeoning body of research on social behaviors, the precise neural circuit mechanisms underpinning these phenomena remain to be elucidated. In this paper, we review the pivotal role of the prefrontal cortex (PFC) in modulating social behaviors, as well as its functional alteration in social disorders in ASD or SCZ. We posit that PFC dysfunction may represent a critical hub in the pathogenesis of psychiatric disorders characterized by shared social deficits. Furthermore, we delve into the intricate connectivity of the mPFC with other cortical areas and subcortical brain regions in rodents, which exerts a profound influence on social behaviors. Notably, a substantial body of evidence underscores the role of N-methyl-D-aspartate (NMDA) receptors and the proper functioning of parvalbumin-positive interneurons within the mPFC for social regulation. Our overarching goal is to furnish a comprehensive understanding of these intricate circuits, and thereby contribute to the enhancement of both research endeavors and clinical practices concerning social behavior deficits.

Key words: Prefrontal cortex; Social behaviors; Autism spectrum disorder; Schizophrenia; Parvalbumin-positive interneurons; NMDA receptors

unedited

1 Introduction

Generally, social behaviors can be defined as any form of communication and interaction between two conspecifics. These behaviors are indispensable and complex in many species, essential for survival and reproduction (Ebstein et al., 2010; Mcgraw and Young, 2010). However, when exhibited at inappropriate times, places or intensities, social behaviors can have detrimental effects on both individuals and the social group as a whole. Impairments in social functioning are prominent features of several neuropsychiatric disorders,

✉ Jianhong LUO, luojianhong@zju.edu.cn
Jianhong LUO, https://orcid.org/0000-0001-7832-496X
Wei CAO, https://orcid.org/0000-0001-6196-5680

Received Oct. 12, 2023; Revision accepted Jan. 2, 2024;
Crosschecked xxx. xx, 20xx; Published online xxx. xx, 20xx

© Zhejiang University Press 2024

including autism spectrum disorder (ASD) and schizophrenia (SCZ) (Chen and Hong, 2018).

The prefrontal cortex (PFC) and its extensive bidirectional connections, forming a top-down modulatory system for social behaviors, are components of the intricate and vast neural networks related to social behaviors (Anastasiades and Carter, 2021; Klune et al., 2021). Throughout the evolution of mammals, the prefrontal region has expanded in size relative to the rest of the cortex, reaching its largest volume in the human brain. Present-day neuroscientists have increasingly turned to research involving mice to gain insights into the underlying mechanisms taking place in the prefrontal cortex for social behaviors. Studies in both humans and animals have underscored the central role of the frontal brain regions in cognition (Carlen, 2017).

Despite the rapid advancements in molecular, cellular and genetic methodologies, alongside the implementation of cutting-edge imaging technologies, the precise neural circuit mechanisms responsible for social behavior remain elusive. In this paper, we aim to provide a concise review of the role of PFC in mediating a wide range of social behaviors in ASD/SCZ patients and animal models.

2 The prefrontal cortex and social regulation in humans

2.1 Structure and function of the prefrontal cortex

The prefrontal cortex plays an indispensable role in higher brain function, including cognition, motivation, reward, and emotion. It also governs goal-directed behaviors, social behavior and moral judgement (Forbes and Grafman, 2010; Buschman and Miller, 2014; Hanganu-Opatz et al., 2023). In humans, the PFC can be divided into distinct regions, including dorsolateral (dlPFC), ventrolateral (vlPFC), dorsomedial (dmPFC), ventromedial (vmPFC), and orbitofrontal cortex (OFC) (Forbes and Grafman, 2010). While the medial regions of the PFC (mPFC) are specifically associated with social behavior, the lateral regions, including the dlPFC and vlPFC, may become active during social tasks while they are generally regarded as ‘domain general’ (Amodio and Frith, 2006; Mitchell et al., 2006). Other brain regions commonly recruited by social behaviors in humans include the anterior cingulate cortex (ACC), nucleus accumbens (NAc), amygdala, hippocampus (HPC), superior temporal sulcus, and temporal parietal junction (Gangopadhyay et al., 2021; Chen et al., 2023). Regions of the PFC exhibit dense interconnections. Major glutamatergic projections arise from the PFC to the ACC, thalamus, ventral tegmental area (VTA), HPC and NAc. Additionally, glutamatergic neurons originate from the HPC and innervate into the hypothalamus (HT), VTA, NAc, and PFC, and from the amygdala to HT, ACC and NAc (Sarawagi et al., 2021). From birth to early adulthood, PFC cells and circuits undergo physiological changes, altering their strength of connectivity with distant brain regions. The extended maturation of the PFC likely enables the emergence of complex behaviors but also renders them more susceptible to disruptions (Kolb et al., 2012; Klune, et al., 2021).

Studies in humans have highlighted the central role of the frontal brain regions in cognition. Higher-order cognitive abilities encompass attention, salience detection, working memory, strategy shifting, and inhibitory control, all of which facilitate adaptation to varying conditions (Diamond, 2013; Buschman and Miller, 2014). These abilities necessitate the coordination of cellular ensembles that balance the stability and flexibility of neural representations. During baseline activity, prefrontal neurons lack the functional coordination of activity; however, when task-related demands arise, the temporal coordination of activity dynamically binds prefrontal neurons into functional units through oscillations and polychrony (Hanganu-Opatz, et al., 2023). This synchronization influences communication between neuronal groups and ensures that presynaptic activation patterns reach postsynaptic neurons in a temporally coordinated manner. Oscillatory phase locking is primarily driven by bottom-up-directed gamma band activity (30-90 Hz) (Fries, 2015). In each oscillatory cycle, representing a transition between an averaged depolarized and hyperpolarized state, neurons have a temporal window to synchronize their firing, establishing connections (Buzsaki and Draguhn, 2004). Oscillations are hypothesized to create a temporal scaffold for intra- and interbrain area communication (Buzsaki and Llinas, 2017). The gamma rhythm primarily arises from the synchronized fast inhibition of excitatory neurons by parvalbumin (PV)-expressing interneurons, demonstrating its role as a resonant property in local networks (Cardin et al.,

Unedited

2009; Sohal et al., 2009). It has been suggested that local gamma oscillations mirror the transient activation of PFC neuronal assemblies (Fujisawa et al., 2008). The gamma rhythm usually occurs along with the theta rhythm (4-12 Hz). The synchronization of theta rhythm between PFC and hippocampus can influence various cognitive states, including memory integration and sequential working memory (Backus et al., 2016; Su et al., 2023). To this end, research has illuminated the vital role of oscillatory synchrony, particularly gamma and theta rhythms, in facilitating communication within the prefrontal cortex and between various brain areas, which underscores their significance in orchestrating complex behaviors, including social interactions.

2.2 Cognitive and emotional processing in social regulation

Social behavior deficits represent a fundamental dimension of many psychiatric disorders, such as the neurodevelopmental disorders ASD and SCZ. In social decision-making contexts, individuals frequently need to recognize other individuals, infer their intentions and emotions, and weigh the values of both social and non-social outcomes before choosing an action. To a significant extent, these aspects of social information processing rely on the prefrontal cortex, specifically the medial prefrontal cortex (Kietzman and Gourley, 2023). Consequently, social cognition and emotion play critical roles in social regulation.

Social cognition encompasses a wide array of mental operations used for identifying and interpreting social signals, as well as the utilization of these signals to flexibly guide appropriate social behaviors in changing environments (Millan and Bales, 2013). Social cognition has three major facets: social motivation, self and other knowledge, and group dynamics. Research using both humans and translational animal models has demonstrated that these aspects of social cognition relate to psychiatric disorders. Social motivation can be described as biasing the individual to preferentially orient to the social world (social orienting), to seek and take pleasure in social interactions (social reward), and to strive to foster and maintain social bonds (social maintaining). The social motivation theory of autism suggests that a lack of early social interest may contribute to subsequent social cognitive deficits that manifest later in development (Chevallier et al., 2012). Thus, social motivation can be seen as a developmental and evolutionary foundation for other social behaviors. Self and other knowledge forms the core of social cognition and encompasses concepts explored in basic processes like facial recognition, action perception and empathy, as well as more sophisticated social behaviors such as cooperation and intergroup interaction (Catmur et al., 2016). Living in groups is common in mammalian societies, and dominance hierarchy is a fundamental organizing mechanism based on repeated social competitions. The result of social competition is influenced by personality traits under the regulation of high cortical functions (Zhou et al., 2018). Among the high cortical regions, the mPFC has been specifically associated with social dominance (Wang et al., 2011; Zhang et al., 2022; Fan et al., 2023). Hence, social cognition encompasses a wide range of mental processes crucial for our understanding of social interactions. These facets of social cognition have not only been linked to psychiatric disorders in humans and animal models but also offer insights into the fundamental building blocks of social behavior.

Emotion regulation involves processes that affect both the experience and expression of emotions; it is essential for successful social interactions and functioning (Mcrae and Gross, 2020). Effective emotion regulation is associated with an increased capacity to engage in socially appropriate emotions and behaviors, contributing to adaptive social interactions and social competence (Eisenberg et al., 1995; Eisenberg et al., 2000). Conversely, disruptions in effective emotion regulation strategies may result from negative social experiences, leading to difficulties in social behaviors and negative social encounters (Eisenberg et al., 1999; Geckeler et al., 2022). These intricate processes of emotion regulation play pivotal roles in shaping the quality of social interactions and overall social functioning.

Taken together, the prefrontal cortex, with its intricate network of regions and dynamic neural activity patterns, serves as the cornerstone of higher brain functions, influencing cognition, emotion, social behavior, and more. At the same time, addressing social behavior deficits in psychiatric disorders like ASD and SCZ necessitates a comprehensive exploration of the interconnected dimensions of social cognition and emotion regulation. These conditions often involve complex processes of social information processing that heavily rely

Unedited

on the prefrontal cortex, particularly the medial prefrontal cortex.

3 Prefrontal dysfunction in autism and schizophrenia

3.1 Brain imaging evidence of impaired PFC networks in ASD

Numerous researchers have underscored the pivotal role of the PFC in the manifestation of various ASD symptoms. For instance, functional magnetic resonance imaging (fMRI) studies have revealed that dysfunctional networks within the frontal lobes may affect not only social cognition but also repetitive behavior, executive function and verbal communication—functions primarily controlled by the frontal lobe (Baron-Cohen et al., 1999; Carper and Courchesne, 2000; Ohnishi et al., 2000; Luna et al., 2002; Herbert et al., 2003; Carper and Courchesne, 2005; Chandana et al., 2005; Rafiee et al., 2022). Furthermore, reduced serotonergic synthesis in the frontal lobes, along with concomitant abnormalities of whole-brain serotonin synthesis, has been observed in individuals with ASD (Chugani et al., 1999; Chandana, et al., 2005). Serotonin, similar to the brain-derived neurotrophic factor (BDNF), modulates axonal arborization, and dysfunction in serotonin synthesis has been linked to frontal lobe connectivity anomalies (Kumar et al., 2010).

Disorders of neural connectivity, which explain a substantial portion of ASD variance, enable the integration of cognitive theories (e.g., central coherence) with a comprehensive understanding of ASD behaviors at the neuroanatomical, neurophysiological and neuropsychological levels (Hughes, 2007,2008). Impaired integration in ASD may arise from functional dysconnectivity among brain systems. Existing data suggest both cortico-subcortical and inter- and intra-hemispheric hyper- and hypoconnectivity (Vissers et al., 2012; Uddin et al., 2013; Kleinhans et al., 2016; Martinez et al., 2020). The role of frontal streams, as presented in the following section, has been validated through postmortem studies of individuals with ASD. An important finding was the observation of brain overgrowth postpartum early in life, particularly of the PFC, which correlated with significantly larger head circumferences in children under three years of age (Courchesne et al., 2003; Sacco et al., 2015). Subcortical brain regions involved in sensory processing that project to the PFC, including the cerebellum and the thalamus, also exhibit anatomical differences compared to neurotypical individuals (Hazlett et al., 2005; Courchesne et al., 2011a; Courchesne et al., 2011b). The underconnectivity observed between frontal-thalamic areas has been associated with the severity of symptoms in individuals with ASD (Martinez-Sanchis, 2014; Rane et al., 2015). Moreover, there are strong connections between the PFC and the cerebellum, and abnormalities in both regions have been linked to symptom severity (Carper and Courchesne, 2000; Kumar, et al., 2010).

Unedited

3.2 Brain imaging as evidence for impaired PFC networks in SCZ

Numerous reports have underscored the significant similarities between the cognitive deficits observed in patients with frontal lesions and those encountered in schizophrenia (Muller et al., 2002). Furthermore, functional imaging studies have shed light on altered PFC activation during cognitive testing in schizophrenia patients, supporting the hypothesis that disruptions in PFC activity cause deficits in working memory and other cognitive functions (Weinberger and Berman, 1996; Karlsgodt et al., 2009). In fact, SCZ patients exhibit reduced blood flow in the PFC during task performance and diminished functional connectivity between the PFC and other brain regions when engaged in cognitive tasks (Goldman-Rakic and Selemon, 1997; Fornito et al., 2009).

A crucial source of excitatory input to the PFC originates from the thalamus, including the mediodorsal nucleus (MD). A reduction in correlated activity between the thalamus and the PFC during cognitive testing could be a potential factor contributing to cognitive deficits (Mitelman et al., 2005; Woodward et al., 2012; Giraldo-Chica et al., 2018). Similarly, assessments of resting-state functional connectivity have indicated reduced thalamoprefrontal connectivity in various stages of schizophrenia, including in adolescents with early-onset schizophrenia, young adolescents at clinical high risk, and adults in early stages of the disorder (Anticevic et al., 2015; Cho et al., 2016; Woodward and Heckers, 2016; Huang et al., 2021; Zhang et al., 2021).

Notably, reduction in thalamoprefrontal connectivity among high-risk individuals becomes the most pronounced in those who are diagnosed with the disorder later (Anticevic, et al., 2015). These findings intriguingly suggest that thalamo-PFC dysconnectivity during adolescence might contribute to the developmental etiology of schizophrenia (Anticevic, et al., 2015; Woodward and Heckers, 2016).

Schizophrenia is associated with both aberrant PFC development during embryonic stages and disrupted PFC maturation in later adolescence driven by environmental stressors. Due to this unique aspect, SCZ can be characterized by two critical susceptibility periods (Selemon and Zecevic, 2015). The disruption of cortical synaptic pruning has been proposed as a key factor in the etiology of SCZ, given that its symptoms typically manifest in late adolescence, which coincides with the developmental period when PFC connectivity matures (Feinberg, 1990; Lewis, 1997). Gray matter loss, reflecting cortical maturation, tends to occur later in higher-order association cortices such as the prefrontal and temporal cortices compared to lower-order somatosensory and visual cortices, indicating a delayed maturation of these cortical regions (Gogtay et al., 2004). Cortical gray matter loss is more pronounced in individuals at clinical high risk for schizophrenia who develop the disorder later, in contrast to others (Cannon et al., 2015; Chung et al., 2017). These observations suggest that alterations in prefrontal maturation may play a role in the etiology of the disorder.

Findings from postmortem studies consistently reveal changes in elements of GABAergic neurotransmission within the PFC of individuals with schizophrenia. The effectiveness of GABAergic neurotransmission correlates with the availability of GABA in the synapse. In schizophrenia, both mRNA and protein levels of GAD67, the principal enzyme responsible for cortical GABA synthesis, are diminished in the PFC (Vawter et al., 2002; Woo et al., 2008; Curley et al., 2011; Kimoto et al., 2014). This aligns with the notion that the availability of presynaptic GABA is reduced in this illness. Additionally, the number of PV^+ (Beasley and Reynolds, 1997; Beasley et al., 2002; Hashimoto et al., 2003), CCK^+ (Hashimoto et al., 2008; Fung et al., 2010) and SST^+ (Hashimoto, et al., 2008; Morris et al., 2008; Fung, et al., 2010) interneurons is decreased in the PFC of individuals with schizophrenia. Interestingly, GABA concentration has been found to predict the gamma oscillatory frequency (Edden et al., 2009). There is substantial evidence supporting N-methyl-D-aspartate receptor (NMDAR) hypofunction in SCZ, as treatment with NMDAR antagonists, such as MK-801 or phencyclidine, could induce SCZ-like symptoms in human subjects. Consequently, it has been postulated that the NMDAR hypofunction in SCZ primarily occurs in cortical GABAergic interneurons (Nakazawa et al., 2017).

In brief, mounting evidence regarding PFC impairment in both ASD and SCZ patients highlights the critical role of this brain region in the manifestation of symptoms in these disorders. Brain imaging studies have unveiled disrupted PFC and PFC network functionality, enhancing our understanding of the cognitive deficits and behavioral challenges seen in these conditions. Additionally, insights from studies examining neural connectivity and neurotransmission further emphasize the significance of the PFC in the pathophysiology of ASD and SCZ. Understanding these neural underpinnings is crucial for developing targeted interventions and treatments to improve the lives of individuals affected by these disorders.

4 Prefrontal mechanisms of social deficits in rodents

4.1 Measuring social behaviors in rodents

A multitude of studies have demonstrated that rodents exhibit behaviors closely aligned with ethological aspects of social processing demands. These behaviors encompass sociability, social memory/recognition and dominance, offering conceptual parallelism with human social categories (Moy et al., 2004; Bariselli et al., 2018; Zhou, et al., 2018). Given their inherent social nature, mice serve as invaluable research models, facilitating the exploration of aberrant social behaviors associated with ASD and SCZ.

In order to assess sociability in mice, behavioral paradigms like social preference tests have often been employed. These tests assess the amount of time spent in proximity to an unfamiliar mouse compared to an empty side within a three-chamber apparatus. Typically, the novel social stimulus is confined to a compartment that allows sniffing and interaction but prevents physical contact (Moy, et al., 2004). Alternatively, sociability

Unedited

can be evaluated by measuring the time spent engaged in unconstrained interactions.

In order to quantify social recognition in mice, researchers can measure the time spent with a novel versus a familiar conspecific in a three-chamber apparatus (Moy, et al., 2004). Typically, wild-type mice exhibit a preference for social novelty by spending more time with the novel conspecific. Other paradigms for measuring social recognition take advantage of mice's natural tendency to habituate to a familiar conspecific while showing greater interest in exploring a novel mouse. A reduction in sniffing time during repeated trials indicates recognition of the familiar mouse. After repeated presentations, a novel mouse is introduced, and an increased investigation of the novel mouse reflects a preference for social novelty (Bariselli, et al., 2018).

Dominance hierarchy serves as a fundamental organizing mechanism in most animal societies. Once established, the hierarchical rank remains relatively stable and reduces intense conflicts among group members. To assess the dominance tendencies of mice, the tube test was developed, in which one mouse, termed as 'loser', is compelled to back out of a narrow tube by another mouse, referred to as 'winner' (Wang, et al., 2011; Zhang, et al., 2022). A simpler method is to observe the behaviors of animals within their home cages or to monitor aggressive interactions, which typically occur when a group is placed into a new cage (Shemesh et al., 2013).

Play behavior can be observed in various animals, including rodents. In rats, most play behavior is social, involving play fighting, where animals compete for an advantage (Palagi et al., 2016; Pellis et al., 2023). Play fighting activity peaks during the juvenile period, and juveniles are highly motivated to engage peers in this behavior (Varlinskaya et al., 1999). Consequently, social play interactions are a major source of social experience during this developmental stage. It is important to note, however, that play fighting does not specifically train particular motor actions; instead, it enhances a skill set applicable in various social and non-social contexts. Depriving juvenile rats of typical peer-to-peer play experiences results in adults displaying socio-cognitive deficiencies, which correlate with physiological and anatomical changes in neurons, notably in the prefrontal cortex, especially the mPFC (Pellis, et al., 2023).

Over the last decade, there has been a surge of investigations focused on how mPFC circuits and pathways modulate social behaviors. While the hypothalamus and other subcortical circuits play a central role in generating social behaviors, the mPFC is uniquely positioned to integrate pertinent information such as social rank, memories, and context to modulate these behaviors. In the subsequent sections, we review the available evidence indicating that activity within the rodent mPFC can lead to alterations in social behaviors.

Unedited

4.2 Involvement of the medial prefrontal cortex in social deficits in rodents

The medial prefrontal cortex of rodents consists of the anterior cingulate cortex, the prelimbic cortex (PL), and the infralimbic cortex (IL), each of them with unique connectivity and functional properties. Multiple lines of evidence indicate the significance of neural activity originating from the mPFC in shaping social behavior in rodents. Although the medial aspect of the secondary motor cortex (M2) is sometimes considered to be a part of the rodent mPFC, we will not include it in this review (Barthas and Kwan, 2017).

4.2.1 Altered synaptic transmission in the mPFCs

Studies involving rodents have shown that dysfunction in the prefrontal cortex (PFC) reduces social interest (Murray et al., 2015), impairs social memory (Rudebeck et al., 2007), hinders the processing of social hierarchy information, and lowers social rank (Wang et al., 2014). Research has also delved into synaptic transmission in mouse models with genetic modification derived from risk genes for ASD and SCZ, and revealed general synaptic dysfunctions in the cortex, and in some cases, specifically in the PFC. These studies have involved the abnormal expression of various molecules, including MeCP2 (Sceniak et al., 2016), b2 nicotinic receptor (Avale et al., 2011), insulin receptor substrate protein (IPSp53) (Chung et al., 2015), Shank3 (Duffney et al., 2015; Lee et al., 2015), and Neuroligin 3 (NL3) (Cao et al., 2018), which have all been linked to altered synaptic transmission in the mPFC and abnormal social behavior.

Excitation/inhibition imbalance and synaptic dysfunction have been identified as common pathophysiological features in various mouse models of ASD and SCZ. For instance, the conditional knockout of Neuroligin

2 (NL2) in the adult mPFC induced significant reductions in synaptic inhibition and caused parallel impairments in anxiety, fear memory and social interaction behaviors (Liang et al., 2015). Also, NMDAR hypofunction and dysfunction of PV^+ interneurons have been observed in NL3-deficient mice, and the micro-infusion of NMDAR co-agonist D-cycloserine (DCS) into the mPFC rescued the social deficits in these mice (Cao et al., 2022). Similar findings have been reported in other mouse models. For example, a 16p11.2 deletion mouse model exhibited NMDAR hypofunction in PFC pyramidal neurons and cognitive and social impairments, which could be rescued by the chemogenetic activation of PFC pyramidal neurons (Wang et al., 2018). Moreover, treatment with NMDAR antagonists such as MK-801 or phencyclidine could induce schizophrenia-like symptoms in rodents. These symptoms included hyperlocomotion as an index of positive symptoms, a social behavioral deficit in a social interaction test, and enhanced immobility in a forced swimming test as indices of negative symptoms (Kehrer et al., 2008; Nakazawa, et al., 2017; Kruse and Bustillo, 2022). This strongly supports the NMDA receptor hypofunction hypothesis of schizophrenia. Overall, the above findings suggest the importance of normal synaptic function in the mPFC, particularly the role of NMDAR, and provide valuable insights into potential therapeutic approaches targeting specific molecular and synaptic mechanisms, such as NMDARs, for the development of more effective treatments for ASD and SCZ.

Excitation/inhibition imbalance within the mouse mPFC has been linked to profound social deficits (Yizhar et al., 2011). Studies of mouse models lacking the *CNTNAP2* gene, implicated in autism, have demonstrated that modulating excitation/inhibition in the mPFC through neuron type-selective, real-time, reversible optogenetic manipulations can acutely rescue deficits in social behavior and hyperactivity (Selimbeyoglu et al., 2017). Moreover, a 16p11.2 duplication mouse model exhibited deficient GABAergic synaptic transmission and elevated excitability in PFC pyramidal neurons, as well as social and cognitive deficits. These deficits could be restored by the expression of Npas4 in PFC, a key regulator of GABA synapses (Rein et al., 2021). In line with this finding, decreased cortical inhibition and/or PV^+ interneuron number have been commonly observed in mouse models with social impairments (Han et al., 2012; Filice et al., 2016; Cao, et al., 2018; Cao, et al., 2022). These results indicate that maintaining normal synaptic transmission and proper inhibition within the mPFC neuronal network is essential for normal social behaviors.

Unedited

4.2.2 Neuronal oscillations in the mPFC during social interactions

The neural activity of principal neurons of the mPFC is crucial for the social behavior of rodents. Studies have shown that certain mPFC principal neurons increase their activity during social interactions in mice (Lee et al., 2016; Brumback et al., 2018). Calcium activity recordings in mPFC principal neurons have identified distinct and dynamic ON and OFF neural ensembles that encode social exploration, and linked dysfunctions in the activity of these ensembles to abnormal social exploration (Liang et al., 2018). Furthermore, the transient excitation of excitatory neurons in the mPFC through optogenetic manipulation led to reduced sociability (Yizhar, et al., 2011) but improved social competition (Zhang, et al., 2022).

In addition, neural oscillations in the low gamma frequency range (30-50 Hz) have been found to be disturbed in ASD and SCZ (Gandal et al., 2012; David et al., 2016). Optogenetic studies have revealed that activating PV^+ interneurons is both necessary and adequate for inducing gamma oscillations in the cortex, thereby influencing cortical information processing (Cardin, et al., 2009; Sohal, et al., 2009). Modulating these oscillations, especially through patterned optogenetic stimulation (40-8 Hz) of mPFC PV^+ interneurons, has been shown to rescue social deficit in an NL3-deficient mouse model of autism (Cao, et al., 2018). Interestingly, the excitation of vasoactive intestinal polypeptide-positive interneurons or the inhibition of PV^+ interneurons induced winning, and vice versa (Zhang, et al., 2022). The optogenetic synchronization of either PV+ interneurons or somatostatin-positive interneurons at a low gamma frequency improved sociability in wild-type mice (Liu et al., 2020). These findings underscore the close correlation between social interaction and elevated gamma rhythms in the prefrontal local field potentials, controlled by PV^+ interneurons. As a result, targeted interventions aimed at restoring neuronal oscillations hold great promise for future therapeutic strategies in neurodevelopmental and neuropsychiatric disorders.

4.2.3 Distinct mPFC circuits are involved in social behaviors

The mPFC receives a variety of long-range inputs and sends diverse outputs to many other brain regions, connected via local circuits involving different excitatory and inhibitory neurons and communicating with each other based on specific wiring rules.

Cell type-specific optogenetic and chemogenetic manipulations, behavioral testing, and electrophysiological recordings have been used to investigate the functional coupling of the mPFC with subcortical systems in social behavior. Studies have shown that projection neurons of the basolateral amygdala (BLA) in the IL are preferentially activated in response to a social cue as compared with BLA-projecting neurons in the PL, and the chemogenetic activation of PL-BLA or the inhibition of IL-BLA circuits impairs social behavior (Huang et al., 2020). Moreover, the activation of BLA input to the mPFC leads to an increase in anxiety-like behavior and a decrease in sociability (Felix-Ortiz et al., 2016). Interestingly, PL to NAc stimulation decreased the preference for a social target (Murugan et al., 2017). The ventral hippocampus (vHPC) has been implicated in social memory, and the inhibition of direct projections from vHPC to mPFC impaired social memory expression (Okuyama et al., 2016; Sun et al., 2020). In addition, the firing rates of VTA dopamine neurons projecting to the mPFC were dramatically decreased in a mouse model of repeated social defeat-induced depression. The activation of VTA projections to the NAc or the inhibition of VTA projections to the mPFC induced susceptibility to social defeat stress (Liu et al., 2018). The mPFC is also involved in processing the affective state of others through non-verbal communication. As visualized by *in vivo* single-cell microendoscopic Ca^{2+} imaging, an increase in the synchronous activity of mPFC somatostatin-positive interneurons, guiding the inhibition of pyramidal neurons, has been associated with affective state discrimination (Scheggia et al., 2020). Interestingly, prior social experience incentivizes later instrumental choices such as food, and the social incentivization of future choices requires the PL (Kietzman et al., 2022).

Moreover, serotonergic neurons in the dorsal raphe nucleus (DRN) play a role in social behavior modulation through their interaction with the mPFC. The manipulation of synaptic inputs from the mPFC to the DRN influences social avoidance behaviors (Michelsen et al., 2008; Challis et al., 2013; Challis and Berton, 2015). The oxytocinergic system, originating from the hypothalamus and acting through SST^+ interneurons expressing oxytocin receptors (OXTRs) in the mPFC, also has an important function in regulating sociosexual behaviors in female mice but anxiety-related behaviors in male mice (Donaldson and Young, 2008; Nakajima et al., 2014; Li et al., 2016). Furthermore, OXTRs are also expressed on glutamatergic neurons in the mPFC, and the optogenetic stimulation of axons in the BLA arising from OXTR-expressing neurons in the PFC eliminated the ability to distinguish novel conspecifics from familiar ones (Tan et al., 2019). Thus, this section provides a comprehensive overview of the intricate neural circuits and systems that regulate social behaviors, emphasizing the central role of the mPFC and its interactions with various brain regions and neurotransmitter systems. The mPFC receives inputs from the vHPC, BLA and VTA, and it sends outputs to the NAc, BLA, VTA, and DRN, all of which are implicated in social memory, sociability, social defeat, or anxiety. The modulation of PFC circuits by monoaminergic systems profoundly affects PFC-mediated social behaviors.

In conclusion, the rodent studies summarized in this section strongly support the notion that PFC neurons and their interconnected neural circuits play a central role in the regulation of social behaviors. Within the microcircuits of the mPFC, NMDA receptors and PV^+ interneurons have merged as critical components that potentially serve as pivotal elements in the regulation of social behaviors.

5 Perspectives

The PFC remains a central focus of inquiry in the field of neuroscience, with the potential to unveil deep insights into the complexities of social behavior. Deciphering the pivotal role of the PFC in governing social behaviors not only enriches our comprehension of fundamental brain mechanisms but also carries profound implications for addressing social deficits witnessed in psychiatric disorders such as ASD and SCZ.

Future research endeavors are predicted to illuminate the dynamic interplay within the PFC and its impact

Unedited

on our social interactions. There is a necessity for continued exploration into the neural circuitry of the PFC, with the objective of unraveling how this region choreographs a multitude of processes integral to social cognition and emotion regulation. Furthermore, it is essential to approach the neural mechanisms of social activities by considering social networks as a cohesive entity, including the PFC. In rodents, the unique input and output from diverse subregions, layers and cell types within the PFC govern various facets of social behavior. For example, neurons in the layer 2/3 IL that express the neuropeptide corticotropin-releasing hormone (CRH) respond to social interaction with familiar over novel mice and release CRH into the rostral lateral septum (rLS) to suppress social interactions with familiar mice through LS disinhibition (De Leon Reyes et al., 2023). The mPFC cells that project to the lateral hypothalamus promote dominance behavior during reward competition (Padilla-Coreano et al., 2022). Decreased PV^+ interneuron excitability in the mPFC is a causative factor in social novelty deficits (Cao, et al., 2018). Hence, forthcoming research should focus on meticulously categorizing PFC cells according to their precise spatial localization, physical connections and protein markers. Subsequently, examining the regulatory impact of these smaller cell groupings on social behavior is imperative. A more comprehensive investigation into the intricate microcirculatory networks within the PFC is also warranted; a deeper understanding of PFC circuits will not only expand our knowledge of social behaviors but also provide new treatment strategies for social disorders.

In the preceding discussion, we emphasized the significance of gamma oscillations in the mPFC concerning social behavior, and suggested that targeted interventions aimed at restoring neuronal oscillations might be a promising route of addressing the social disorders of ASD and SCZ. There are three potential approaches for implementing oscillation interventions. The first method involves directly enhancing the weakened oscillations in social behavior through physical interventions. Optogenetics are commonly used in animal studies, but the introduction of exogeneous genes into the brain hinders its application in humans. Currently, electrode stimulations based on information decoding (Li et al., 2023) provide a more physiological and precise approach, potentially adaptable for translation in patients if the concern about invasive surgery can be addressed. Additionally, non-invasive neuromodulations such as transcranial direct current stimulation or transcranial magnetic stimulation can be evaluated in clinical trials, with observations on whether social behavior and gamma oscillation are improved in subjects. The second method comprises pharmacological intervention, utilizing specific drugs that target PV^+ interneurons to regulate their excitability, thereby impacting gamma oscillation. For instance, potential options are drugs targeting Nav1.1 (Ogiwara et al., 2007) and Kv3.1 (Kaczmarek and Zhang, 2017), specifically expressed on PV^+ interneurons. Moreover, it is noteworthy that the excitability of PV^+ interneurons can also be modulated by NMDAR (Cao, et al., 2022), although this modulation is not exclusive to PV interneurons. The third method entails neuron transplantation. Studies have shown that transplanting Nav1.1-enhanced interneurons can increase gamma oscillation and enhance cognition in a mouse model of Alzheimer's disease (Martinez-Losa et al., 2018). As research progresses, this approach could potentially prove effective in human patients. In summary, the exploration of the above varied approaches underscores the potential for targeted interventions in manipulating gamma oscillations, opening up promising avenues for addressing disorders with social disfunction, such as ASD and SCZ. Continued research in these areas may pave the way for novel therapeutic strategies in the realm of neurobiological interventions for social behavior-related conditions.

Uncorrected

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China (81801355 to W.C., U22A20306 and 3192010300 to J.-h. LUO), the Autism Research Special Fund of Zhejiang Foundation for Disabled Persons (2022001 and 2023002 to W.C.), Research and Development Program of Guangdong Province (2019B030335001 to J.-h LUO), and the Non-profit Central Research Institute Fund of Chinese Academy of Medical Sciences (2023-PT310-01 to J.-h. LUO).

Author contributions

Jianhong LUO determined the topic of the article and proposed the program. Wei CAO was responsible for reference searching and manuscript writing, and Jianhong LUO for supervision and manuscript revision. All authors have read and approved the final manuscript.

Compliance with ethics guidelines

Wei CAO and Jianhong LUO declare that they have no conflicts of interest.

This review does not contain any studies with human or animal subjects performed by any of the authors.

References

Amodio DM, Frith CD, 2006. Meeting of minds: The medial frontal cortex and social cognition. *Nat Rev Neurosci*, 7(4):268-277. https://doi.org/10.1038/nrn1884

Anastasiades PG, Carter AG, 2021. Circuit organization of the rodent medial prefrontal cortex. *Trends Neurosci*, 44(7):550-563. https://doi.org/10.1016/j.tins.2021.03.006

Anticevic A, Haut K, Murray JD, et al., 2015. Association of thalamic dysconnectivity and conversion to psychosis in youth and young adults at elevated clinical risk. *JAMA Psychiatry*, 72(9):882-891. https://doi.org/10.1001/jamapsychiatry.2015.0566

Avale ME, Chabout J, Pons S, et al., 2011. Prefrontal nicotinic receptors control novel social interaction between mice. *FASEB J*, 25(7):2145-2155. https://doi.org/10.1096/fj.10-178558

Backus AR, Schoffelen JM, Szebenyi S, et al., 2016. Hippocampal-prefrontal theta oscillations support memory integration. *Curr Biol*, 26(4):450-457. https://doi.org/10.1016/j.cub.2015.12.048

Bariselli S, Hornberg H, Prevost-Solie C, et al., 2018. Role of vta dopamine neurons and neuroligin 3 in sociability traits related to nonfamiliar conspecific interaction. *Nat Commun*, 9(1):3173. https://doi.org/10.1038/s41467-018-05382-3

Baron-Cohen S, Ring HA, Wheelwright S, et al., 1999. Social intelligence in the normal and autistic brain: An fmri study. *Eur J Neurosci*, 11(6):1891-1898. https://doi.org/10.1046/j.1460-9568.1999.00621.x

Barthas F, Kwan AC, 2017. Secondary motor cortex: Where 'sensory' meets 'motor' in the rodent frontal cortex. *Trends Neurosci*, 40(3):181-193. https://doi.org/10.1016/j.tins.2016.11.006

Beasley CL, Reynolds GP, 1997. Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics. *Schizophr Res*, 24(3):349-355. https://doi.org/10.1016/s0920-9964(96)00122-3

Beasley CL, Zhang ZJ, Patten I, et al., 2002. Selective deficits in prefrontal cortical gabaergic neurons in schizophrenia defined by the presence of calcium-binding proteins. *Biol Psychiatry*, 52(7):708-715. https://doi.org/10.1016/s0006-3223(02)01360-4

Brumback AC, Ellwood IT, Kjaerby C, et al., 2018. Identifying specific prefrontal neurons that contribute to autism-associated abnormalities in physiology and social behavior. *Mol Psychiatry*, 23(10):2078-2089. https://doi.org/10.1038/mp.2017.213

Buschman TJ, Miller EK, 2014. Goal-direction and top-down control. *Philos Trans R Soc Lond B Biol Sci*, 369(1655) https://doi.org/10.1098/rstb.2013.0471

Buzsaki G, Draguhn A, 2004. Neuronal oscillations in cortical networks. *Science*, 304(5679):1926-1929. https://doi.org/10.1126/science.1099745

Buzsaki G, Llinas R, 2017. Space and time in the brain. *Science*, 358(6362):482-485. https://doi.org/10.1126/science.aan8869

Cannon TD, Chung Y, He G, et al., 2015. Progressive reduction in cortical thickness as psychosis develops: A multisite longitudinal neuroimaging study of youth at elevated clinical risk. *Biol Psychiatry*, 77(2):147-157. https://doi.org/10.1016/j.biopsych.2014.05.023

Cao W, Lin S, Xia QQ, et al., 2018. Gamma oscillation dysfunction in mpfc leads to social deficits in neuroligin 3 r451c knockin mice. *Neuron*, 97(6):1253-1260.e1257. https://doi.org/10.1016/j.neuron.2018.02.001

Cao W, Li JH, Lin S, et al., 2022. Nmda receptor hypofunction underlies deficits in parvalbumin interneurons and social behavior in neuroligin 3 r451c knockin mice. *Cell Rep*, 41(10):111771. https://doi.org/10.1016/j.celrep.2022.111771

Cardin JA, Carlen M, Meletis K, et al., 2009. Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature*, 459(7247):663-667. https://doi.org/10.1038/nature08002

Carlen M, 2017. What constitutes the prefrontal cortex? *Science*, 358(6362):478-482. https://doi.org/10.1126/science.aan8868

Carper RA, Courchesne E, 2000. Inverse correlation between frontal lobe and cerebellum sizes in children with

Unedited

autism. *Brain*, 123 (Pt 4):836-844. https://doi.org/10.1093/brain/123.4.836
Carper RA, Courchesne E, 2005. Localized enlargement of the frontal cortex in early autism. *Biol Psychiatry*, 57(2):126-133. https://doi.org/10.1016/j.biopsych.2004.11.005
Catmur C, Cross ES, Over H, 2016. Understanding self and others: From origins to disorders. *Philos Trans R Soc Lond B Biol Sci*, 371(1686):20150066. https://doi.org/10.1098/rstb.2015.0066
Challis C, Boulden J, Veerakumar A, et al., 2013. Raphe gabaergic neurons mediate the acquisition of avoidance after social defeat. *J Neurosci*, 33(35):13978-13988, 13988a. https://doi.org/10.1523/JNEUROSCI.2383-13.2013
Challis C, Berton O, 2015. Top-down control of serotonin systems by the prefrontal cortex: A path toward restored socioemotional function in depression. *ACS Chem Neurosci*, 6(7):1040-1054. https://doi.org/10.1021/acschemneuro.5b00007
Chandana SR, Behen ME, Juhasz C, et al., 2005. Significance of abnormalities in developmental trajectory and asymmetry of cortical serotonin synthesis in autism. *Int J Dev Neurosci*, 23(2-3):171-182. https://doi.org/10.1016/j.ijdevneu.2004.08.002
Chen P, Hong W, 2018. Neural circuit mechanisms of social behavior. *Neuron*, 98(1):16-30. https://doi.org/10.1016/j.neuron.2018.02.026
Chen X, Liu J, Luo YJ, et al., 2023. Brain systems underlying fundamental motivations of human social conformity. *Neurosci Bull*, 39(2):328-342. https://doi.org/10.1007/s12264-022-00960-4
Chevallier C, Kohls G, Troiani V, et al., 2012. The social motivation theory of autism. *Trends Cogn Sci*, 16(4):231-239. https://doi.org/10.1016/j.tics.2012.02.007
Cho KI, Shenton ME, Kubicki M, et al., 2016. Altered thalamo-cortical white matter connectivity: Probabilistic tractography study in clinical-high risk for psychosis and first-episode psychosis. *Schizophr Bull*, 42(3):723-731. https://doi.org/10.1093/schbul/sbv169
Chugani DC, Muzik O, Behen M, et al., 1999. Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Ann Neurol*, 45(3):287-295. https://doi.org/10.1002/1531-8249(199903)45:3<287::aid-ana3>3.0.co;2-9
Chung W, Choi SY, Lee E, et al., 2015. Social deficits in irsp53 mutant mice improved by nmdar and mglur5 suppression. *Nat Neurosci*, 18(3):435-443. https://doi.org/10.1038/nn.3927
Chung Y, Haut KM, He G, et al., 2017. Ventricular enlargement and progressive reduction of cortical gray matter are linked in prodromal youth who develop psychosis. *Schizophr Res*, 189:169-174. https://doi.org/10.1016/j.schres.2017.02.014
Courchesne E, Carper R, Akshoomoff N, 2003. Evidence of brain overgrowth in the first year of life in autism. *JAMA*, 290(3):337-344. https://doi.org/10.1001/jama.290.3.337
Courchesne E, Campbell K, Solso S, 2011a. Brain growth across the life span in autism: Age-specific changes in anatomical pathology. *Brain Res*, 1380:138-145. https://doi.org/10.1016/j.brainres.2010.09.101
Courchesne E, Mouton PR, Calhoun ME, et al., 2011b. Neuron number and size in prefrontal cortex of children with autism. *JAMA*, 306(18):2001-2010. https://doi.org/10.1001/jama.2011.1638
Curley AA, Arion D, Volk DW, et al., 2011. Cortical deficits of glutamic acid decarboxylase 67 expression in schizophrenia: Clinical, protein, and cell type-specific features. *Am J Psychiatry*, 168(9):921-929. https://doi.org/10.1176/appi.ajp.2011.11010052
David N, Schneider TR, Peiker I, et al., 2016. Variability of cortical oscillation patterns: A possible endophenotype in autism spectrum disorders? *Neurosci Biobehav Rev*, 71:590-600. https://doi.org/10.1016/j.neubiorev.2016.09.031
De Leon Reyes NS, Sierra Diaz P, Nogueira R, et al., 2023. Corticotropin-releasing hormone signaling from prefrontal cortex to lateral septum suppresses interaction with familiar mice. *Cell*, 186(19):4152-4171 e4131. https://doi.org/10.1016/j.cell.2023.08.010
Diamond A, 2013. Executive functions. *Annu Rev Psychol*, 64:135-168. https://doi.org/10.1146/annurev-psych-113011-143750
Donaldson ZR, Young LJ, 2008. Oxytocin, vasopressin, and the neurogenetics of sociality. *Science*, 322(5903):900-904. https://doi.org/10.1126/science.1158668
Duffney LJ, Zhong P, Wei J, et al., 2015. Autism-like deficits in shank3-deficient mice are rescued by targeting actin regulators. *Cell Rep*, 11(9):1400-1413. https://doi.org/10.1016/j.celrep.2015.04.064
Ebstein RP, Israel S, Chew SH, et al., 2010. Genetics of human social behavior. *Neuron*, 65(6):831-844. https://doi.org/10.1016/j.neuron.2010.02.020
Edden RA, Muthukumaraswamy SD, Freeman TC, et al., 2009. Orientation discrimination performance is predicted by gaba concentration and gamma oscillation frequency in human primary visual cortex. *J Neurosci*,

Unedited

29(50):15721-15726. https://doi.org/10.1523/JNEUROSCI.4426-09.2009
Eisenberg N, Fabes RA, Murphy B, et al., 1995. The role of emotionality and regulation in children's social functioning: A longitudinal study. *Child Dev*, 66(5):1360-1384.
Eisenberg N, Guthrie IK, Murphy BC, et al., 1999. Consistency and development of prosocial dispositions: A longitudinal study. *Child Dev*, 70(6):1360-1372. https://doi.org/10.1111/1467-8624.00100
Eisenberg N, Fabes RA, Guthrie IK, et al., 2000. Dispositional emotionality and regulation: Their role in predicting quality of social functioning. *J Pers Soc Psychol*, 78(1):136-157. https://doi.org/10.1037//0022-3514.78.1.136
Fan Z, Chang J, Liang Y, et al., 2023. Neural mechanism underlying depressive-like state associated with social status loss. *Cell*, 186(3):560-576 e517. https://doi.org/10.1016/j.cell.2022.12.033
Feinberg I, 1990. Cortical pruning and the development of schizophrenia. *Schizophr Bull*, 16(4):567-570. https://doi.org/10.1093/schbul/16.4.567
Filice F, Vorckel KJ, Sungur AO, et al., 2016. Reduction in parvalbumin expression not loss of the parvalbumin-expressing gaba interneuron subpopulation in genetic parvalbumin and shank mouse models of autism. *Mol Brain*, 9:10. https://doi.org/10.1186/s13041-016-0192-8
Forbes CE, Grafman J, 2010. The role of the human prefrontal cortex in social cognition and moral judgment. *Annu Rev Neurosci*, 33:299-324. https://doi.org/10.1146/annurev-neuro-060909-153230
Fornito A, Yucel M, Dean B, et al., 2009. Anatomical abnormalities of the anterior cingulate cortex in schizophrenia: Bridging the gap between neuroimaging and neuropathology. *Schizophr Bull*, 35(5):973-993. https://doi.org/10.1093/schbul/sbn025
Fries P, 2015. Rhythms for cognition: Communication through coherence. *Neuron*, 88(1):220-235. https://doi.org/10.1016/j.neuron.2015.09.034
Fujisawa S, Amarasingham A, Harrison MT, et al., 2008. Behavior-dependent short-term assembly dynamics in the medial prefrontal cortex. *Nat Neurosci*, 11(7):823-833. https://doi.org/10.1038/nn.2134
Fung SJ, Webster MJ, Sivagnanasundaram S, et al., 2010. Expression of interneuron markers in the dorsolateral prefrontal cortex of the developing human and in schizophrenia. *Am J Psychiatry*, 167(12):1479-1488. https://doi.org/10.1176/appi.ajp.2010.09060784
Gandal MJ, Edgar JC, Klook K, et al., 2012. Gamma synchrony: Towards a translational biomarker for the treatment-resistant symptoms of schizophrenia. *Neuropharmacology*, 62(3):1504-1518. https://doi.org/10.1016/j.neuropharm.2011.02.007
Gangopadhyay P, Chawla M, Dal Monte O, et al., 2021. Prefrontal-amygdala circuits in social decision-making. *Nat Neurosci*, 24(1):5-18. https://doi.org/10.1038/s41593-020-00738-9
Geckeler KC, Barch DM, Karcher NR, 2022. Associations between social behaviors and experiences with neural correlates of implicit emotion regulation in middle childhood. *Neuropsychopharmacology*, 47(6):1169-1179. https://doi.org/10.1038/s41386-022-01286-5
Giraldo-Chica M, Rogers BP, Damon SM, et al., 2018. Prefrontal-thalamic anatomical connectivity and executive cognitive function in schizophrenia. *Biol Psychiatry*, 83(6):509-517. https://doi.org/10.1016/j.biopsych.2017.09.022
Gogtay N, Giedd JN, Lusk L, et al., 2004. Dynamic mapping of human cortical development during childhood through early adulthood. *Proc Natl Acad Sci U S A*, 101(21):8174-8179. https://doi.org/10.1073/pnas.0402680101
Goldman-Rakic PS, Selemon LD, 1997. Functional and anatomical aspects of prefrontal pathology in schizophrenia. *Schizophr Bull*, 23(3):437-458. https://doi.org/10.1093/schbul/23.3.437
Han S, Tai C, Westenbroek RE, et al., 2012. Autistic-like behaviour in scn1a+/- mice and rescue by enhanced gaba-mediated neurotransmission. *Nature*, 489(7416):385-390. https://doi.org/10.1038/nature11356
Hanganu-Opatz IL, Klausberger T, Sigurdsson T, et al., 2023. Resolving the prefrontal mechanisms of adaptive cognitive behaviors: A cross-species perspective. *Neuron*, 111(7):1020-1036. https://doi.org/10.1016/j.neuron.2023.03.017
Hashimoto T, Volk DW, Eggan SM, et al., 2003. Gene expression deficits in a subclass of gaba neurons in the prefrontal cortex of subjects with schizophrenia. *J Neurosci*, 23(15):6315-6326. https://doi.org/10.1523/JNEUROSCI.23-15-06315.2003
Hashimoto T, Arion D, Unger T, et al., 2008. Alterations in gaba-related transcriptome in the dorsolateral prefrontal cortex of subjects with schizophrenia. *Mol Psychiatry*, 13(2):147-161. https://doi.org/10.1038/sj.mp.4002011
Hazlett HC, Poe M, Gerig G, et al., 2005. Magnetic resonance imaging and head circumference study of brain size in autism: Birth through age 2 years. *Arch Gen Psychiatry*, 62(12):1366-1376.

Unedited

https://doi.org/10.1001/archpsyc.62.12.1366
Herbert MR, Ziegler DA, Deutsch CK, et al., 2003. Dissociations of cerebral cortex, subcortical and cerebral white matter volumes in autistic boys. *Brain*, 126(Pt 5):1182-1192. https://doi.org/10.1093/brain/awg110
Huang AS, Rogers BP, Sheffield JM, et al., 2021. Characterizing effects of age, sex and psychosis symptoms on thalamocortical functional connectivity in youth. *Neuroimage*, 243:118562. https://doi.org/10.1016/j.neuroimage.2021.118562
Huang WC, Zucca A, Levy J, et al., 2020. Social behavior is modulated by valence-encoding mpfc-amygdala sub-circuitry. *Cell Rep*, 32(2):107899. https://doi.org/10.1016/j.celrep.2020.107899
Hughes JR, 2007. Autism: The first firm finding = underconnectivity? *Epilepsy Behav*, 11(1):20-24. https://doi.org/10.1016/j.yebeh.2007.03.010
Hughes JR, 2008. A review of recent reports on autism: 1000 studies published in 2007. *Epilepsy Behav*, 13(3):425-437. https://doi.org/10.1016/j.yebeh.2008.06.015
Kaczmarek LK, Zhang Y, 2017. Kv3 channels: Enablers of rapid firing, neurotransmitter release, and neuronal endurance. *Physiol Rev*, 97(4):1431-1468. https://doi.org/10.1152/physrev.00002.2017
Karlsgodt KH, Sanz J, Van Erp TG, et al., 2009. Re-evaluating dorsolateral prefrontal cortex activation during working memory in schizophrenia. *Schizophr Res*, 108(1-3):143-150. https://doi.org/10.1016/j.schres.2008.12.025
Kehrer C, Maziashvili N, Dugladze T, et al., 2008. Altered excitatory-inhibitory balance in the nmda-hypofunction model of schizophrenia. *Front Mol Neurosci*, 1:6. https://doi.org/10.3389/neuro.02.006.2008
Kietzman HW, Trinoskey-Rice G, Blumenthal SA, et al., 2022. Social incentivization of instrumental choice in mice requires amygdala-prelimbic cortex-nucleus accumbens connectivity. *Nat Commun*, 13(1):4768. https://doi.org/10.1038/s41467-022-32388-9
Kietzman HW, Gourley SL, 2023. How social information impacts action in rodents and humans: The role of the prefrontal cortex and its connections. *Neurosci Biobehav Rev*, 147:105075. https://doi.org/10.1016/j.neubiorev.2023.105075
Kimoto S, Bazmi HH, Lewis DA, 2014. Lower expression of glutamic acid decarboxylase 67 in the prefrontal cortex in schizophrenia: Contribution of altered regulation by zif268. *Am J Psychiatry*, 171(9):969-978. https://doi.org/10.1176/appi.ajp.2014.14010004
Kleinhans NM, Reiter MA, Neuhaus E, et al., 2016. Subregional differences in intrinsic amygdala hyperconnectivity and hypoconnectivity in autism spectrum disorder. *Autism Res*, 9(7):760-772. https://doi.org/10.1002/aur.1589
Klune CB, Jin B, Denardo LA, 2021. Linking mpfc circuit maturation to the developmental regulation of emotional memory and cognitive flexibility. *Elife*, 10 https://doi.org/10.7554/eLife.64567
Kolb B, Mychasiuk R, Muhammad A, et al., 2012. Experience and the developing prefrontal cortex. *Proc Natl Acad Sci U S A*, 109 Suppl 2(Suppl 2):17186-17193. https://doi.org/10.1073/pnas.1121251109
Kruse AO, Bustillo JR, 2022. Glutamatergic dysfunction in schizophrenia. *Transl Psychiatry*, 12(1):500. https://doi.org/10.1038/s41398-022-02253-w
Kumar A, Sundaram SK, Sivaswamy L, et al., 2010. Alterations in frontal lobe tracts and corpus callosum in young children with autism spectrum disorder. *Cereb Cortex*, 20(9):2103-2113. https://doi.org/10.1093/cercor/bhp278
Lee E, Rhim I, Lee JW, et al., 2016. Enhanced neuronal activity in the medial prefrontal cortex during social approach behavior. *J Neurosci*, 36(26):6926-6936. https://doi.org/10.1523/JNEUROSCI.0307-16.2016
Lee J, Chung C, Ha S, et al., 2015. Shank3-mutant mice lacking exon 9 show altered excitation/inhibition balance, enhanced rearing, and spatial memory deficit. *Front Cell Neurosci*, 9:94. https://doi.org/10.3389/fncel.2015.00094
Lewis DA, 1997. Development of the prefrontal cortex during adolescence: Insights into vulnerable neural circuits in schizophrenia. *Neuropsychopharmacology*, 16(6):385-398. https://doi.org/10.1016/S0893-133X(96)00277-1
Li K, Nakajima M, Ibanez-Tallon I, et al., 2016. A cortical circuit for sexually dimorphic oxytocin-dependent anxiety behaviors. *Cell*, 167(1):60-72 e11. https://doi.org/10.1016/j.cell.2016.08.067
Li Q, Takeuchi Y, Wang J, et al., 2023. Reinstating olfactory bulb-derived limbic gamma oscillations alleviates depression-like behavioral deficits in rodents. *Neuron*, 111(13):2065-2075 e2065. https://doi.org/10.1016/j.neuron.2023.04.013
Liang B, Zhang L, Barbera G, et al., 2018. Distinct and dynamic on and off neural ensembles in the prefrontal cortex code social exploration. *Neuron*, 100(3):700-714 e709. https://doi.org/10.1016/j.neuron.2018.08.043
Liang J, Xu W, Hsu YT, et al., 2015. Conditional neuroligin-2 knockout in adult medial prefrontal cortex links chronic

Unedited

changes in synaptic inhibition to cognitive impairments. *Mol Psychiatry*, 20(7):850-859. https://doi.org/10.1038/mp.2015.31

Liu D, Tang QQ, Yin C, et al., 2018. Brain-derived neurotrophic factor-mediated projection-specific regulation of depressive-like and nociceptive behaviors in the mesolimbic reward circuitry. *Pain*, 159(1):175. https://doi.org/10.1097/j.pain.0000000000001083

Liu L, Xu H, Wang J, et al., 2020. Cell type-differential modulation of prefrontal cortical gabaergic interneurons on low gamma rhythm and social interaction. *Sci Adv*, 6(30):eaay4073. https://doi.org/10.1126/sciadv.aay4073

Luna B, Minshew NJ, Garver KE, et al., 2002. Neocortical system abnormalities in autism: An fmri study of spatial working memory. *Neurology*, 59(6):834-840. https://doi.org/10.1212/wnl.59.6.834

Martinez K, Martinez-Garcia M, Marcos-Vidal L, et al., 2020. Sensory-to-cognitive systems integration is associated with clinical severity in autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry*, 59(3):422-433. https://doi.org/10.1016/j.jaac.2019.05.033

Martinez-Losa M, Tracy TE, Ma K, et al., 2018. Nav1.1-overexpressing interneuron transplants restore brain rhythms and cognition in a mouse model of alzheimer's disease. *Neuron*, 98(1):75-89 e75. https://doi.org/10.1016/j.neuron.2018.02.029

Martinez-Sanchis S, 2014. Neurobiological foundations of multisensory integration in people with autism spectrum disorders: The role of the medial prefrontal cortex. *Front Hum Neurosci*, 8:970. https://doi.org/10.3389/fnhum.2014.00970

Mcgraw LA, Young LJ, 2010. The prairie vole: An emerging model organism for understanding the social brain. *Trends Neurosci*, 33(2):103-109. https://doi.org/10.1016/j.tins.2009.11.006

Mcrae K, Gross JJ, 2020. Emotion regulation. *Emotion*, 20(1):1-9. https://doi.org/10.1037/emo0000703

Michelsen KA, Prickaerts J, Steinbusch HW, 2008. The dorsal raphe nucleus and serotonin: Implications for neuroplasticity linked to major depression and alzheimer's disease. *Prog Brain Res*, 172:233-264. https://doi.org/10.1016/S0079-6123(08)00912-6

Millan MJ, Bales KL, 2013. Towards improved animal models for evaluating social cognition and its disruption in schizophrenia: The cntrics initiative. *Neurosci Biobehav Rev*, 37(9 Pt B):2166-2180. https://doi.org/10.1016/j.neubiorev.2013.09.012

Mitchell JP, Macrae CN, Banaji MR, 2006. Dissociable medial prefrontal contributions to judgments of similar and dissimilar others. *Neuron*, 50(4):655-663. https://doi.org/10.1016/j.neuron.2006.03.040

Mitelman SA, Byne W, Kemether EM, et al., 2005. Metabolic disconnection between the mediodorsal nucleus of the thalamus and cortical brodmann's areas of the left hemisphere in schizophrenia. *Am J Psychiatry*, 162(9):1733-1735. https://doi.org/10.1176/appi.ajp.162.9.1733

Morris HM, Hashimoto T, Lewis DA, 2008. Alterations in somatostatin mrna expression in the dorsolateral prefrontal cortex of subjects with schizophrenia or schizoaffective disorder. *Cereb Cortex*, 18(7):1575-1587. https://doi.org/10.1093/cercor/bhm186

Moy SS, Nadler JJ, Perez A, et al., 2004. Sociability and preference for social novelty in five inbred strains: An approach to assess autistic-like behavior in mice. *Genes Brain Behav*, 3(5):287-302. https://doi.org/10.1111/j.1601-1848.2004.00076.x

Muller NG, Machado L, Knight RT, 2002. Contributions of subregions of the prefrontal cortex to working memory: Evidence from brain lesions in humans. *J Cogn Neurosci*, 14(5):673-686. https://doi.org/10.1162/08989290260138582

Murray AJ, Woloszynowska-Fraser MU, Ansel-Bollepalli L, et al., 2015. Parvalbumin-positive interneurons of the prefrontal cortex support working memory and cognitive flexibility. *Sci Rep*, 5:16778. https://doi.org/10.1038/srep16778

Murugan M, Jang HJ, Park M, et al., 2017. Combined social and spatial coding in a descending projection from the prefrontal cortex. *Cell*, 171(7):1663-1677 e1616. https://doi.org/10.1016/j.cell.2017.11.002

Nakajima M, Gorlich A, Heintz N, 2014. Oxytocin modulates female sociosexual behavior through a specific class of prefrontal cortical interneurons. *Cell*, 159(2):295-305. https://doi.org/10.1016/j.cell.2014.09.020

Nakazawa K, Jeevakumar V, Nakao K, 2017. Spatial and temporal boundaries of nmda receptor hypofunction leading to schizophrenia. *NPJ Schizophr*, 3:7. https://doi.org/10.1038/s41537-016-0003-3

Ogiwara I, Miyamoto H, Morita N, et al., 2007. Nav1.1 localizes to axons of parvalbumin-positive inhibitory interneurons: A circuit basis for epileptic seizures in mice carrying an scn1a gene mutation. *J Neurosci*, 27(22):5903-5914. https://doi.org/10.1523/JNEUROSCI.5270-06.2007

Ohnishi T, Matsuda H, Hashimoto T, et al., 2000. Abnormal regional cerebral blood flow in childhood autism. *Brain*, 123 (Pt 9):1838-1844. https://doi.org/10.1093/brain/123.9.1838

Okuyama T, Kitamura T, Roy DS, et al., 2016. Ventral ca1 neurons store social memory. *Science*,

Unedited

353(6307):1536-1541. https://doi.org/10.1126/science.aaf7003
Padilla-Coreano N, Batra K, Patarino M, et al., 2022. Cortical ensembles orchestrate social competition through hypothalamic outputs. *Nature*, 603(7902):667-671. https://doi.org/10.1038/s41586-022-04507-5
Palagi E, Burghardt GM, Smuts B, et al., 2016. Rough-and-tumble play as a window on animal communication. *Biol Rev Camb Philos Soc*, 91(2):311-327. https://doi.org/10.1111/brv.12172
Pellis SM, Pellis VC, Ham JR, et al., 2023. Play fighting and the development of the social brain: The rat's tale. *Neurosci Biobehav Rev*, 145:105037. https://doi.org/10.1016/j.neubiorev.2023.105037
Rafiee F, Rezvani Habibabadi R, Motaghi M, et al., 2022. Brain mri in autism spectrum disorder: Narrative review and recent advances. *J Magn Reson Imaging*, 55(6):1613-1624. https://doi.org/10.1002/jmri.27949
Rane P, Cochran D, Hodge SM, et al., 2015. Connectivity in autism: A review of mri connectivity studies. *Harv Rev Psychiatry*, 23(4):223-244. https://doi.org/10.1097/HRP.0000000000000072
Rein B, Tan T, Yang F, et al., 2021. Reversal of synaptic and behavioral deficits in a 16p11.2 duplication mouse model via restoration of the gaba synapse regulator npas4. *Mol Psychiatry*, 26(6):1967-1979. https://doi.org/10.1038/s41380-020-0693-9
Rudebeck PH, Walton ME, Millette BH, et al., 2007. Distinct contributions of frontal areas to emotion and social behaviour in the rat. *Eur J Neurosci*, 26(8):2315-2326. https://doi.org/10.1111/j.1460-9568.2007.05844.x
Sacco R, Gabriele S, Persico AM, 2015. Head circumference and brain size in autism spectrum disorder: A systematic review and meta-analysis. *Psychiatry Res*, 234(2):239-251. https://doi.org/10.1016/j.pscychresns.2015.08.016
Sarawagi A, Soni ND, Patel AB, 2021. Glutamate and gaba homeostasis and neurometabolism in major depressive disorder. *Front Psychiatry*, 12:637863. https://doi.org/10.3389/fpsyt.2021.637863
Sceniak MP, Lang M, Enomoto AC, et al., 2016. Mechanisms of functional hypoconnectivity in the medial prefrontal cortex of mecp2 null mice. *Cereb Cortex*, 26(5):1938-1956. https://doi.org/10.1093/cercor/bhv002
Scheggia D, Manago F, Maltese F, et al., 2020. Somatostatin interneurons in the prefrontal cortex control affective state discrimination in mice. *Nat Neurosci*, 23(1):47-60. https://doi.org/10.1038/s41593-019-0551-8
Selemon LD, Zecevic N, 2015. Schizophrenia: A tale of two critical periods for prefrontal cortical development. *Transl Psychiatry*, 5(8):e623. https://doi.org/10.1038/tp.2015.115
Selimbeyoglu A, Kim CK, Inoue M, et al., 2017. Modulation of prefrontal cortex excitation/inhibition balance rescues social behavior in cntnap2-deficient mice. *Sci Transl Med*, 9(401) https://doi.org/10.1126/scitranslmed.aah6733
Shemesh Y, Sztainberg Y, Forkosh O, et al., 2013. High-order social interactions in groups of mice. *Elife*, 2:e00759. https://doi.org/10.7554/eLife.00759
Sohal VS, Zhang F, Yizhar O, et al., 2009. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature*, 459(7247):698-702. https://doi.org/10.1038/nature07991
Su M, Hu K, Liu W, et al., 2023. Theta oscillations support prefrontal-hippocampal interactions in sequential working memory. *Neurosci Bull*, https://doi.org/10.1007/s12264-023-01134-6
Sun Q, Li X, Li A, et al., 2020. Ventral hippocampal-prefrontal interaction affects social behavior via parvalbumin positive neurons in the medial prefrontal cortex. *iScience*, 23(3):100894. https://doi.org/10.1016/j.isci.2020.100894
Tan Y, Singhal SM, Harden SW, et al., 2019. Oxytocin receptors are expressed by glutamatergic prefrontal cortical neurons that selectively modulate social recognition. *J Neurosci*, 39(17):3249-3263. https://doi.org/10.1523/JNEUROSCI.2944-18.2019
Uddin LQ, Supekar K, Menon V, 2013. Reconceptualizing functional brain connectivity in autism from a developmental perspective. *Front Hum Neurosci*, 7:458. https://doi.org/10.3389/fnhum.2013.00458
Varlinskaya EI, Spear LP, Spear NE, 1999. Social behavior and social motivation in adolescent rats: Role of housing conditions and partner's activity. *Physiol Behav*, 67(4):475-482. https://doi.org/10.1016/s0031-9384(98)00285-6
Vawter MP, Crook JM, Hyde TM, et al., 2002. Microarray analysis of gene expression in the prefrontal cortex in schizophrenia: A preliminary study. *Schizophr Res*, 58(1):11-20. https://doi.org/10.1016/s0920-9964(01)00377-2
Vissers ME, Cohen MX, Geurts HM, 2012. Brain connectivity and high functioning autism: A promising path of research that needs refined models, methodological convergence, and stronger behavioral links. *Neurosci Biobehav Rev*, 36(1):604-625. https://doi.org/10.1016/j.neubiorev.2011.09.003
Wang F, Zhu J, Zhu H, et al., 2011. Bidirectional control of social hierarchy by synaptic efficacy in medial prefrontal cortex. *Science*, 334(6056):693-697. https://doi.org/10.1126/science.1209951
Wang F, Kessels HW, Hu H, 2014. The mouse that roared: Neural mechanisms of social hierarchy. *Trends Neurosci*,

Accepted

37(11):674-682. https://doi.org/10.1016/j.tins.2014.07.005

Wang W, Rein B, Zhang F, et al., 2018. Chemogenetic activation of prefrontal cortex rescues synaptic and behavioral deficits in a mouse model of 16p11.2 deletion syndrome. *J Neurosci*, 38(26):5939-5948. https://doi.org/10.1523/JNEUROSCI.0149-18.2018

Weinberger DR, Berman KF, 1996. Prefrontal function in schizophrenia: Confounds and controversies. *Philos Trans R Soc Lond B Biol Sci*, 351(1346):1495-1503. https://doi.org/10.1098/rstb.1996.0135

Woo TU, Kim AM, Viscidi E, 2008. Disease-specific alterations in glutamatergic neurotransmission on inhibitory interneurons in the prefrontal cortex in schizophrenia. *Brain Res*, 1218:267-277. https://doi.org/10.1016/j.brainres.2008.03.092

Woodward ND, Karbasforoushan H, Heckers S, 2012. Thalamocortical dysconnectivity in schizophrenia. *Am J Psychiatry*, 169(10):1092-1099. https://doi.org/10.1176/appi.ajp.2012.12010056

Woodward ND, Heckers S, 2016. Mapping thalamocortical functional connectivity in chronic and early stages of psychotic disorders. *Biol Psychiatry*, 79(12):1016-1025. https://doi.org/10.1016/j.biopsych.2015.06.026

Yizhar O, Fenno LE, Prigge M, et al., 2011. Neocortical excitation/inhibition balance in information processing and social dysfunction. *Nature*, 477(7363):171-178. https://doi.org/10.1038/nature10360

Zhang C, Zhu H, Ni Z, et al., 2022. Dynamics of a disinhibitory prefrontal microcircuit in controlling social competition. *Neuron*, 110(3):516-531 e516. https://doi.org/10.1016/j.neuron.2021.10.034

Zhang M, Palaniyappan L, Deng M, et al., 2021. Abnormal thalamocortical circuit in adolescents with early-onset schizophrenia. *J Am Acad Child Adolesc Psychiatry*, 60(4):479-489. https://doi.org/10.1016/j.jaac.2020.07.903

Zhou T, Sandi C, Hu H, 2018. Advances in understanding neural mechanisms of social dominance. *Curr Opin Neurobiol*, 49:99-107. https://doi.org/10.1016/j.conb.2018.01.006

unedited