



## Review

<https://doi.org/10.1631/jzus.B2400127>



# Endomitosis: a new cell fate in the cell cycle leading to polyploidy in megakaryocytes and hepatocytes

Qi-Hua HUA<sup>1</sup>, Xuechun ZHANG<sup>1</sup>, Ruifeng TIAN<sup>2</sup>, Zhigang SHE<sup>2,3</sup>, Zan HUANG<sup>1</sup>✉

<sup>1</sup>College of Life Sciences, Key Laboratory of Cell Homeostasis of Hubei Province, Wuhan University, Wuhan 430060, China

<sup>2</sup>Department of Cardiology, Renmin Hospital of Wuhan University, Wuhan 430060, China

<sup>3</sup>Gannan Innovation and Translational Medicine Research Institute, Key Laboratory of Prevention and Treatment of Cardiovascular and Cerebrovascular Diseases, Ministry of Education, Gannan Medical University, Ganzhou 341000, China

**Abstract:** Megakaryocytes and hepatocytes are unique cells in mammals that undergo polyploidization through endomitosis in terminal differentiation. Many polyploidization regulators and underlying mechanisms have been reported, most of which are tightly coupled with development, organogenesis, and cell differentiation. However, the nature of endomitosis, which involves successful entry into and exit from mitosis without complete cytokinesis, has not yet been fully elucidated. We highlight that endomitosis is a new cell fate in the cell cycle, and tetraploidy is a critical stage at the bifurcation of cell fate decision. This review summarizes the recent research progress in this area and provides novel insights into how cells manipulate mitosis toward endomitosis. Endomitotic cells can evade the tetraploidy restrictions and proceed to multiple rounds of the cell cycle. This knowledge not only deepens our understanding of endomitosis as a fundamental biological process but also offers new perspectives on the physiological and pathophysiological implications of polyploidization.

**Key words:** Polyploidy; Endomitosis; Megakaryocyte; Hepatocyte; Cell cycle; Tetraploidy checkpoint; Whole-genome doubling

## 1 Introduction

Whole-organism polyploidy is commonly found in plants, fungi, and lower animals, where it is linked to diversity and evolution (Albertin and Marullo, 2012; van de Peer et al., 2017). In contrast, whole-organism polyploidy in mammals is not common and is highly debated, as it typically leads to embryonic resorption or spontaneous abortion (Svartman et al., 2005). At the cellular level, polyploidy is a distinct feature of specific and highly differentiated mammalian cell types, such as skeletal muscle cells (Abmayr and Pavlath, 2012), osteoclasts (Loutit and Nisbet, 1982), trophoblasts (Pandit et al., 2013), megakaryocytes (Ravid et al., 2002; Mazzi et al., 2018), hepatocytes (Guidotti et al., 2003; Wang et al., 2017; Donne et al., 2020; Sladky

et al., 2021), and cardiomyocytes (Gan et al., 2020). Polyploidy is tightly relevant to the physiological functions of these cells (Pandit et al., 2013). For instance, the degree of polyploidy is associated with the capability of megakaryocytes to produce platelets (Zimmet and Ravid, 2000; Bluteau et al., 2009). Polyploidy in the liver is believed to be associated with increased metabolic capacity (Fox and Duronio, 2013; van de Peer et al., 2017). Meanwhile, the impact of polyploidy on hepatocellular carcinoma remains controversial (Zhang et al., 2018b; Lin et al., 2020, 2021; Sladky et al., 2020b), which is attributed to the different genetic landscapes, such as *p53* mutation status (Bou-Nader et al., 2020; Sladky et al., 2020b). Osteoclasts depend on polyploidy for their effector function in bone resorption (Loutit and Nisbet, 1982). In the lactating mammary gland, binucleated polyploid epithelial cells are required for efficient milk production (Rios et al., 2016). Precise regulation of polyploidization in these cells is critical for health, and dysregulation of polyploidization involves many diseases including cancer.

✉ Zan HUANG, z-huang@whu.edu.cn

Zan HUANG, <https://orcid.org/0000-0001-5623-6950>

Qi-Hua HUA, <https://orcid.org/0009-0007-8599-5834>

Received Mar. 7, 2024; Revision accepted Aug. 31, 2024;  
Crosschecked Sept. 3, 2025

© Zhejiang University Press 2025

## 2 Route to polyploidy in mammalian cells

Polyploidy can be achieved through cell fusion, which is not dependent on the cell cycle or cell division. Endoreplication is another way to produce polyploidy, which continuously multiplies the genome without cell division. Endocycle (cycling cells proliferate without M phase) and endomitosis (cycling cells exit M phase and re-enter G1 phase without cell division) are two major forms of endoreplication. Trophoblast giant cell polyploidization is a typical example of endocycle (Zielke et al., 2013). Endomitosis is mainly found in megakaryocytes and hepatocytes (Nevzorova et al., 2009; Mazzi et al., 2018; Donne et al., 2020; Vainchenker and Raslova, 2020; Sladky et al., 2021; Sanz-Gómez et al., 2023). Notably, cytokinesis failure is sometimes regarded as a subtype of endomitosis (Zielke et al., 2013). Therefore, we will use “endomitosis” as a collective term to refer to a diversity of cell cycle programs in which cells progress to and exit M phase without complete cytokinesis (Herriage et al., 2024). In this review, we choose megakaryocytes and hepatocytes as two representative cell types that achieve their polyploidization via endomitosis. The canonical mechanisms have been well reviewed in other articles (Suraneni and Crispino, 2016; Mazzi et al., 2018; Donne et al., 2020; Vainchenker and Raslova, 2020; Sladky et al., 2021), and are concisely summarized in this review. We will further highlight endomitosis as a new cell fate in the cell cycle.

## 3 Canonical mechanisms of polyploidization

### 3.1 Megakaryocytes

Megakaryocytes are large (50–100  $\mu\text{m}$  in diameter), rare (0.05%–0.10%), polyploid, and platelet-producing hematopoietic cells that predominantly reside in the bone marrow in adults (Ebaugh and Bird, 1951). Recently, megakaryocytes have also been found in the lung and produce about 30% of platelets (Scheinin and Koivuniemi, 1963; Lefrançois et al., 2017). Megakaryocytes have a ploidy ranging from 2n to 128n, with a modal ploidy of 16n (Ravid et al., 2002). Megakaryocytes are differentiated from hematopoietic stem cells (HSCs), and megakaryocyte polyploidization is tightly coupled with differentiation and maturation, which are precisely regulated by many cytokines,

signaling pathways, transcription factors, and cell cycle regulators.

#### 3.1.1 Cytokines and signaling pathways

Thrombopoietin (TPO) is the primary cytokine regulating the entire process of megakaryopoiesis and thrombopoiesis (Nagata et al., 1997). TPO signals primarily through the Janus kinase (JAK)/signal transducer and activator of transcription (STAT), phosphoinositide 3-kinase (PI3K)/protein kinase B (AKT), and mitogen-activated protein kinase (MAPK) pathways (Chanprasert et al., 2006). TPO signaling through STAT proteins regulates the expression of cyclins D and E along with B-cell lymphoma-extra large (BCL-XL) (Eliades et al., 2010; Kaushansky, 2016), and induces modest increases in p21<sup>CIP1/WAF1</sup>, which may function as a licensing factor at low levels (Kikuchi et al., 1997). *Stat1*-deficient mice exhibit defects in ploidy, suggesting the importance of STAT1 in megakaryocyte polyploidization (Huang et al., 2007). STAT3 and STAT5 are two important downstream mediators of myeloproliferative leukemia virus oncogene (MPL)/TPO signaling that promote megakaryocyte development and proliferation (Yu et al., 2016). Furthermore, TPO signals to PI3K to increase polyploidy by facilitating cell cycle progression through the PI3K/AKT/forkhead box protein O3a (FOXO3a)/p27<sup>KIP1</sup> pathway (Nakao et al., 2008). The PI3K, STAT, and MAPK pathways also regulate *c-myc* expression (Chanprasert et al., 2006). The roles of mechanistic target of rapamycin (mTOR) and MAPK signaling remain controversial. One study demonstrated that mTOR inhibition had no effect on polyploidization (Drayer et al., 2006). However, another study demonstrated that mTOR induced enhanced polyploidization by inducing cyclin D3 and causing moderate upregulation of p21<sup>CIP1/WAF1</sup> (Raslova et al., 2006). Previous studies have reported that MAPK/extracellular signal-regulated kinase (ERK) inhibition restricts ploidy and differentiation in human cell lines and mouse primary cells (Rouyez et al., 1997; Mazharian et al., 2009), but it does not show any effect or enhancement in human primary cells (Guerriero et al., 2006). Distinctions between human and murine megakaryocyte polyploidization responses to ERK signaling cannot be excluded. The negative effects of the lysyl oxidase propeptide on polyploidization may be associated with the suppression of MAPK (Eliades et al., 2013). We and others have identified

regulators in these signaling pathways, including acidic nuclear phosphoprotein 32 family member A (ANP32A), proline-serine-threonine phosphatase-interacting protein 2 (PSTPIP2), and lymphocyte adaptor protein (LNK), which affect normal and abnormal megakaryopoiesis (Tong and Lodish, 2004; Liu et al., 2014; Sun et al., 2017). Finally, TPO may also modulate ploidy by inducing the formation of reactive oxygen species (ROS) through the upregulation of NADPH oxidase 1 (NOX1) (McCran et al., 2009). Inhibition of various NOX isoforms decreases polyploidization and the expression of cyclins D3 and E (McCran et al., 2009).

Many other hematopoietic cytokines also participate in megakaryocyte polyploidization. Members of the interleukin-6 (IL-6) family, including IL-6 and IL-11, promote polyploidy, whereas cytokines such as IL-3 can inhibit polyploidization (Burstein et al., 1992). Although all of these signals converge on JAK2 or JAK1 to activate STAT proteins, differences in the activation of STAT isoforms, signal kinetics, or receptor expression levels may contribute to these varied effects. In vitro studies have shown that the inhibition of transforming growth factor  $\beta$ -1 (TGF $\beta$ 1) signaling enhances TPO-driven polyploidization (Kuter et al., 1992). This effect is possibly mediated through increased expression of cyclin-dependent kinase (CDK) inhibitors (CDKIs), such as p27<sup>KIP1</sup>, p57<sup>KIP2</sup>, or p15<sup>INK4B</sup>. The C-X-C motif chemokine 12 (CXCL12)/C-X-C chemokine receptor type 4 (CXCR4) pathway may regulate polyploidization, but the role of CXCL12 in megakaryopoiesis remains contradictory (Hamada et al., 1998; Guerriero et al., 2001; Berthebaud et al., 2005; Stegner et al., 2017).

Diverse signaling pathways may also be involved in megakaryopoiesis and polyploidization, for instance, nuclear factor- $\kappa$ B (NF- $\kappa$ B) and Notch signaling (Zhang et al., 2002; Sun et al., 2004; Mercher et al., 2008; Weiss-Gayet et al., 2016). p21-activated kinase 2 (PAK2), the cell division control protein 42 homolog (CDC42)/Rac family small GTPase (RAC) pathway target, is suggested to restrain polyploidization in mice (Kosoff et al., 2015). Nicotinamide has been reported to elevate megakaryocyte ploidy in vitro, potentially through Sirtuin 1/2 inhibition (Giammona et al., 2009). The depletion of MAPK-interacting serine/threonine kinase 1 (MNK1) in cultured megakaryocytes reduces megakaryocyte ploidy in both mouse and human models (Manne et al., 2022).

### 3.1.2 Transcription factors

Megakaryocyte differentiation and polyploidization are tightly regulated by lineage-specific transcription factors, including GATA-binding factor 1 (GATA1), T-cell acute lymphocytic leukemia 1 (TAL1, also known as SCL), friend of GATA1 (FOG-1), growth factor-independent 1B (GFI1B), and nuclear factor, erythroid 2 (NF-E2) (Tijssen and Ghevaert, 2013). Monolineage production of megakaryocytes is driven by friend leukemia integration 1 transcription factor (FLI1), Runt-related transcription factor 1 (RUNX1), and GAB-binding protein  $\alpha$  chain (GABP $\alpha$ ) (Tijssen et al., 2011; Crispino and Weiss, 2014). The loss of GATA1, RUNX1, and FLI1 leads to defects in megakaryocyte differentiation and polyploidization (Tijssen and Ghevaert, 2013). The polyploidization defect caused by the loss of GATA1 can be partly attributed to the dysregulation of multiple genes. For instance, GATA1 upregulates cyclin D1 expression, and the megakaryopoiesis impairment as a result of GATA1 deficiency can be restored by the overexpression of cyclin D1-CDK4 in ex vivo cultured mouse megakaryocytes (Muntean et al., 2007). GATA1 also upregulates STAT1 (Huang et al., 2007) and cell division cycle 6 (CDC6) (Vilaboa et al., 2004) expression in megakaryocytes. Interestingly, estrogen increases polyploidization by elevating GATA1 transcriptional activity (Vilaboa et al., 2004). Furthermore, RUNX1 promotes polyploidization by silencing myosin heavy chain 10 (*MYH10*) and induces an arrest of endomitosis through the upregulation of p19<sup>INK4D</sup> in later stages of differentiation (Gilles et al., 2008; Lordier et al., 2012). Recently, studies have shown that *RUNX1* deficiency causes reduced polyploidization in both in vitro and in vivo human megakaryocytes, as well as in a rhesus macaque model (Bluteau et al., 2012; Lee BC et al., 2023; Lee K et al., 2023). FLI1 also regulates *MYH10* and possibly p19<sup>INK4D</sup> expression levels (Antony-Debré et al., 2012). RUNX1 and FLI1 not only regulate the transition from mitosis to endomitosis but also couple the arrest of DNA replication to megakaryocyte terminal differentiation (Huang H et al., 2009). TAL1/SCL has been shown to affect polyploidization by regulating p21<sup>CIP1/WAF1</sup> expression (Chagraoui et al., 2011). The role of NF-E2, a transcriptional target of GATA1, in megakaryocyte polyploidization is more complex. *Nfe2*-knockout megakaryocytes fail to undergo proplatelet formation but exhibit normal ploidy, which causes severe thrombocytopenia in mice

(Shivdasani et al., 1995). On the other hand, NF-E2 can induce polyploidization of cell lines and reprogram nonhematopoietic cells into polyploid megakaryocytes (Kobayashi et al., 1998; Ono et al., 2012). NF-E2 also co-regulates FLI1 and RUNX1 in late megakaryocyte differentiation (Zang et al., 2016). Indeed, these transcription factors collaborate in megakaryocyte differentiation and polyploidization. Genome-wide analyses have demonstrated combinatorial interactions among GATA1, GATA2, RUNX1, FLI1, and TAL1 in the early stages of megakaryopoiesis and cooperation among RUNX1, FLI1, and NF-E2 at later stages (Tijssen et al., 2011; Tijssen and Ghevaert, 2013).

GATA2, enhancer of zeste homolog 2 (EZH2), and aryl hydrocarbon receptor (AHR) play critical roles in HSC biology and regulate megakaryocyte polyploidization. Knockdown of *Gata2* impairs the polyploidization of megakaryocytes in vitro, and GATA2 promotes polyploidization of megakaryocytes in part by repressing the expression of myeloid transcription factors like purine-rich box 1 (PU.1) and CCAAT/enhancer-binding protein  $\alpha$  (C/EBP $\alpha$ ) (Huang Z et al., 2009). A recent ex vitro study demonstrates that EZH2 inhibition decreases the mean ploidy of megakaryocytes derived from human cord blood cluster of differentiation 34-positive (CD34<sup>+</sup>) cells (Mazzi et al., 2021). Specifically, EZH2 contributes to the effect of TPO on megakaryocyte commitment and permits normal polyploidization at the late stages of megakaryocyte differentiation by regulating *CDKN1A* and *CDKN2D* (Mazzi et al., 2021). *Ahr*-null mice exhibit a decrease in the percentage of high-ploidy megakaryocytes (>16n), likely due to the downregulation of hairy and enhancer of split 1 (HES1), a regulator of Notch signaling (Lindsey and Papoutsakis, 2011).

Some ubiquitous transcription factors are also involved in megakaryocyte polyploidization. Specifically, the megakaryoblastic leukemia 1 (MKL1)/serum response factor (SRF) complex plays a crucial role in megakaryocyte maturation by regulating the actomyosin cytoskeleton (Cheng et al., 2009; Gilles et al., 2009; Ragu et al., 2010; Elagib et al., 2022; Reed et al., 2022). Its transcriptional activity is regulated by actin polymerization through its cellular localization, connecting cytoskeleton remodeling to transcription (Cheng et al., 2009; Gilles et al., 2009; Smith et al., 2013). MKL1 downregulates guanine nucleotide exchange factor-H1 (GEF-H1, encoded by *ARHGEF2*)

and promotes megakaryocyte polyploidization in mouse models, possibly by inducing the transition from mitosis to endomitosis (Gao et al., 2012; Reed et al., 2022). The level of epithelial cell-transforming sequence 2 (ECT2) is reduced in higher-ploidy (>4n) megakaryocytes to regulate further polyploidization (Gao et al., 2012). Interestingly, RUNX1 and FLI1 bind to the *ARHGEF2* gene and may also regulate its expression (Tijssen et al., 2011).

### 3.1.3 Cell cycle regulators

Multiple cell cycle regulators are involved in megakaryocyte differentiation and polyploidization. Megakaryocytes express high levels of cyclin D1/D3, which elevate megakaryocyte number/ploidy but not platelet production (Wang et al., 1995; Sun et al., 2001). Conversely, cyclin E is essential for endomitosis as its overexpression increases megakaryocyte ploidy up to a modal 32n by regulating the pre-replication complex proteins minichromosome maintenance factor (MCM) and CDC6 (Bermejo et al., 2002; Geng et al., 2003; Eliades et al., 2010). CDC6 overexpression in megakaryocyte cell lines similarly elevates ploidy (Bermejo et al., 2002). Interestingly, cyclin B1 is present in megakaryocytes and undergoes conserved timing degradation during endomitosis (Zhang et al., 1996; Vitrat et al., 1998; Roy et al., 2001).

Megakaryocytes also express CDKs of the CDK-inhibitory protein/kinase inhibitory protein (Cip/Kip) (CDKN1) and inhibitor of CDK4 (INK4) (CDKN2) families, blocking G1/S progression. The role of INK4 family members during endomitosis remains unclear. One study found elevated p15<sup>INK4B</sup> expression in CD41<sup>+</sup> cells derived from human CD34<sup>+</sup> cells (Teofili et al., 2001), while another report showed decreased p16 expression during megakaryopoiesis (Furukawa et al., 2000). While all Cip/Kip family members (p21<sup>CIP1/WAF1</sup>, p27<sup>KIP1</sup>, and p57<sup>KIP2</sup>) are present (Taniguchi et al., 1999), only p19<sup>INK4D</sup> expression correlates positively with polyploidization (Raslova et al., 2007). p19<sup>INK4D</sup> uniquely impacts basal polyploidization and megakaryocyte maturation, as p19<sup>INK4D</sup> deficiency leads to an increase in the mean ploidy of both human megakaryocytes in vitro and mouse megakaryocytes in vivo (Gilles et al., 2008). Although p21 overexpression inhibits polyploidization (Kikuchi et al., 1997), p21 deletion in mice exerts no effect on polyploidization (Baccini et al., 2001). Furthermore, the loss of p53 in mouse and

cultured human megakaryocytes is associated with increased megakaryocyte ploidy levels, and this effect is exacerbated under stress conditions (Apostolidis et al., 2012; Roy et al., 2016b). p16 acts as an inhibitor of polyploidization by blocking the cyclin D-CDK4/6 complex, and its overexpression leads to the inhibition of endomitosis in ex vivo cultured mouse megakaryocytes (Muntean et al., 2007).

Chromosome passenger proteins may regulate endomitosis because of their functions in metaphase/anaphase, cytokinesis, and abscission. Aurora kinase A inhibition enhances polyploidization and differentiation in acute megakaryoblastic leukemia (Krause and Crispino, 2013). However, Aurora kinase A is necessary for mitosis but dispensable for endomitosis in normal megakaryopoiesis (Goldenson et al., 2015). Although Aurora kinase B is localized appropriately throughout endomitosis, it is non-essential for polyploidization (Lordier et al., 2010). Similarly, the inhibition of Polo-like kinase 1 (PLK1) causes polyploidization in the acute megakaryoblastic leukemia cell line (Wen et al., 2012), whereas *Plkl* knockout in mice induces megakaryocyte cell cycle arrest as well as spindle checkpoint activation (Trakala et al., 2015a). *Cdc20* null causes the failure of mitotic exit, leading to death and thrombocytopenia in mouse megakaryocytes (Trakala et al., 2015b). Interestingly, *Cdk1* knockout does not inhibit polyploidization but causes an endocycle for polyploidization in mouse megakaryocytes (Trakala et al., 2015b).

### 3.2 Hepatocytes

Hepatocyte polyploidization during development has been studied almost exclusively in rodent models (Guidotti et al., 2003; Margall-Ducos et al., 2007; Celton-Morizur et al., 2009). While all hepatocytes are diploid at birth, postnatal liver growth leads to the development of hepatocytes with varying ploidy levels, including diploid, tetraploid, and octoploid. The ploidy of a hepatocyte is determined by both its nuclear ploidy (diploid, tetraploid, and octoploid) and the number of nuclei (mononucleate and binucleate). Up to 90% of hepatocytes are polyploid in adult rodents, while adult humans have approximately 30% polyploid hepatocytes (Carriere, 1967; Kudryavtsev et al., 1993; Guidotti et al., 2003; Duncan et al., 2010; Donne et al., 2020). In the early postnatal stage, the liver of rodents consists of diploid and proliferative hepatocytes. Diploid

hepatocytes follow either a conventional cell cycle or endomitosis that involves incomplete cytokinesis (Margall-Ducos et al., 2007). In the following cell cycle, polyploid hepatocytes can undergo incomplete or complete cytokinesis, the latter increasing the number of polyploids (Guidotti et al., 2003). After this initial wave, the degree of hepatocyte polyploidy increases only slightly throughout the organism's lifespan.

#### 3.2.1 Cytokines and signaling pathways

During the postnatal growth stage, cytokinesis failure in hepatocytes occurs during weaning, a period characterized by significant changes in feeding behavior, hormone levels, and metabolic pathways. Insulin has been identified as a crucial factor in the development of binucleate tetraploid hepatocytes (Celton-Morizur et al., 2009, 2010). The transition from suckling milk to taking solid food in rodents causes an increase in insulin levels. Increased insulin activates the PI3K/AKT pathway and results in the delocalization of Ras homolog family member A (RhoA) and other essential cytokinesis effectors, which prevents the formation of cleavage furrows and ingression (Celton-Morizur et al., 2009). The exact mechanism linking insulin and RhoA impairment is not fully understood. In rodents, reduced insulin signaling significantly decreases the production of binucleate progenies, while repeated insulin injections promote the generation of polyploid hepatocytes (Celton-Morizur et al., 2009). The PI3K/AKT pathway is the primary downstream pathway that mediates the effects of insulin under these conditions. Multiple studies have revealed that PI3K/AKT regulates actin cytoskeleton polarization and reorganization during processes such as cell migration and invasion. In dividing hepatocytes, inhibiting AKT activity results in proper actin reorganization during mitosis and corrects RhoA localization at the division site, which are crucial for successful cytokinesis (Celton-Morizur et al., 2009). Phosphatase and tensin homolog (PTEN) is a negative regulator of the PI3K/AKT signaling pathway. Several studies have verified that *Pten*-deficient mouse hepatocytes exhibit increased ploidy in vivo (Gentric et al., 2015; Moreno et al., 2022).

Several other pathways have been found to negatively regulate polyploidization. Deficiencies in liver kinase B1 (LKB1) lead to hyperpolyploidy in mouse hepatocytes (Maillet et al., 2018). Mice lacking the

E3 ligase S-phase kinase-associated protein 2 (SKP2) accumulate p27, resulting in extensive liver polyploidization (Minamishima et al., 2002). Interestingly, the Hippo pathway also regulates hepatocyte polyploidy through SKP2 (Zhang et al., 2017). The activation of Yes-associated protein (YAP) causes AKT phosphorylation, leading to SKP2 cytoplasmic retention and, consequently, p27 accumulation in the nucleus (Zhang et al., 2017). Although previous studies in cell lines demonstrated the role of the Hippo pathway in preventing the cell cycle progression of tetraploids by activating p53, YAP activation in the liver results in increased ploidy independent of p53 (Ganem et al., 2014; Zhang et al., 2017). In addition, dysregulation of the degradation of the transcription factor FOXO1/3 results in hepatic hyperploidy (Zhang et al., 2017). Hepatocyte-specific *Foxo3* knockout reduced polyploidization of hepatocytes in a mouse model (Liang et al., 2022). In addition, wntless-type mouse mammary tumor virus (MMTV) integration site family (WNT) signaling inhibits binucleation and polyploidization of hepatocytes in an E2F7/8-dependent manner both in a mouse model and a human hepatocyte organoid model (Jin et al., 2022; Darmasaputra et al., 2024).

### 3.2.2 Transcription factors and cell cycle regulators

E2F transcription factors play important roles in the regulation of polyploidization during liver development (Conner et al., 2003; Chen et al., 2012; Pandit et al., 2012; Sladky et al., 2020a). This family of transcription factors consists of both activators (E2F1–3) and repressors (E2F4–6), as well as two atypical repressors (E2F7 and E2F8). Interestingly, the levels of E2F1–6 remain low throughout the lifespan of mice, while E2F7 and E2F8 are highly expressed during the initial seven weeks after birth (Pandit et al., 2012). This pattern coincides with liver polyploidization. Deficiencies in these atypical repressors result in a significant increase in diploid hepatocytes, possibly by dysregulating genes involved in cytokinesis, such as *Ect2*, mitotic kinesin-like protein 1 (*Mklp1*), and Rac GTPase-activating protein 1 (*Racgap1*, also known as *MgcRacGAP*) (Margall-Ducos et al., 2007). This is particularly interesting as these proteins are known to play crucial roles in the communication between the spindle and the cortex during hepatocyte polyploidization, prevent the formation of a functional contractile

actomyosin ring, and ultimately lead to impaired cytokinesis (Margall-Ducos et al., 2007). On the other hand, activators E2F1–3 initiate the transcription of their target genes, including S-phase cyclins E and A, various cytokinesis effectors, and their antagonists E2F7 and E2F8, which, in turn, repress the same set of target genes (Conner et al., 2003; Chen et al., 2012; Pandit et al., 2012; Bertoli et al., 2013). This feedback loop between E2F1 and E2F7/8 results in oscillating protein levels throughout the cell cycle. In mice, E2F1 deficiency alone leads to an increase in hepatocyte ploidy, while overexpression of E2F1 results in hypoploidy (Conner et al., 2003; Chen et al., 2012; Pandit et al., 2012).

Deficiencies in multiple cell cycle regulator genes, including *Cdk1* (Diril et al., 2012), *Trp53* (Sheahan et al., 2004; Kurinna et al., 2013), *Cdkn1a* (encoding p21) (Wu et al., 1996; Sheahan et al., 2004), *c-myc* (Baena et al., 2005), *Ccne* (encoding cyclin E) (Nevzorova et al., 2009), *Birc5* (encoding survivin) (Li et al., 2013), *Ssu72* (Kim et al., 2016), *Stk3* (Mst1/2) (Zhang et al., 2017), *Tgfb1* (de Santis Puzzon et al., 2016), *Skp2* (Minamishima et al., 2002; Zhang et al., 2017), and *Rb1* (Sheahan et al., 2004; Mayhew et al., 2005), have been shown to affect ploidy levels in hepatocytes in various mouse models. For instance, *Cdk1*-deficient hepatocytes exhibit enlarged nuclei with increased ploidy (Diril et al., 2012), indicating that endoreplication has occurred without cell division. Cyclins E1 and E2 have distinct and non-redundant functions in hepatocyte S phase entry and polyploidy during liver regeneration, with cyclin E1 promoting and cyclin E2 repressing polyploidization (Nevzorova et al., 2009).

### 3.2.3 Additional regulators

Anillin actin-binding protein (ANLN) is required for effective hepatocyte division (Zhang et al., 2018a, 2018b). *Anln*-deficient cells progress through metaphase but are unable to form or ingress the cleavage furrow in anaphase/telophase, leading to increased polyploidy (Zhang et al., 2018a). Interestingly, microRNAs (miRs) have also been identified as new regulators of hepatocyte polyploidy (Hsu et al., 2016). In particular, miR-122 exhibits differential expression during post-natal development and mice lacking miR-122 showed a significant decrease in the number of polyploid hepatocytes (Hsu et al., 2016). miR-122 directly targets


and inhibits pro-cytokinesis factors during liver development, and reduces the transcriptional levels of RhoA and cut-like homeobox 1 (CUX1), the latter being a transcriptional activator of ECT2 and central spindlein (Hsu et al., 2016).

#### 4 Endomitosis as a variant of mitosis in megakaryocytes and hepatocytes

Polyplloid megakaryocytes can enter mitosis and exhibit the formation of a complex multipolar spindle (with two poles between  $2n$  and  $4n$ , four poles between  $4n$  and  $8n$ , and so on) and an asymmetrical segregation of chromosomes towards the different poles (Vitrat et al., 1998; Roy et al., 2001; Geddis et al., 2007; Papadantonakis et al., 2008). Endomitosis was once believed to be an incomplete mitosis that skips telophase and cytokinesis (Nagata et al., 1997; Vitrat et al., 1998). In fact, endomitotic cells can proceed to telophase, and the two daughter cells are nearly separated with a seemingly normal midzone and the formation of a cleavage furrow, particularly in the transition from  $2n$  to  $4n$  (Geddis et al., 2007). However, this furrow regresses rapidly in endomitotic megakaryocytes, and the two daughter cells fuse together (Geddis et al., 2007). The abnormalities in furrow ingression are more significant in higher ploidy, leading to minimal cell elongation and chromosome segregation (Papadantonakis et al., 2008). A significantly prolonged M phase has been observed in endomitotic megakaryocytes (Trakala et al., 2015a), which may consequently lead to cell cycle exit without cytokinesis. In rodent hepatocytes, disruption of canonical cytokinesis resulting from the absence of actomyosin ring formation and impaired actin organization during anaphase-telophase has been observed (Celton-Morizur and Desdouets, 2010; Fortier et al., 2017). Consequently, essential microtubule-associated proteins, such as Aurora B, protein regulator of cytokinesis 1 (PRC1), MgcRacGAP, and PLK1, fail to localize properly, impeding RhoA concentration and contractile ring formation at the division site. Disorganization of microtubules during incomplete cytokinesis impairs molecular signaling to the equatorial cortex (Margall-Ducos et al., 2007). Active RhoA concentrates equatorially during normal cytokinesis but remains diffuse throughout the center of cells undergoing cytokinesis failure (Margall-Ducos et al., 2007).

These observations demonstrate that improper regulation of RhoA signaling stemming from disorganized microtubules during incomplete cytokinesis ultimately leads to the production of polyploid hepatocytes. Similar to observations in megakaryocytes, diploid hepatocytes are found to enter and complete the cell cycle more rapidly than polyploid hepatocytes (Wilkinson et al., 2019). Differences are also noted among polyploid hepatocytes, with tetraploids initiating cell cycle progression at a faster pace than octoploids, suggesting that cell cycle regulation may exhibit subtle variances between diploidy and polyploidy (Wilkinson et al., 2019). Notably, recent live-imaging analyses of fetal tissue-derived human hepatocyte organoids revealed that human hepatocytes undergoing endomitosis inhibit cell division during a late step in cytokinesis (Darmasaputra et al., 2024). More similar to megakaryocytes than rodent hepatocytes, endomitotic human hepatocytes exhibit normal cytokinetic furrow ingression; however, the ingressed membrane detaches from the midbody during late cytokinesis. Proteins involved in anchoring the midbody to the cell cortex are present but lose their association with the cell membrane during endomitotic furrow regression (Darmasaputra et al., 2024). These observations imply that endomitosis exhibits some typical characteristics of mitosis. Endomitosis essentially may be a variant of mitosis.

Endomitotic megakaryocytes and hepatocytes exhibit notable distinctions, particularly in the number of nuclei (Fig. 1). The majority of polyploid megakaryocytes are large mononucleated cells, while polyploid hepatocytes can be either mononucleated or binucleated. It has been hypothesized that the progression of endomitosis may determine the mono- or binucleation status: cytokinesis failure at late M phase results in binucleation, whereas exiting M phase at the early stage leads to mononucleation (Darmasaputra et al., 2024). In fact, megakaryocytes display significant furrow ingression and regression during the  $2n$  to  $4n$  transition, and a substantial proportion of  $4n$  megakaryocytes are truly binucleated (Lordier et al., 2008; Leysi-Derilou et al., 2010), indicating a late cytokinesis defect. In contrast, most high-ploidy megakaryocytes are mononucleated and exhibit measurable furrowing followed by regression (Lordier et al., 2008; Leysi-Derilou et al., 2010). In sharp contrast, rodent endomitotic hepatocytes lack a central spindle, cleavage furrow ingression, and anaphase cell elongation (Guidotti et al., 2003;



**Fig. 1** Illustrations of different nucleations of endomitosis in megakaryocytes and hepatocytes. Immunofluorescence images of endomitotic megakaryocytes and hepatocytes. (a) Polyploid megakaryocytes are typically mononucleated and possess a single nucleus that is multilobulated. Nuclei of polyploid megakaryocytes stained with 4',6-diamidino-2-phenylindole (DAPI) (left), a bright field image (middle), and a merged image (right). (b) Polyploid hepatocytes are mononucleated or binucleated. Nuclei of polyploid hepatocytes stained with DAPI (left), a bright field image (middle), and a merged image (right).

Margall-Ducos et al., 2007), suggesting that endomitosis in rodent hepatocytes corresponds to early cytokinesis failure, but still produces binucleated polyploid cells. Moreover, the formation of mononucleation in megakaryocytes has been proposed to be a karyokinesis defect with the persistent presence of nucleoplasmic bridges (Lordier et al., 2012), which is not observed in hepatocyte endomitosis (Margall-Ducos et al., 2007; Darmasaputra et al., 2024). Indeed, during liver growth, mononucleated polyploid hepatocytes are essentially derived from the division of binucleated polyploid hepatocytes (Donne et al., 2020). Whether cells become mononucleated or binucleated is not critical for endomitosis leading to polyploidy, while there might be functional distinctions between different nucleations (Katsuda et al., 2020; Richter et al., 2021; Dehn and Losick, 2022; van Rijnberk et al., 2022). Despite these differences, endomitotic cells exhibit remarkable similarities in their transition from diploidy to polyploidy (Ravid et al., 2002; Guidotti et al., 2003; Ganem et al., 2014; Gentric and Desdouets, 2014; Mazzi et al., 2018; Øvrebø and Edgar, 2018).

In general, commonalities and specificities of endomitosis exist in megakaryocytes and hepatocytes. Undoubtedly, endomitosis is a fundamental biological process rather than merely a developmentally coupled phenomenon. Whether cells decide to divide or not involves a cell fate decision during the cell cycle. The

fundamental mechanisms responsible for this cell fate decision involve elegant regulation of the M phase and remain to be addressed. How endomitotic cells manipulate mitosis to achieve this in megakaryocytes and hepatocytes warrants further clarification, which may reveal commonalities and specificities.


## 5 Mechanisms rewiring mitosis toward endomitosis for polyploidization

As summarized above, most studies on polyploidization regulators are tightly coupled with development, organogenesis, and cell differentiation. However, these studies have not addressed the essence of endomitosis, a fundamental biological process involving cell cycle regulation. Several key events involving the mitosis phase have to be modified for cells to progress to this cell fate (Fig. 2).

### 5.1 Endomitosis requires functional mechanisms regulating M-phase entry and exit

CDK1 is known to be a critical regulator of M-phase entry and is also indispensable for endomitosis (Santamaría et al., 2007). An initial study showed that the expression of cyclin B1, the CDK1 activity regulator, was undetectable or decreased in megakaryocytes (Zhang et al., 1996). However, later studies confirmed that both cyclin B1 and CDK1 are expressed in endomitotic megakaryocytes and are normally degraded by anaphase-promoting complex or cyclosome (APC/C) along with its co-activators, CDC20 and cadherin 1 (CDH1) (Vitrat et al., 1998; Roy et al., 2001). *Cdk1*-null megakaryocytes and hepatocytes both fail to enter M phase and proceed to endocycle rather than endomitosis (Diril et al., 2012; Trakala et al., 2015b).

As the activation of the spindle assembly checkpoint (SAC) is imperative for mitosis, where CDK1/cyclin B serves as a linchpin, the SAC may play a crucial role in endomitosis as well. CDK1 contributes to the SAC, which ensures the integrity of the chromosome and allows the M phase to proceed to anaphase (Bentley et al., 2007; Chen et al., 2008; Alfonso-Pérez et al., 2019). In fact, the SAC is present and functions in megakaryocytes undergoing endomitosis (Roy et al., 2001; Trakala et al., 2015a). Typically, the SAC will be activated and arrest the mitotic cells in metaphase until all chromosomes are fully attached (Musacchio



**Fig. 2** Crossroads of endomitotic cell fate decision. Endomitotic cells can enter and exit mitosis without undergoing complete cytokinesis. During endomitosis, cells manipulate multiple critical events in the cell cycle and bypass the tetraploidy checkpoint, thereby achieving polyploidy. The solid arrows in the schematic illustration represent the canonical route of endomitosis cell fate, while the dashed arrows indicate alternative cell fates, such as endocycle, mitosis, and cell death induced by the tetraploidy checkpoint. CDK1: cyclin-dependent kinase 1; SAC: spindle assembly checkpoint; RhoA: Ras homology family member A.

and Salmon, 2007). However, failure to satisfy the SAC sometimes causes cells to slip out of arrest, a phenomenon called mitotic slippage (Sinha et al., 2019). In this context, progressive cyclin B1 degradation allows cells to escape mitosis and reach the G1 phase, even when the SAC is active (Brito and Rieder, 2006). Furthermore, reduced SAC activity is suggested to promote polyploidization in megakaryocytes (Wang et al., 2004; Brito and Rieder, 2006; Trakala et al., 2015a). The level of SAC activity in endomitotic cells and its impact on polyploidization are attractive questions to explore.

PLK1 is believed to be involved in these processes. In megakaryocytes undergoing endomitosis, sister chromatid separation appears to occur asymmetrically toward opposite poles rather than symmetrically, with chromosomes clustering around each pole into rings, ultimately forming multilobed megakaryocytes (Roy et al., 2001). This asymmetrical separation into multiple poles suggests that the SAC allows megakaryocyte chromosomes to attach complexly to multiple spindle poles. This phenomenon is altered in *Plk1*-deficient megakaryocytes. With a deficiency of PLK1, all chromosomes form single or multiple rings around the sole or minority of poles, which is consistent with megakaryocytes having single or fewer lobes in vivo (Trakala et al., 2015a). These spindle assembly defects during formation could lead to SAC-dependent mitotic arrest, which typically triggers cell death (Trakala et al., 2015a; Sinha et al., 2019).

APC/C targeting cyclin B for its degradation also contributes to endomitosis, as a low level of cyclin B is a key factor for M-phase exit. The APC/C-CDC20 complex is an E3-ubiquitin ligase, which is essential for exiting mitosis by triggering the degradation of the CDK1 activator cyclin B1 (Peters, 2006). CDC20 ablation in mice does not affect interphase or mitotic entry, but these *Cdc20*-deficient cells are trapped in mitosis due to the absence of APC/C-CDC20 activity (Manchado et al., 2010). Deletion of *Cdc20* in mice causes the accumulation of abnormal small megakaryocytes that display hyper-condensed chromosomes and mitotic arrest (Trakala et al., 2015b). Over 80% of *Cdc20*-null megakaryocytes are arrested in metaphase and cannot exit from the M phase (Trakala et al., 2015b). In late mitosis and G1, APC/C associates with CDH1, which continues to keep mitotic cyclins low until the next G2/M, and this event is also conserved in endomitotic megakaryocytes (Roy et al., 2001; Peters, 2006).

In the conserved mitotic exit mechanism, the orchestration of kinases and phosphatases plays a pivotal role (Wurzenberger and Gerlich, 2011; Manic et al., 2017). The Aurora kinases A and B have both been proven to be dispensable for megakaryocyte polyploidization (Lordier et al., 2010; Goldenson et al., 2015). However, the inhibition of Aurora B profoundly modifies the endomitotic process by inducing a missegregation of chromosomes and a mitotic failure in anaphase (Lordier et al., 2010). This defect may be related

to the abnormal localization of Aurora B due to the inhibition of its kinase activity. Clarifying the role of other kinases and phosphatases in regulating endomitotic entry and exit will advance our understanding of endomitosis.

## 5.2 Incomplete cytokinesis is a characteristic of endomitosis

Endomitotic cells need to initiate and exit M phase without cytokinesis due to cytokinesis failure or mitotic slippage, as completing cytokinesis would result in the production of either two faithfully separated daughter cells or two aneuploid cells rather than a polyploid cell (Leysi-Derilou et al., 2010; Fededa and Gerlich, 2012; Lacroix and Maddox, 2012). Cytokinesis failure can arise through defects in any of the four stages of the process: (1) positioning of the division plane, (2) ingression of the cleavage furrow, (3) formation of the midbody, and (4) abscission (Wang et al., 2017). Although cytokinesis is inhibited at different stages of the M phase in megakaryocytes (Geddis et al., 2007; Gao et al., 2012), cardiomyocytes (Engel et al., 2006; Leone et al., 2018), and hepatocytes (Guidotti et al., 2003; Margall-Ducos et al., 2007; Celton-Morizur et al., 2009; Pandit et al., 2012), the downregulation or inhibition of key cytokinesis regulators seems to be a common feature.

The absence of active RhoA in the cleavage furrow seems to be the most critical factor, which can be attributed to either the abnormal localization of RhoA or a defect in RhoA activation. The RhoA pathway plays a key role in cleavage furrow formation and ingression by facilitating actomyosin ring assembly, which generates contractile forces for abscission (Vainchenker et al., 2021). RhoA activation at the cleavage furrow induces localized actin polymerization and myosin II accumulation/activation, allowing furrow progression and abscission. This is mediated via formin/profilin for F-actin polymerization, as well as Rho kinase (ROCK) and citron kinase phosphorylating myosin light chain 2 (MLC2) for myosin activation. In megakaryocyte endomitosis, there is defective accumulation of myosin and F-actin, despite residual myosin activity remaining (Geddis and Kaushansky, 2006; Lordier et al., 2008). The RhoA/ROCK pathway is downregulated during the process of megakaryocyte endomitosis (Lordier et al., 2008; Gao et al., 2012; Avanzi et al., 2014). However, megakaryocytes from mice conditionally lacking RhoA or Cdc42 do not show impaired

megakaryocyte endomitosis (Heib et al., 2021), suggesting a critical role of RhoA/Cdc42 in mitosis but not endomitosis, similar to survivin (Wen et al., 2009). Although RhoA properly localizes in the cleavage furrows, particularly during the transition from 2n to 4n, there is a defect in RhoA activation in megakaryocytes (Geddis and Kaushansky, 2006; Lordier et al., 2008). Consistently, incomplete cytokinesis in hepatocyte endomitosis is also characterized by an absence of active RhoA localization at the putative cleavage plane (Margall-Ducos et al., 2007). RhoA activity relies on its GEFs like GEF-H1 and ECT2, which localize to cleavage furrows. GEF-H1 downregulation coincides with 2n to 4n transitions, while ECT2 markedly decreases at higher-ploidy stages during megakaryocyte polyploidization (Gao et al., 2012). The downregulation of ECT2 in both hepatocytes and high-ploidy (>4n) megakaryocytes may explain the lack of significant ingression of the cleavage furrow during cytokinesis failure in both cell types (Papadantonakis et al., 2008; Gao et al., 2012; Pandit et al., 2012). During megakaryocyte differentiation, two types of myosin II are synthesized: myosin IIA (MYH9) and myosin IIB (MYH10). However, in megakaryocyte precursors, myosin IIB is almost the only myosin II to be recruited at the cleavage furrow due to the structural differences between myosin IIA and myosin IIB (Badirou et al., 2014; Roy et al., 2016a). Furthermore, during megakaryocyte endomitosis, the transcriptional downregulation of GEF-H1 and MYH10 leads to a marked defect of contractile forces in the cleavage furrow, potentially explaining the abortive cytokinesis (Gao et al., 2012; Lordier et al., 2012; Badirou et al., 2014). Additional mitotic spindle components may influence endomitosis, such as downregulated microtubule-destabilizing stathmin (Iancu-Rubin et al., 2005, 2011).

## 5.3 Tetraploidy is a critical stage for endomitosis

Tetraploid cells at the G2/M phase may proceed to cytokinesis for the next round of the cell cycle, whereas a prolonged G2/M phase causes apoptosis. Under rare conditions, a cell may bypass cell division and transit to G1 for the next cell cycle, an event termed whole-genome doubling (WGD). Unscheduled WGD cells typically face replicative stress and DNA damage and exhibit genomic instability due to unbalanced chromosome segregation in subsequent mitosis (Quinton et al.,

2021; Gemble et al., 2022; Lambuta et al., 2023). Typically, unscheduled WGD cells are arrested in the subsequent G1 phase due to the activation of p53 signaling and the Hippo pathway, a mechanism known as the “tetraploidy checkpoint,” and the associated cell death is termed “mitotic catastrophe” (Margolis et al., 2003; Vitale et al., 2011; Ganem et al., 2014). p53 activation prevents cells from re-entering the cell cycle, leading to senescence (Notterman et al., 1998; Andreassen et al., 2001; Aylon and Oren, 2011). Furthermore, the p53 and Hippo pathways are interconnected via large tumor suppressor homolog 2 (LATS2, a Hippo effector), ensuring post-WGD G1 arrest (Aylon et al., 2006; Ganem et al., 2014). The tetraploidy checkpoint is recognized as a significant barrier to WGD (Margolis et al., 2003; Ganem et al., 2014). Additionally, extra centrosomes may influence WGD. The PIDDosome acts as a sensor for supernumerary centrosomes, a hallmark of tetraploid cells (Fava et al., 2017). The PIDDosome, a multiprotein complex that activates caspase-2 (CASP2), includes the death domain (DD)-containing proteins p53-induced DD protein 1 (PIDD1) and caspase and RIP adapter with death domain (CRADD, also known as RIP-associated protein with a death domain (RAIDD)). The accumulation of extra centrosomes triggers the PIDDosome signaling cascade, resulting in cell cycle arrest (Fava et al., 2017). The Hippo pathway also senses extra centrosomes (Sanz-Gómez et al., 2023). The presence of supernumerary centrosomes in RPE1 cells inhibits RhoA GTPase activity and activates RAC1 (Ganem et al., 2014). This modification of the cytoskeleton GTPases RhoA and RAC1 activates the Hippo pathway as a protective mechanism (Yu et al., 2012; Ganem et al., 2014). In fact, p53, LATS2, and the PIDDosome are considered components of the “tetraploidy checkpoint” that prevents tetraploid cells from re-entering the cell cycle, as the inactivation of any of these genes reverses G1 arrest (Aylon et al., 2006; Aylon and Oren, 2011; Ganem et al., 2014; Sladky et al., 2020b).

Endomitotic cells must overcome tetraploidy restrictions to transit from diploidy to polyploidy. Consequently, it is plausible that hepatocytes and megakaryocytes have a unique ability to bypass the tetraploidy checkpoint and proceed to polyploidization (Roy et al., 2016b; Zhang et al., 2017). Although the precise mechanism remains largely unknown, low RhoA activity during megakaryocyte polyploidization fails to

activate the Hippo pathway *ex vivo*, resulting in reduced levels of phosphorylated YAP but a high overall level of total YAP protein. However, *YAP* knock-down does not affect the ploidy of *ex vivo* cultured human megakaryocytes (Roy et al., 2016b). In contrast, tetraploid hepatocytes in mice have been shown to activate the Hippo pathway *in vivo* (Ganem et al., 2014). The activation of YAP/tafazzin (TAZ) signaling, either by inhibiting the Hippo pathway or by direct YAP overexpression *in vivo*, leads to increased ploidy in hepatocytes (Lee et al., 2016; Zhang et al., 2017). Specifically, the inhibition of the Hippo pathway in hepatocytes *in vivo* results in the accumulation of supernumerary centrosomes and cytokinesis failure (Zhang et al., 2017).

p53 is often referred to as the “guardian of the ploidy” because of its role in preventing WGD cells from re-entering the cell cycle by arresting them in the G1 phase and eventually triggering a non-proliferative state (Cross et al., 1995; Andreassen et al., 2001; Aylon and Oren, 2011). To bypass the tetraploid barrier, endomitotic hepatocytes regulate ploidy via the PIDDosome-p53 pathway (Fava et al., 2017; Sladky et al., 2021, 2022). Unlike typical cells that undergo PIDDosome-mediated apoptosis following multiple rounds of centrosome duplication, hepatocytes downregulate CASP2 via E2F7 and E2F8 to attain a physiologically normal polyploid state (Sladky et al., 2020a). The E2F transcriptional circuit further promotes polyploidization by downregulating cytokinesis machinery. PIDDosome-deficient mice exhibit significantly increased hepatocyte ploidy levels. Similarly, the loss of p53 in mice and *p53* knockdown in cultured human megakaryocytes are both associated with increased megakaryocyte ploidy levels (Fuhrken et al., 2008; Apostolidis et al., 2012; Roy et al., 2016b). However, the roles of the PIDDosome-p53 pathway in regulating megakaryocyte ploidy and endomitosis are still under investigation. Recent studies have shown that PIDD1 is recruited to the older centriole by the centriolar distal appendage protein ankyrin repeat domain-containing protein 26 (ANKRD26), which is essential for PIDDosome activation in response to excess centrosomes (Burigotto et al., 2021; Evans et al., 2021). Interestingly, *ANKRD26* is typically silenced during the late stages of healthy megakaryopoiesis, and mutations in *ANKRD26* found in thrombocytopenic patients disrupt this repression (Bluteau et al., 2014; Burigotto et al.,

2021; Evans et al., 2021). Consequently, megakaryocytes from these patients exhibit reductions in ploidy (Bluteau et al., 2014). It is very likely that the reduction in megakaryocyte ploidy in these patients arises due to ANKRD26-mediated activation of the PIDDosome in megakaryocytes with extra centrosomes (Burigotto et al., 2021; Evans et al., 2021). Further research in this area may elucidate how megakaryocytes tolerate supernumerary centrosomes and naturally prevent PIDDosome activation during endomitosis.

Tetraploidy checkpoint may not be the only challenge that endomitotic cells face. A recent study indicated that significant replication stress is a major source of genetic and chromosomal instability in newly formed WGD cells (Gemble et al., 2022). Mechanistically, these defects arise from a shortage of essential proteins during the first G1/S transition after WGD induction, compromising the fidelity of DNA replication and leading to high levels of DNA damage and abnormal karyotypes (Gemble et al., 2022). Furthermore, a subsequent study suggested that the oncogenic transformation of tetraploid cells is linked to this protein insufficiency (Lambuta et al., 2023). As endomitosis is a process leading to a WGD event in a physiological context, G1 arrest triggered by the tetraploidy checkpoint is suggested to be crucial for endomitotic cells to restore protein levels (Lambuta et al., 2023). An earlier study supported this hypothesis, confirming that p21 is highly expressed in hypoploid (2n and 4n) and polyploid (at least 8n) megakaryocytes derived from human CD34<sup>+</sup> cells (Baccini et al., 2001). High levels of p27, p16, cyclin E, and cyclin D3 were also observed in these populations, along with a hypophosphorylated form of retinoblastoma protein (Rb), indicating that most hypoploid and polyploid megakaryocytes are G1-arrested cells (Baccini et al., 2001; Besancenot et al., 2010). Thus, it appears that endomitotic cells must overcome the tetraploidy restrictions to transition from the canonical cell cycle to endomitosis while also mitigating the detrimental effects on cell fitness to achieve a stable polyploid state.

## 6 Perspectives

Endomitosis is essentially a variant form of mitosis, and cells rewire mitosis toward endomitosis for polyploidization. Dysregulation of polyploidization is

associated with multiple types of diseases. For instance, myeloproliferative neoplasms exhibit increased proliferation and decreased polyploidization in megakaryocytes, leading to thrombocythemia (Wen et al., 2015). Loss of polyploidization causes acute megakaryoblastic leukemia, and the induction of polyploidization has been suggested as a novel differentiation therapy (Wen et al., 2012; Krause and Crispino, 2013). Further studies should focus on characterizing the unique features of the M phase in endomitotic cells. How tetraploid cells re-enter the G1 phase for the next round of the endomitotic cell cycle is a critical point. Recent technical developments allow us to distinguish polyploid cells in the M phase from those in the G1 phase (Sakaue-Sawano et al., 2008, 2013). With the emergence of single-cell and nuclear sequencing (Katsuda et al., 2020; Richter et al., 2021), further studies may decipher the gene expression programs in polyploid cells at the G1 and M phases.

Previous studies used nocodazole or other drugs to induce WGD (Gemble et al., 2022). It cannot be ruled out that the results of these studies are add-on effects of the drug treatment (Uetake and Sluder, 2004; Wong and Stearns, 2005). Additionally, the tetraploid cells in these studies did not truly enter the endomitosis cell fate, as these cells are prone to dividing in the following cell cycle rather than undergoing consecutive cycles of endomitosis to accomplish polyploidization. WGD has been observed in early and pre-malignant lesions of various tissues, and it is estimated to occur in approximately 30% of human cancers (Zack et al., 2013; Bielski et al., 2018; Quinton et al., 2021; Zeng et al., 2023; McKenney et al., 2024). These observations raise intriguing questions. Why do tetraploid cells originating from WGD not proceed to endomitosis? Why do megakaryocytes achieve such high ploidy via consecutive cycles of endomitosis but rarely undergo malignant transformation? Indeed, paradoxically, polyploidy has been shown to both promote and suppress tumorigenesis (Ganem et al., 2007; Zhang et al., 2018b, 2019; Bou-Nader et al., 2020). The formation of tetraploid and polyploid cells in physiological contexts, such as in megakaryocytes and hepatocytes, can serve as perfect models for studying these problems. Dissecting the mechanisms underlying endomitosis is important not only for understanding physiological polyploidization in normal cells but also for understanding related diseases.

## Acknowledgments

This research was supported by the National Natural Science Foundation of China (Nos. 32270643, 91957109, and 81870427).

## Author contributions

Qi-Hua HUA wrote the original manuscript. Qi-Hua HUA and Zan HUANG conceived, designed, and discussed the work, and revised the manuscript. Xuechun ZHANG and Ruifeng TIAN performed the experiments and prepared the immunofluorescence images for Fig. 1. Qi-Hua HUA created Fig. 2 and processed the immunofluorescence images in Fig. 1. Zan HUANG and Zhigang SHE administered and supervised this work. All authors have read and approved the final manuscript.

## Compliance with ethics guidelines

Qi-Hua HUA, Xuechun ZHANG, Ruifeng TIAN, Zhigang SHE, and Zan HUANG declare they have no conflicts of interest.

This review does not contain any studies with human or animal subjects performed by any of the authors.

## References

- Abmayr SM, Pavlath GK, 2012. Myoblast fusion: lessons from flies and mice. *Development*, 139(4):641-656. <https://doi.org/10.1242/dev.068353>
- Albertin W, Marullo P, 2012. Polyploidy in fungi: evolution after whole-genome duplication. *Proc Biol Sci*, 279(1738):2497-2509. <https://doi.org/10.1098/rspb.2012.0434>
- Alfonso-Pérez T, Hayward D, Holder J, et al., 2019. MAD1-dependent recruitment of CDK1-CCNB1 to kinetochores promotes spindle checkpoint signaling. *J Cell Biol*, 218(4):1108-1117. <https://doi.org/10.1083/jcb.201808015>
- Andreassen PR, Lohez OD, Lacroix FB, et al., 2001. Tetraploid state induces p53-dependent arrest of nontransformed mammalian cells in G1. *Mol Biol Cell*, 12(5):1315-1328. <https://doi.org/10.1091/mbc.12.5.1315>
- Antony-Debré I, Bluteau D, Itzykson R, et al., 2012. MYH10 protein expression in platelets as a biomarker of RUNX1 and FLI1 alterations. *Blood*, 120(13):2719-2722. <https://doi.org/10.1182/blood-2012-04-422352>
- Apostolidis PA, Woulfe DS, Chavez M, et al., 2012. Role of tumor suppressor p53 in megakaryopoiesis and platelet function. *Exp Hematol*, 40(2):131-142.e4. <https://doi.org/10.1016/j.exphem.2011.10.006>
- Avanzi MP, Goldberg F, Davila J, et al., 2014. Rho kinase inhibition drives megakaryocyte polyploidization and proplatelet formation through MYC and NFE2 downregulation. *Br J Haematol*, 164(6):867-876. <https://doi.org/10.1111/bjh.12709>
- Aylon Y, Oren M, 2011. p53: guardian of ploidy. *Mol Oncol*, 5(4):315-323. <https://doi.org/10.1016/j.molonc.2011.07.007>
- Aylon Y, Michael D, Shmueli A, et al., 2006. A positive feedback loop between the P53 and Lats2 tumor suppressors prevents tetraploidization. *Genes Dev*, 20(19):2687-2700. <https://doi.org/10.1101/gad.1447006>
- Baccini V, Roy L, Vitrat N, et al., 2001. Role of p21<sup>Cip1/Waf1</sup> in cell-cycle exit of endomitotic megakaryocytes. *Blood*, 98(12):3274-3282. <https://doi.org/10.1182/blood.v98.12.3274>
- Badirou I, Pan JJ, Legrand C, et al., 2014. Carboxyl-terminal-dependent recruitment of nonmuscle myosin II to megakaryocyte contractile ring during polyploidization. *Blood*, 124(16):2564-2568. <https://doi.org/10.1182/blood-2014-06-584995>
- Baena E, Gandarillas A, Vallespinós M, et al., 2005. c-Myc regulates cell size and ploidy but is not essential for post-natal proliferation in liver. *Proc Natl Acad Sci USA*, 102(20):7286-7291. <https://doi.org/10.1073/pnas.0409260102>
- Bentley AM, Normand G, Hoyt J, et al., 2007. Distinct sequence elements of cyclin B1 promote localization to chromatin, centrosomes, and kinetochores during mitosis. *Mol Biol Cell*, 18(12):4847-4858. <https://doi.org/10.1091/mbc.e06-06-0539>
- Bermejo R, Vilaboa N, Calés C, 2002. Regulation of CDC6, geminin, and CDT1 in human cells that undergo polyploidization. *Mol Biol Cell*, 13(11):3989-4000. <https://doi.org/10.1091/mbc.e02-04-0217>
- Berthebaud M, Riviere C, Jarrier P, et al., 2005. RGS16 is a negative regulator of SDF-1-CXCR4 signaling in megakaryocytes. *Blood*, 106(9):2962-2968. <https://doi.org/10.1182/blood-2005-02-0526>
- Bertoli C, Skotheim JM, de Bruin RAM, 2013. Control of cell cycle transcription during G1 and S phases. *Nat Rev Mol Cell Biol*, 14(8):518-528. <https://doi.org/10.1038/nrm3629>
- Besancenot R, Chaligné R, Tonetti C, et al., 2010. A senescence-like cell-cycle arrest occurs during megakaryocytic maturation: implications for physiological and pathological megakaryocytic proliferation. *PLoS Biol*, 8(9):e1000476. <https://doi.org/10.1371/journal.pbio.1000476>
- Bielski CM, Zehir A, Penson AV, et al., 2018. Genome doubling shapes the evolution and prognosis of advanced cancers. *Nat Genet*, 50(8):1189-1195. <https://doi.org/10.1038/s41588-018-0165-1>
- Bluteau D, Lordier L, di Stefano A, et al., 2009. Regulation of megakaryocyte maturation and platelet formation. *J Thromb Haemost*, 7(Suppl 1):227-234. <https://doi.org/10.1111/j.1538-7836.2009.03398.x>
- Bluteau D, Glembotsky AC, Raimbault A, et al., 2012. Dysmegakaryopoiesis of FPD/AML pedigrees with constitutional RUNX1 mutations is linked to myosin II deregulated expression. *Blood*, 120(13):2708-2718. <https://doi.org/10.1182/blood-2012-04-422337>
- Bluteau D, Balduini A, Balayn N, et al., 2014. Thrombocytopenia-associated mutations in the ANKRD26 regulatory region induce MAPK hyperactivation. *J Clin Invest*, 124(2):580-591. <https://doi.org/10.1172/jci71861>

- Bou-Nader M, Caruso S, Donne R, et al., 2020. Polyploidy spectrum: a new marker in HCC classification. *Gut*, 69(2): 355-364.  
<https://doi.org/10.1136/gutjnl-2018-318021>
- Brito DA, Rieder CL, 2006. Mitotic checkpoint slippage in humans occurs via cyclin B destruction in the presence of an active checkpoint. *Curr Biol*, 16(12):1194-1200.  
<https://doi.org/10.1016/j.cub.2006.04.043>
- Burigotto M, Mattivi A, Migliorati D, et al., 2021. Centriolar distal appendages activate the centrosome-PIDDosome-p53 signalling axis via ANKRD26. *EMBO J*, 40(4):e104844.  
<https://doi.org/10.15252/embj.2020104844>
- Burstein SA, Mei RL, Henthorn J, et al., 1992. Leukemia inhibitory factor and interleukin-11 promote maturation of murine and human megakaryocytes in vitro. *J Cell Physiol*, 153(2):305-312.  
<https://doi.org/10.1002/jcp.1041530210>
- Carriere R, 1967. Polyploid cell reproduction in normal adult rat liver. *Exp Cell Res*, 46(3):533-540.  
[https://doi.org/10.1016/0014-4827\(67\)90379-5](https://doi.org/10.1016/0014-4827(67)90379-5)
- Celton-Morizur S, Desdouets C, 2010. Polyploidization of liver cells. In: Poon RYC (Ed.), *Polyploidization and Cancer*. Springer, New York, p.123-135.  
[https://doi.org/10.1007/978-1-4419-6199-0\\_8](https://doi.org/10.1007/978-1-4419-6199-0_8)
- Celton-Morizur S, Merlen G, Couton D, et al., 2009. The insulin/Akt pathway controls a specific cell division program that leads to generation of binucleated tetraploid liver cells in rodents. *J Clin Invest*, 119(7):1880-1887.  
<https://doi.org/10.1172/jci38677>
- Celton-Morizur S, Merlen G, Couton D, et al., 2010. Polyploidy and liver proliferation: central role of insulin signaling. *Cell Cycle*, 9(3):460-466.  
<https://doi.org/10.4161/cc.9.3.10542>
- Chagraoui H, Kassouf M, Banerjee S, et al., 2011. SCL-mediated regulation of the cell-cycle regulator p21 is critical for murine megakaryopoiesis. *Blood*, 118(3):723-735.  
<https://doi.org/10.1182/blood-2011-01-328765>
- Chanprasert S, Geddis AE, Barroga C, et al., 2006. Thrombopoietin (TPO) induces *c-myc* expression through a PI3K- and MAPK-dependent pathway that is not mediated by Akt, PKC $\zeta$  or mTOR in TPO-dependent cell lines and primary megakaryocytes. *Cell Signal*, 18(8):1212-1218.  
<https://doi.org/10.1016/j.cellsig.2005.09.010>
- Chen HZ, Ouseph MM, Li J, et al., 2012. Canonical and atypical E2Fs regulate the mammalian endocycle. *Nat Cell Biol*, 14(11):1192-1202.  
<https://doi.org/10.1038/ncb2595>
- Chen Q, Zhang XY, Jiang Q, et al., 2008. Cyclin B1 is localized to unattached kinetochores and contributes to efficient microtubule attachment and proper chromosome alignment during mitosis. *Cell Res*, 18(2):268-280.  
<https://doi.org/10.1038/cr.2008.11>
- Cheng EC, Luo Q, Bruscia EM, et al., 2009. Role for MKL1 in megakaryocytic maturation. *Blood*, 113(12):2826-2834.  
<https://doi.org/10.1182/blood-2008-09-180596>
- Conner EA, Lemmer ER, Sánchez A, et al., 2003. E2F1 blocks and c-Myc accelerates hepatic ploidy in transgenic mouse models. *Biochem Biophys Res Commun*, 302(1):114-120.  
[https://doi.org/10.1016/s0006-291x\(03\)00125-6](https://doi.org/10.1016/s0006-291x(03)00125-6)
- Crispino JD, Weiss MJ, 2014. Erythro-megakaryocytic transcription factors associated with hereditary anemia. *Blood*, 123(20):3080-3088.  
<https://doi.org/10.1182/blood-2014-01-453167>
- Cross SM, Sanchez CA, Morgan CA, et al., 1995. A p53-dependent mouse spindle checkpoint. *Science*, 267(5202): 1353-1356.  
<https://doi.org/10.1126/science.7871434>
- Darmasaputra GS, Geerlings CC, Chuva de Sousa Lopes SM, et al., 2024. Binucleated human hepatocytes arise through late cytokinetic regression during endomitosis M phase. *J Cell Biol*, 223(8):e202403020.  
<https://doi.org/10.1083/jcb.202403020>
- Dehn AS, Losick VP, 2022. Binucleation ramps up gene expression meeting the physiological demands of an organism. *PLoS Biol*, 20(5):e3001639.  
<https://doi.org/10.1371/journal.pbio.3001639>
- de Santis Puzzon M, Cozzolino AM, Grassi G, et al., 2016. TGFbeta induces binucleation/polyploidization in hepatocytes through a Src-dependent cytokinesis failure. *PLoS ONE*, 11(11):e0167158.  
<https://doi.org/10.1371/journal.pone.0167158>
- Diril MK, Ratnacaram CK, Padmakumar VC, et al., 2012. Cyclin-dependent kinase 1 (Cdk1) is essential for cell division and suppression of DNA re-replication but not for liver regeneration. *Proc Natl Acad Sci USA*, 109(10):3826-3831.  
<https://doi.org/10.1073/pnas.1115201109>
- Donne R, Saroul-Ainama M, Cordier P, et al., 2020. Polyploidy in liver development, homeostasis and disease. *Nat Rev Gastroenterol Hepatol*, 17(7):391-405.  
<https://doi.org/10.1038/s41575-020-0284-x>
- Drayer AL, Olthof SGM, Vellenga E, 2006. Mammalian target of rapamycin is required for thrombopoietin-induced proliferation of megakaryocyte progenitors. *Stem Cells*, 24(1):105-114.  
<https://doi.org/10.1634/stemcells.2005-0062>
- Duncan AW, Taylor MH, Hickey RD, et al., 2010. The ploidy conveyor of mature hepatocytes as a source of genetic variation. *Nature*, 467(7316):707-710.  
<https://doi.org/10.1038/nature09414>
- Ebaugh FG Jr, Bird RM, 1951. The normal megakaryocyte concentration in aspirated human bone marrow. *Blood*, 6(1): 75-80.  
<https://doi.org/10.1182/blood.v6.1.75.75>
- Elagib KE, Brock A, Clementelli CM, et al., 2022. Relieving DYRK1A repression of MKL1 confers an adult-like phenotype to human infantile megakaryocytes. *J Clin Invest*, 132(19):e154839.  
<https://doi.org/10.1172/jci154839>
- Eliades A, Papadantonakis N, Ravid K, 2010. New roles for cyclin E in megakaryocytic polyploidization. *J Biol Chem*, 285(24):18909-18917.  
<https://doi.org/10.1074/jbc.M110.102145>
- Eliades A, Papadantonakis N, Matsuura S, et al., 2013. Megakaryocyte polyploidy is inhibited by lysyl oxidase propeptide. *Cell Cycle*, 12(8):1242-1250.  
<https://doi.org/10.4161/cc.24312>

- Engel FB, Schebesta M, Keating MT, 2006. Anillin localization defect in cardiomyocyte binucleation. *J Mol Cell Cardiol*, 41(4):601-612.  
<https://doi.org/10.1016/j.yjmcc.2006.06.012>
- Evans LT, Anglen T, Scott P, et al., 2021. ANKRD26 recruits PIDD1 to centriolar distal appendages to activate the PIDDosome following centrosome amplification. *EMBO J*, 40(4):e105106.  
<https://doi.org/10.15252/embj.2020105106>
- Fava LL, Schuler F, Sladky V, et al., 2017. The PIDDosome activates p53 in response to supernumerary centrosomes. *Genes Dev*, 31(1):34-45.  
<https://doi.org/10.1101/gad.289728.116>
- Fededa JP, Gerlich DW, 2012. Molecular control of animal cell cytokinesis. *Nat Cell Biol*, 14(5):440-447.  
<https://doi.org/10.1038/ncb2482>
- Fortier M, Celton-Morizur S, Desdouets C, 2017. Incomplete cytokinesis/binucleation in mammals: the powerful system of hepatocytes. *Methods Cell Biol*, 137:119-142.  
<https://doi.org/10.1016/bs.mcb.2016.04.006>
- Fox DT, Duronio RJ, 2013. Endoreplication and polyploidy: insights into development and disease. *Development*, 140(1):3-12.  
<https://doi.org/10.1242/dev.080531>
- Fuhrken PG, Apostolidis PA, Lindsey S, et al., 2008. Tumor suppressor protein p53 regulates megakaryocytic polyploidization and apoptosis. *J Biol Chem*, 283(23):15589-15600.  
<https://doi.org/10.1074/jbc.M801923200>
- Furukawa Y, Kikuchi J, Nakamura M, et al., 2000. Lineage-specific regulation of cell cycle control gene expression during haematopoietic cell differentiation. *Br J Haematol*, 110(3):663-673.  
<https://doi.org/10.1046/j.1365-2141.2000.02253.x>
- Gan PH, Patterson M, Sucof HM, 2020. Cardiomyocyte polyploidy and implications for heart regeneration. *Annu Rev Physiol*, 82:45-61.  
<https://doi.org/10.1146/annurev-physiol-021119-034618>
- Ganem NJ, Storchova Z, Pellman D, 2007. Tetraploidy, aneuploidy and cancer. *Curr Opin Genet Dev*, 17(2):157-162.  
<https://doi.org/10.1016/j.gde.2007.02.011>
- Ganem NJ, Cornils H, Chiu SY, et al., 2014. Cytokinesis failure triggers Hippo tumor suppressor pathway activation. *Cell*, 158(4):833-848.  
<https://doi.org/10.1016/j.cell.2014.06.029>
- Gao Y, Smith E, Ker E, et al., 2012. Role of RhoA-specific guanine exchange factors in regulation of endomitosis in megakaryocytes. *Dev Cell*, 22(3):573-584.  
<https://doi.org/10.1016/j.devcel.2011.12.019>
- Geddis AE, Kaushansky K, 2006. Endomitotic megakaryocytes form a midzone in anaphase but have a deficiency in cleavage furrow formation. *Cell Cycle*, 5(5):538-545.  
<https://doi.org/10.4161/cc.5.5.2537>
- Geddis AE, Fox NE, Tkachenko E, et al., 2007. Endomitotic megakaryocytes that form a bipolar spindle exhibit cleavage furrow ingression followed by furrow regression. *Cell Cycle*, 6(4):455-460.  
<https://doi.org/10.4161/cc.6.4.3836>
- Gemble S, Wardenaar R, Keuper K, et al., 2022. Genetic instability from a single S phase after whole-genome duplication. *Nature*, 604(7904):146-151.  
<https://doi.org/10.1038/s41586-022-04578-4>
- Geng Y, Yu QY, Sicinska E, et al., 2003. Cyclin E ablation in the mouse. *Cell*, 114(4):431-443.  
[https://doi.org/10.1016/s0092-8674\(03\)00645-7](https://doi.org/10.1016/s0092-8674(03)00645-7)
- Gentric G, Desdouets C, 2014. Polyploidization in liver tissue. *Am J Pathol*, 184(2):322-331.  
<https://doi.org/10.1016/j.ajpath.2013.06.035>
- Gentric G, Maillet V, Paradis V, et al., 2015. Oxidative stress promotes pathologic polyploidization in nonalcoholic fatty liver disease. *J Clin Invest*, 125(3):981-992.  
<https://doi.org/10.1172/jci73957>
- Giammona LM, Panuganti S, Kemper JM, et al., 2009. Mechanistic studies on the effects of nicotinamide on megakaryocytic polyploidization and the roles of NAD<sup>+</sup> levels and SIRT inhibition. *Exp Hematol*, 37(11):1340-1352.e3.  
<https://doi.org/10.1016/j.exphem.2009.08.004>
- Gilles L, Guïze R, Bluteau D, et al., 2008. P19INK4D links endomitotic arrest and megakaryocyte maturation and is regulated by AML-1. *Blood*, 111(8):4081-4091.  
<https://doi.org/10.1182/blood-2007-09-113266>
- Gilles L, Bluteau D, Boukour S, et al., 2009. MAL/SRF complex is involved in platelet formation and megakaryocyte migration by regulating MYL9 (MLC2) and MMP9. *Blood*, 114(19):4221-4232.  
<https://doi.org/10.1182/blood-2009-03-209932>
- Goldenson B, Kirsammer G, Stankiewicz MJ, et al., 2015. Aurora kinase A is required for hematopoiesis but is dispensable for murine megakaryocyte endomitosis and differentiation. *Blood*, 125(13):2141-2150.  
<https://doi.org/10.1182/blood-2014-12-615401>
- Guerrero R, Mattia G, Testa U, et al., 2001. Stromal cell-derived factor 1 $\alpha$  increases polyploidization of megakaryocytes generated by human hematopoietic progenitor cells. *Blood*, 97(9):2587-2595.  
<https://doi.org/10.1182/blood.v97.9.2587>
- Guerrero R, Parolini I, Testa U, et al., 2006. Inhibition of TPO-induced MEK or mTOR activity induces opposite effects on the ploidy of human differentiating megakaryocytes. *J Cell Sci*, 119(4):744-752.  
<https://doi.org/10.1242/jcs.02784>
- Guidotti JE, Br gerie O, Robert A, et al., 2003. Liver cell polyploidization: a pivotal role for binuclear hepatocytes. *J Biol Chem*, 278(21):19095-19101.  
<https://doi.org/10.1074/jbc.M300982200>
- Hamada T, M hle R, Hesselgesser J, et al., 1998. Transendothelial migration of megakaryocytes in response to stromal cell-derived factor 1 (SDF-1) enhances platelet formation. *J Exp Med*, 188(3):539-548.  
<https://doi.org/10.1084/jem.188.3.539>
- Heib T, Hermanns HM, Manukjan G, et al., 2021. RhoA/Cdc42 signaling drives cytoplasmic maturation but not endomitosis in megakaryocytes. *Cell Rep*, 35(6):109102.  
<https://doi.org/10.1016/j.celrep.2021.109102>
- Herriage HC, Huang YT, Calvi BR, 2024. The antagonistic relationship between apoptosis and polyploidy in development

- and cancer. *Semin Cell Dev Biol*, 156:35-43.  
<https://doi.org/10.1016/j.semcdb.2023.05.009>
- Hsu SH, Delgado ER, Otero PA, et al., 2016. MicroRNA-122 regulates polyploidization in the murine liver. *Hepatology*, 64(2):599-615.  
<https://doi.org/10.1002/hep.28573>
- Huang H, Yu M, Akie TE, et al., 2009. Differentiation-dependent interactions between RUNX-1 and FLI-1 during megakaryocyte development. *Mol Cell Biol*, 29(15):4103-4115.  
<https://doi.org/10.1128/MCB.00090-09>
- Huang Z, Richmond TD, Muntean AG, et al., 2007. STAT1 promotes megakaryopoiesis downstream of GATA-1 in mice. *J Clin Invest*, 117(12):3890-3899.  
<https://doi.org/10.1172/JCI33010>
- Huang Z, Dore LC, Li Z, et al., 2009. GATA-2 reinforces megakaryocyte development in the absence of GATA-1. *Mol Cell Biol*, 29(18):5168-5180.  
<https://doi.org/10.1128/MCB.00482-09>
- Iancu-Rubin C, Nasrallah CA, Atweh GF, 2005. Stathmin prevents the transition from a normal to an endomitotic cell cycle during megakaryocytic differentiation. *Cell Cycle*, 4(12):1774-1782.  
<https://doi.org/10.4161/cc.4.12.2171>
- Iancu-Rubin C, Gajzer D, Tripodi J, et al., 2011. Down-regulation of stathmin expression is required for megakaryocyte maturation and platelet production. *Blood*, 117(17):4580-4589.  
<https://doi.org/10.1182/blood-2010-09-305540>
- Jin YH, Anbarchian T, Wu P, et al., 2022. Wnt signaling regulates hepatocyte cell division by a transcriptional repressor cascade. *Proc Natl Acad Sci USA*, 119(30):e2203849119.  
<https://doi.org/10.1073/pnas.2203849119>
- Katsuda T, Hosaka K, Matsuzaki J, et al., 2020. Transcriptomic dissection of hepatocyte heterogeneity: linking ploidy, zonation, and stem/progenitor cell characteristics. *Cell Mol Gastroenterol Hepatol*, 9(1):161-183.  
<https://doi.org/10.1016/j.jcmgh.2019.08.011>
- Kaushansky K, 2016. Thrombopoietin and its receptor in normal and neoplastic hematopoiesis. *Thromb J*, 14(Suppl 1):40.  
<https://doi.org/10.1186/s12959-016-0095-z>
- Kikuchi J, Furukawa Y, Iwase S, et al., 1997. Polyploidization and functional maturation are two distinct processes during megakaryocytic differentiation: involvement of cyclin-dependent kinase inhibitor p21 in polyploidization. *Blood*, 89(11):3980-3990.  
<https://doi.org/10.1182/blood.V89.11.3980>
- Kim SH, Jeon Y, Kim HS, et al., 2016. Hepatocyte homeostasis for chromosome ploidy and liver function is regulated by Ssu72 protein phosphatase. *Hepatology*, 63(1):247-259.  
<https://doi.org/10.1002/hep.28281>
- Kobayashi S, Teramura M, Ito K, et al., 1998. Transcription factor NF-E2 is essential for the polyploidization of a human megakaryoblastic cell line, Meg-J. *Biochem Biophys Res Commun*, 247(1):65-69.  
<https://doi.org/10.1006/bbrc.1998.8736>
- Kosoff RE, Aslan JE, Kostyak JC, et al., 2015. Pak2 restrains endomitosis during megakaryopoiesis and alters cytoskeleton organization. *Blood*, 125(19):2995-3005.  
<https://doi.org/10.1182/blood-2014-10-604504>
- Krause DS, Crispino JD, 2013. Molecular pathways: induction of polyploidy as a novel differentiation therapy for leukemia. *Clin Cancer Res*, 19(22):6084-6088.  
<https://doi.org/10.1158/1078-0432.CCR-12-2604>
- Kudryavtsev BN, Kudryavtseva MV, Sakuta GA, et al., 1993. Human hepatocyte polyploidization kinetics in the course of life cycle. *Virchows Arch B Cell Pathol Incl Mol Pathol*, 64(6):387-393.  
<https://doi.org/10.1007/BF02915139>
- Kurinna S, Stratton SA, Coban Z, et al., 2013. p53 regulates a mitotic transcription program and determines ploidy in normal mouse liver. *Hepatology*, 57(5):2004-2013.  
<https://doi.org/10.1002/hep.26233>
- Kuter DJ, Gminski DM, Rosenberg RD, 1992. Transforming growth factor beta inhibits megakaryocyte growth and endomitosis. *Blood*, 79(3):619-626.  
<https://doi.org/10.1182/blood.V79.3.619.619>
- Lacroix B, Maddox AS, 2012. Cytokinesis, ploidy and aneuploidy. *J Pathol*, 226(2):338-351.  
<https://doi.org/10.1002/path.3013>
- Lambuta RA, Nanni L, Liu YL, et al., 2023. Whole-genome doubling drives oncogenic loss of chromatin segregation. *Nature*, 615(7954):925-933.  
<https://doi.org/10.1038/s41586-023-05794-2>
- Lee BC, Zhou YF, Bresciani E, et al., 2023. A RUNX1-FPDMM rhesus macaque model reproduces the human phenotype and predicts challenges to curative gene therapies. *Blood*, 141(3):231-237.  
<https://doi.org/10.1182/blood.2022018193>
- Lee DH, Park JO, Kim TS, et al., 2016. LATS-YAP/TAZ controls lineage specification by regulating TGF $\beta$  signaling and *Hnf4a* expression during liver development. *Nat Commun*, 7:11961.  
<https://doi.org/10.1038/ncomms11961>
- Lee K, Ahn HS, Estevez B, et al., 2023. RUNX1-deficient human megakaryocytes demonstrate thrombopoietic and platelet half-life and functional defects. *Blood*, 141(3):260-270.  
<https://doi.org/10.1182/blood.2022017561>
- Lefrançois E, Ortiz-Muñoz G, Cadrillier A, et al., 2017. The lung is a site of platelet biogenesis and a reservoir for haematopoietic progenitors. *Nature*, 544(7648):105-109.  
<https://doi.org/10.1038/nature21706>
- Leone M, Musa G, Engel FB, 2018. Cardiomyocyte binucleation is associated with aberrant mitotic microtubule distribution, mislocalization of RhoA and IQGAP3, as well as defective actomyosin ring anchorage and cleavage furrow ingression. *Cardiovasc Res*, 114(8):1115-1131.  
<https://doi.org/10.1093/cvr/cvy056>
- Leysi-Derilou Y, Robert A, Duchesne C, et al., 2010. Polyploid megakaryocytes can complete cytokinesis. *Cell Cycle*, 9(13):2589-2599.  
<https://doi.org/10.4161/cc.9.13.12078>
- Li D, Cen J, Chen XT, et al., 2013. Hepatic loss of survivin impairs postnatal liver development and promotes expansion of hepatic progenitor cells in mice. *Hepatology*, 58(6):2109-2121.  
<https://doi.org/10.1002/hep.26601>

- Liang CQ, Zhou DC, Peng WT, et al., 2022. FoxO3 restricts liver regeneration by suppressing the proliferation of hepatocytes. *NPJ Regen Med*, 7:33. <https://doi.org/10.1038/s41536-022-00227-6>
- Lin H, Huang YS, Fustin JM, et al., 2021. Hyperpolyploidization of hepatocyte initiates preneoplastic lesion formation in the liver. *Nat Commun*, 12:645. <https://doi.org/10.1038/s41467-020-20572-8>
- Lin YH, Zhang SY, Zhu M, et al., 2020. Mice with increased numbers of polyploid hepatocytes maintain regenerative capacity but develop fewer hepatocellular carcinomas following chronic liver injury. *Gastroenterology*, 158(6):1698-1712.e14. <https://doi.org/10.1053/j.gastro.2020.01.026>
- Lindsey S, Papoutsakis ET, 2011. The aryl hydrocarbon receptor (AHR) transcription factor regulates megakaryocytic polyploidization. *Br J Haematol*, 152(4):469-484. <https://doi.org/10.1111/j.1365-2141.2010.08548.x>
- Liu L, Wen Q, Gong R, et al., 2014. *PSTPIP2* dysregulation contributes to aberrant terminal differentiation in GATA-1-deficient megakaryocytes by activating LYN. *Cell Death Dis*, 5(1):e988. <https://doi.org/10.1038/cddis.2013.512>
- Lordier L, Jalil A, Aurade F, et al., 2008. Megakaryocyte endomitosis is a failure of late cytokinesis related to defects in the contractile ring and Rho/Rock signaling. *Blood*, 112(8):3164-3174. <https://doi.org/10.1182/blood-2008-03-144956>
- Lordier L, Chang YH, Jalil A, et al., 2010. Aurora B is dispensable for megakaryocyte polyploidization, but contributes to the endomitotic process. *Blood*, 116(13):2345-2355. <https://doi.org/10.1182/blood-2010-01-265785>
- Lordier L, Bluteau D, Jalil A, et al., 2012. RUNX1-induced silencing of non-muscle myosin heavy chain IIB contributes to megakaryocyte polyploidization. *Nat Commun*, 3:717. <https://doi.org/10.1038/ncomms1704>
- Loutit JF, Nisbet NW, 1982. The origin of osteoclasts. *Immunobiology*, 161(3-4):193-203. [https://doi.org/10.1016/s0171-2985\(82\)80074-0](https://doi.org/10.1016/s0171-2985(82)80074-0)
- Maillet V, Boussetta N, Leclerc J, et al., 2018. LKB1 as a gatekeeper of hepatocyte proliferation and genomic integrity during liver regeneration. *Cell Rep*, 22(8):1994-2005. <https://doi.org/10.1016/j.celrep.2018.01.086>
- Manchado E, Guillamot M, de Cárcer G, et al., 2010. Targeting mitotic exit leads to tumor regression in vivo: modulation by Cdk1, Mastl, and the PP2A/B55 $\alpha$ , $\delta$  phosphatase. *Cancer Cell*, 18(6):641-654. <https://doi.org/10.1016/j.ccr.2010.10.028>
- Manic G, Corradi F, Sistigu A, et al., 2017. Molecular regulation of the spindle assembly checkpoint by kinases and phosphatases. In: Galluzzi L (Ed.), *International Review of Cell and Molecular Biology*. Academic Press, Cambridge, p.105-161. <https://doi.org/10.1016/bs.ircmb.2016.08.004>
- Manne BK, Campbell RA, Bhatlekar S, et al., 2022. MAPK-interacting kinase 1 regulates platelet production, activation, and thrombosis. *Blood*, 140(23):2477-2489. <https://doi.org/10.1182/blood.2022015568>
- Margall-Ducos G, Celton-Morizur S, Couton D, et al., 2007. Liver tetraploidization is controlled by a new process of incomplete cytokinesis. *J Cell Sci*, 120(20):3633-3639. <https://doi.org/10.1242/jcs.016907>
- Margolis RL, Lohez OD, Andreassen PR, 2003. G1 tetraploidy checkpoint and the suppression of tumorigenesis. *J Cell Biochem*, 88(4):673-683. <https://doi.org/10.1002/jcb.10411>
- Mayhew CN, Bosco EE, Fox SR, et al., 2005. Liver-specific pRB loss results in ectopic cell cycle entry and aberrant ploidy. *Cancer Res*, 65(11):4568-4577. <https://doi.org/10.1158/0008-5472.CAN-04-4221>
- Mazharian A, Watson SP, Séverin S, 2009. Critical role for ERK1/2 in bone marrow and fetal liver-derived primary megakaryocyte differentiation, motility, and proplatelet formation. *Exp Hematol*, 37(10):1238-1249.e5. <https://doi.org/10.1016/j.exphem.2009.07.006>
- Mazzi S, Lordier L, Debili N, et al., 2018. Megakaryocyte and polyploidization. *Exp Hematol*, 57:1-13. <https://doi.org/10.1016/j.exphem.2017.10.001>
- Mazzi S, Dessen P, Vieira M, et al., 2021. Dual role of EZH2 in megakaryocyte differentiation. *Blood*, 138(17):1603-1614. <https://doi.org/10.1182/blood.2019004638>
- McCran DJ, Eliades A, Makitalo M, et al., 2009. Differential expression of NADPH oxidases in megakaryocytes and their role in polyploidy. *Blood*, 114(6):1243-1249. <https://doi.org/10.1182/blood-2008-12-195883>
- McKenney C, Lendner Y, Zuniga AG, et al., 2024. CDK4/6 activity is required during G<sub>1</sub> arrest to prevent stress-induced endoreplication. *Science*, 384(6695):ead2421. <https://doi.org/10.1126/science.adi2421>
- Mercher T, Cornejo MG, Sears C, et al., 2008. Notch signaling specifies megakaryocyte development from hematopoietic stem cells. *Cell Stem Cell*, 3(3):314-326. <https://doi.org/10.1016/j.stem.2008.07.010>
- Minamishima YA, Nakayama K, Nakayama KI, 2002. Recovery of liver mass without proliferation of hepatocytes after partial hepatectomy in Skp2-deficient mice. *Cancer Res*, 62(4):995-999.
- Moreno E, Matondo AB, Bongiovanni L, et al., 2022. Inhibition of polyploidization in *Pten*-deficient livers reduces steatosis. *Liver Int*, 42(11):2442-2452. <https://doi.org/10.1111/liv.15384>
- Muntean AG, Pang LY, Poncz M, et al., 2007. Cyclin D-Cdk4 is regulated by GATA-1 and required for megakaryocyte growth and polyploidization. *Blood*, 109(12):5199-5207. <https://doi.org/10.1182/blood-2006-11-059378>
- Musacchio A, Salmon ED, 2007. The spindle-assembly checkpoint in space and time. *Nat Rev Mol Cell Biol*, 8(5):379-393. <https://doi.org/10.1038/nrm2163>
- Nagata Y, Muro Y, Todokoro K, 1997. Thrombopoietin-induced polyploidization of bone marrow megakaryocytes is due to a unique regulatory mechanism in late mitosis. *J Cell Biol*, 139(2):449-457. <https://doi.org/10.1083/jcb.139.2.449>

- Nakao T, Geddis AE, Fox NE, et al., 2008. PI3K/Akt/FOXO3a pathway contributes to thrombopoietin-induced proliferation of primary megakaryocytes in vitro and in vivo via modulation of p27Kip1. *Cell Cycle*, 7(2):257-266. <https://doi.org/10.4161/cc.7.2.5148>
- Nevzorova YA, Tschaharganeh D, Gassler N, et al., 2009. Aberrant cell cycle progression and endoreplication in regenerating livers of mice that lack a single E-type cyclin. *Gastroenterology*, 137(2):691-703.e6. <https://doi.org/10.1053/j.gastro.2009.05.003>
- Notterman D, Young S, Wainger B, et al., 1998. Prevention of mammalian DNA reduplication, following the release from the mitotic spindle checkpoint, requires p53 protein, but not p53-mediated transcriptional activity. *Oncogene*, 17(21):2743-2751. <https://doi.org/10.1038/sj.onc.1202210>
- Ono Y, Wang YH, Suzuki H, et al., 2012. Induction of functional platelets from mouse and human fibroblasts by p45NF-E2/Maf. *Blood*, 120(18):3812-3821. <https://doi.org/10.1182/blood-2012-02-413617>
- Øvrebø JI, Edgar BA, 2018. Polyploidy in tissue homeostasis and regeneration. *Development*, 145(14):dev156034. <https://doi.org/10.1242/dev.156034>
- Pandit SK, Westendorp B, Nantasanti S, et al., 2012. E2F8 is essential for polyploidization in mammalian cells. *Nat Cell Biol*, 14(11):1181-1191. <https://doi.org/10.1038/ncb2585>
- Pandit SK, Westendorp B, de Bruin A, 2013. Physiological significance of polyploidization in mammalian cells. *Trends Cell Biol*, 23(11):556-566. <https://doi.org/10.1016/j.tcb.2013.06.002>
- Papadantonakis N, Makitalo M, McCrann D, et al., 2008. Direct visualization of the endomitotic cell cycle in living megakaryocytes: differential patterns in low and high ploidy cells. *Cell Cycle*, 7(15):2352-2356. <https://doi.org/10.4161/cc.6325>
- Peters JM, 2006. The anaphase promoting complex/cyclosome: a machine designed to destroy. *Nat Rev Mol Cell Biol*, 7(9):644-656. <https://doi.org/10.1038/nrm1988>
- Quinton RJ, Didomizio A, Vittoria MA, et al., 2021. Whole-genome doubling confers unique genetic vulnerabilities on tumour cells. *Nature*, 590(7846):492-497. <https://doi.org/10.1038/s41586-020-03133-3>
- Ragu C, Boukour S, Elain G, et al., 2010. The serum response factor (SRF)/megakaryocytic acute leukemia (MAL) network participates in megakaryocyte development. *Leukemia*, 24(6):1227-1230. <https://doi.org/10.1038/leu.2010.80>
- Raslova H, Baccini V, Loussaief L, et al., 2006. Mammalian target of rapamycin (mTOR) regulates both proliferation of megakaryocyte progenitors and late stages of megakaryocyte differentiation. *Blood*, 107(6):2303-2310. <https://doi.org/10.1182/blood-2005-07-3005>
- Raslova H, Kauffmann A, Sekkai D, et al., 2007. Interrelation between polyploidization and megakaryocyte differentiation: a gene profiling approach. *Blood*, 109(8):3225-3234. <https://doi.org/10.1182/blood-2006-07-037838>
- Ravid K, Lu J, Zimmet JM, et al., 2002. Roads to polyploidy: the megakaryocyte example. *J Cell Physiol*, 190(1):7-20. <https://doi.org/10.1002/jcp.10035>
- Reed FE, Eskow NM, Min E, et al., 2022. Structure-function analysis of the role of megakaryoblastic leukemia 1 in megakaryocyte polyploidization. *Haematologica*, 107(12):2972-2976. <https://doi.org/10.3324/haematol.2021.280499>
- Richter ML, Deligiannis IK, Yin K, et al., 2021. Single-nucleus RNA-seq2 reveals functional crosstalk between liver zonation and ploidy. *Nat Commun*, 12:4264. <https://doi.org/10.1038/s41467-021-24543-5>
- Rios AC, Fu NY, Jamieson PR, et al., 2016. Essential role for a novel population of binucleated mammary epithelial cells in lactation. *Nat Commun*, 7:11400. <https://doi.org/10.1038/ncomms11400>
- Rouyez MC, Boucheron C, Gisselbrecht S, et al., 1997. Control of thrombopoietin-induced megakaryocytic differentiation by the mitogen-activated protein kinase pathway. *Mol Cell Biol*, 17(9):4991-5000. <https://doi.org/10.1128/MCB.17.9.4991>
- Roy A, Lordier L, Mazzi S, et al., 2016a. Activity of nonmuscle myosin II isoforms determines localization at the cleavage furrow of megakaryocytes. *Blood*, 128(26):3137-3145. <https://doi.org/10.1182/blood-2016-04-711630>
- Roy A, Lordier L, Pioche-Durieu C, et al., 2016b. Uncoupling of the Hippo and Rho pathways allows megakaryocytes to escape the tetraploid checkpoint. *Haematologica*, 101(12):1469-1478. <https://doi.org/10.3324/haematol.2016.149914>
- Roy L, Coullin P, Vitrat N, et al., 2001. Asymmetrical segregation of chromosomes with a normal metaphase/anaphase checkpoint in polyploid megakaryocytes. *Blood*, 97(8):2238-2247. <https://doi.org/10.1182/blood.v97.8.2238>
- Sakaue-Sawano A, Kurokawa H, Morimura T, et al., 2008. Visualizing spatiotemporal dynamics of multicellular cell-cycle progression. *Cell*, 132(3):487-498. <https://doi.org/10.1016/j.cell.2007.12.033>
- Sakaue-Sawano A, Hoshida T, Yo M, et al., 2013. Visualizing developmentally programmed endoreplication in mammals using ubiquitin oscillators. *Development*, 140(22):4624-4632. <https://doi.org/10.1242/dev.099226>
- Santamaría D, Barrière C, Cerqueira A, et al., 2007. Cdk1 is sufficient to drive the mammalian cell cycle. *Nature*, 448(7155):811-815. <https://doi.org/10.1038/nature06046>
- Sanz-Gómez N, González-Álvarez M, de Las Rivas J, et al., 2023. Whole-genome doubling as a source of cancer: how, when, where, and why? *Front Cell Dev Biol*, 11:1209136. <https://doi.org/10.3389/fcell.2023.1209136>
- Scheinin TM, Koivuniemi AP, 1963. Megakaryocytes in the pulmonary circulation. *Blood*, 22(1):82-87. <https://doi.org/10.1182/blood.V22.1.82.82>
- Sheahan S, Bellamy CO, Treanor L, et al., 2004. Additive effect of p53, p21 and Rb deletion in triple knockout primary hepatocytes. *Oncogene*, 23(8):1489-1497.

- <https://doi.org/10.1038/sj.onc.1207280>
- Shivdasani RA, Rosenblatt MF, Zucker-Franklin D, et al., 1995. Transcription factor NF-E2 is required for platelet formation independent of the actions of thrombopoietin/MGDF in megakaryocyte development. *Cell*, 81(5):695-704.  
[https://doi.org/10.1016/0092-8674\(95\)90531-6](https://doi.org/10.1016/0092-8674(95)90531-6)
- Sinha D, Duijf PHG, Khanna KK, 2019. Mitotic slippage: an old tale with a new twist. *Cell Cycle*, 18(1):7-15.  
<https://doi.org/10.1080/15384101.2018.1559557>
- Sladky VC, Knapp K, Soratroi C, et al., 2020a. E2F-family members engage the PIDDosome to limit hepatocyte ploidy in liver development and regeneration. *Dev Cell*, 52(3):335-349.e7.  
<https://doi.org/10.1016/j.devcel.2019.12.016>
- Sladky VC, Knapp K, Szabo TG, et al., 2020b. PIDDosome-induced p53-dependent ploidy restriction facilitates hepatocarcinogenesis. *EMBO Rep*, 21(12):e50893.  
<https://doi.org/10.15252/embr.202050893>
- Sladky VC, Eichin F, Reiberger T, et al., 2021. Polyploidy control in hepatic health and disease. *J Hepatol*, 75(5):1177-1191.  
<https://doi.org/10.1016/j.jhep.2021.06.030>
- Sladky VC, Akbari H, Tapias-Gomez D, et al., 2022. Centriole signaling restricts hepatocyte ploidy to maintain liver integrity. *Genes Dev*, 36(13-14):843-856.  
<https://doi.org/10.1101/gad.349727.122>
- Smith EC, Teixeira AM, Chen RC, et al., 2013. Induction of megakaryocyte differentiation drives nuclear accumulation and transcriptional function of MKL1 via actin polymerization and RhoA activation. *Blood*, 121(7):1094-1101.  
<https://doi.org/10.1182/blood-2012-05-429993>
- Stegner D, Vaneeuwijk JMM, Angay O, et al., 2017. Thrombopoiesis is spatially regulated by the bone marrow vasculature. *Nat Commun*, 8:127.  
<https://doi.org/10.1038/s41467-017-00201-7>
- Sun L, Tan P, Yap C, et al., 2004. In vitro biological characteristics of human cord blood-derived megakaryocytes. *Ann Acad Med Singap*, 33(5):570-575.  
<https://doi.org/10.47102/annals-acadmedsg.V33N5p570>
- Sun S, Zimmer JM, Toselli P, et al., 2001. Overexpression of cyclin D1 moderately increases ploidy in megakaryocytes. *Haematologica*, 86(1):17-23.
- Sun XQ, Lu B, Han CJ, et al., 2017. ANP32A dysregulation contributes to abnormal megakaryopoiesis in acute megakaryoblastic leukemia. *Blood Cancer J*, 7(12):661.  
<https://doi.org/10.1038/s41408-017-0031-x>
- Suraneni PK, Crispino JD, 2016. The Hippo-p53 pathway in megakaryopoiesis. *Haematologica*, 101(12):1446-1448.  
<https://doi.org/10.3324/haematol.2016.156125>
- Svartman M, Stone G, Stanyon R, 2005. Molecular cytogenetics discards polyploidy in mammals. *Genomics*, 85(4):425-430.  
<https://doi.org/10.1016/j.ygeno.2004.12.004>
- Taniguchi T, Endo H, Chikatsu N, et al., 1999. Expression of p21<sup>Cip1/Waf1/Sdi1</sup> and p27<sup>Kip1</sup> cyclin-dependent kinase inhibitors during human hematopoiesis. *Blood*, 93(12):4167-4178.  
<https://doi.org/10.1182/blood.V93.12.4167>
- Teofili L, Martini M, di Mario A, et al., 2001. Expression of p15<sup>ink4b</sup> gene during megakaryocytic differentiation of normal and myelodysplastic hematopoietic progenitors. *Blood*, 98(2):495-497.  
<https://doi.org/10.1182/blood.v98.2.495>
- Tijssen MR, Ghevaert C, 2013. Transcription factors in late megakaryopoiesis and related platelet disorders. *J Thromb Haemost*, 11(4):593-604.  
<https://doi.org/10.1111/jth.12131>
- Tijssen MR, Cvejic A, Joshi A, et al., 2011. Genome-wide analysis of simultaneous GATA1/2, RUNX1, FLI1, and SCL binding in megakaryocytes identifies hematopoietic regulators. *Dev Cell*, 20(5):597-609.  
<https://doi.org/10.1016/j.devcel.2011.04.008>
- Tong W, Lodish HF, 2004. Lnk inhibits Tpo-mpl signaling and Tpo-mediated megakaryocytopoiesis. *J Exp Med*, 200(5):569-580.  
<https://doi.org/10.1084/jem.20040762>
- Trakala M, Partida D, Salazar-Roa M, et al., 2015a. Activation of the endomitotic spindle assembly checkpoint and thrombocytopenia in Plk1-deficient mice. *Blood*, 126(14):1707-1714.  
<https://doi.org/10.1182/blood-2015-03-634402>
- Trakala M, Rodríguez-Acebes S, Maroto M, et al., 2015b. Functional reprogramming of polyploidization in megakaryocytes. *Dev Cell*, 32(2):155-167.  
<https://doi.org/10.1016/j.devcel.2014.12.015>
- Uetake Y, Sluder G, 2004. Cell cycle progression after cleavage failure: mammalian somatic cells do not possess a "tetraploidy checkpoint". *J Cell Biol*, 165(5):609-615.  
<https://doi.org/10.1083/jcb.200403014>
- Vainchenker W, Raslova H, 2020. Megakaryocyte polyploidization: role in platelet production. *Platelets*, 31(6):707-716.  
<https://doi.org/10.1080/09537104.2019.1667497>
- Vainchenker W, Arkoun B, Basso-Valentina F, et al., 2021. Role of Rho-GTPases in megakaryopoiesis. *Small GTPases*, 12(5-6):399-415.  
<https://doi.org/10.1080/21541248.2021.1885134>
- van de Peer Y, Mizrahi E, Marchal K, 2017. The evolutionary significance of polyploidy. *Nat Rev Genet*, 18(7):411-424.  
<https://doi.org/10.1038/nrg.2017.26>
- van Rijnberk LM, Barrull-Mascaró R, van der Palen RL, et al., 2022. Endomitosis controls tissue-specific gene expression during development. *PLoS Biol*, 20(5):e3001597.  
<https://doi.org/10.1371/journal.pbio.3001597>
- Vilaboa N, Bermejo R, Martinez P, et al., 2004. A novel E2 box-GATA element modulates Cdc6 transcription during human cells polyploidization. *Nucleic Acids Res*, 32(21):6454-6467.  
<https://doi.org/10.1093/nar/gkh981>
- Vitale I, Galluzzi L, Castedo M, et al., 2011. Mitotic catastrophe: a mechanism for avoiding genomic instability. *Nat Rev Mol Cell Biol*, 12(6):385-392.  
<https://doi.org/10.1038/nrm3115>
- Vitrat N, Cohen-Solal K, Pique C, et al., 1998. Endomitosis of human megakaryocytes are due to abortive mitosis. *Blood*, 91(10):3711-3723.

- <https://doi.org/10.1182/blood.V91.10.3711>
- Wang MJ, Chen F, Lau JTY, et al., 2017. Hepatocyte polyploidization and its association with pathophysiological processes. *Cell Death Dis*, 8(5):e2805.  
<https://doi.org/10.1038/cddis.2017.167>
- Wang Q, Liu TY, Fang YQ, et al., 2004. *BUBR1* deficiency results in abnormal megakaryopoiesis. *Blood*, 103(4):1278-1285.  
<https://doi.org/10.1182/blood-2003-06-2158>
- Wang Z, Zhang Y, Kamen D, et al., 1995. Cyclin D3 is essential for megakaryocytopoiesis. *Blood*, 86(10):3783-3788.  
<https://doi.org/10.1182/blood.V86.10.3783.bloodjournal86103783>
- Weiss-Gayet M, Starck J, Chaabouni A, et al., 2016. Notch stimulates both self-renewal and lineage plasticity in a subset of murine CD9<sup>High</sup> committed megakaryocytic progenitors. *PLoS ONE*, 11(4):e0153860.  
<https://doi.org/10.1371/journal.pone.0153860>
- Wen Q, Leung C, Huang Z, et al., 2009. Survivin is not required for the endomitotic cell cycle of megakaryocytes. *Blood*, 114(1):153-156.  
<https://doi.org/10.1182/blood-2008-11-190801>
- Wen Q, Goldenson B, Silver SJ, et al., 2012. Identification of regulators of polyploidization presents therapeutic targets for treatment of AMKL. *Cell*, 150(3):575-589.  
<https://doi.org/10.1016/j.cell.2012.06.032>
- Wen QJ, Yang Q, Goldenson B, et al., 2015. Targeting megakaryocytic-induced fibrosis in myeloproliferative neoplasms by AURKA inhibition. *Nat Med*, 21(12):1473-1480.  
<https://doi.org/10.1038/nm.3995>
- Wilkinson PD, Delgado ER, Alencastro F, et al., 2019. The polyploid state restricts hepatocyte proliferation and liver regeneration in mice. *Hepatology*, 69(3):1242-1258.  
<https://doi.org/10.1002/hep.30286>
- Wong C, Stearns T, 2005. Mammalian cells lack checkpoints for tetraploidy, aberrant centrosome number, and cytokinesis failure. *BMC Cell Biol*, 6:6.  
<https://doi.org/10.1186/1471-2121-6-6>
- Wu H, Wade M, Krall L, et al., 1996. Targeted in vivo expression of the cyclin-dependent kinase inhibitor p21 halts hepatocyte cell-cycle progression, postnatal liver development and regeneration. *Genes Dev*, 10(3):245-260.  
<https://doi.org/10.1101/gad.10.3.245>
- Wurzenberger C, Gerlich DW, 2011. Phosphatases: providing safe passage through mitotic exit. *Nat Rev Mol Cell Biol*, 12(8):469-482.  
<https://doi.org/10.1038/nrm3149>
- Yu CJ, Yang Q, Chen YH, et al., 2016. Tyrosine 625 plays a key role and cooperates with tyrosine 630 in MPL W515L-induced signaling and myeloproliferative neoplasms. *Cell Biosci*, 6:34.  
<https://doi.org/10.1186/s13578-016-0097-3>
- Yu FX, Zhao B, Panupinthu N, et al., 2012. Regulation of the Hippo-YAP pathway by G-protein-coupled receptor signaling. *Cell*, 150(4):780-791.  
<https://doi.org/10.1016/j.cell.2012.06.037>
- Zack TI, Schumacher SE, Carter SL, et al., 2013. Pan-cancer patterns of somatic copy number alteration. *Nat Genet*, 45(10):1134-1140.  
<https://doi.org/10.1038/ng.2760>
- Zang CZ, Luyten A, Chen J, et al., 2016. NF-E2, FLI1 and RUNX1 collaborate at areas of dynamic chromatin to activate transcription in mature mouse megakaryocytes. *Sci Rep*, 6:30255.  
<https://doi.org/10.1038/srep30255>
- Zeng JK, Hills SA, Ozono E, et al., 2023. Cyclin E-induced replicative stress drives p53-dependent whole-genome duplication. *Cell*, 186(3):528-542.e14.  
<https://doi.org/10.1016/j.cell.2022.12.036>
- Zhang SH, Chen QH, Liu QX, et al., 2017. Hippo signaling suppresses cell ploidy and tumorigenesis through Skp2. *Cancer Cell*, 31(5):669-684.e7.  
<https://doi.org/10.1016/j.ccell.2017.04.004>
- Zhang SY, Nguyen LH, Zhou KJ, et al., 2018a. Knockdown of Anillin actin binding protein blocks cytokinesis in hepatocytes and reduces liver tumor development in mice without affecting regeneration. *Gastroenterology*, 154(5):1421-1434.  
<https://doi.org/10.1053/j.gastro.2017.12.013>
- Zhang SY, Zhou KJ, Luo X, et al., 2018b. The polyploid state plays a tumor-suppressive role in the liver. *Dev Cell*, 44(4):447-459.e5.  
<https://doi.org/10.1016/j.devcel.2018.01.010>
- Zhang SY, Lin YH, Tarlow B, et al., 2019. The origins and functions of hepatic polyploidy. *Cell Cycle*, 18(12):1302-1315.  
<https://doi.org/10.1080/15384101.2019.1618123>
- Zhang Y, Wang ZY, Ravid K, 1996. The cell cycle in polyploid megakaryocytes is associated with reduced activity of cyclin B1-dependent Cdc2 kinase. *J Biol Chem*, 271(8):4266-4272.  
<https://doi.org/10.1074/jbc.271.8.4266>
- Zhang Y, Sun S, Wang ZY, et al., 2002. Signaling by the Mpl receptor involves IKK and NF- $\kappa$ B. *J Cell Biochem*, 85(3):523-535.  
<https://doi.org/10.1002/jcb.10141>
- Zielke N, Edgar BA, DePamphilis ML, 2013. Endoreplication. *Cold Spring Harb Perspect Biol*, 5(1):a012948.  
<https://doi.org/10.1101/cshperspect.a012948>
- Zimmet J, Ravid K, 2000. Polyploidy: occurrence in nature, mechanisms, and significance for the megakaryocyte-platelet system. *Exp Hematol*, 28(1):3-16.  
[https://doi.org/10.1016/s0301-472x\(99\)00124-1](https://doi.org/10.1016/s0301-472x(99)00124-1)