



## Review

<https://doi.org/10.1631/jzus.B2500085>



# Cell proliferation and differentiation during epidermis renewal

Kai LIU<sup>1,2</sup>, Kaidi REN<sup>3</sup>, Dengwen LI<sup>2</sup>✉, Yang YANG<sup>1</sup>✉

<sup>1</sup>Department of Translational Medicine Center, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

<sup>2</sup>Department of Genetics and Cell Biology, State Key Laboratory of Medicinal Chemical Biology, College of Life Sciences, Nankai University, Tianjin 300071, China

<sup>3</sup>Department of Pharmacy, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

**Abstract:** Residing at the outermost layer of the skin, the epidermis is composed of stratified squamous epithelial cells. Regular renewal of the epidermis is essential for maintaining its barrier function, which is dependent on the orchestrated proliferation and differentiation of stem cells located in the basal epidermis. This process necessitates precise dual regulation through the intrinsic control of cell division orientation and external microenvironmental influences. In this comprehensive review, we delve into the critical processes underlying epidermis renewal, emphasizing the balance between symmetric and asymmetric cell fate and the integration of differentiated cells into the suprabasal layer. Our paper highlights the pivotal roles of single-cell omics, live imaging, and artificial intelligence (AI)-driven modeling techniques in elucidating the molecular mechanisms governing cell proliferation and differentiation during epidermis renewal.

**Key words:** Epidermis renewal; Stem cell proliferation; Asymmetric cell division; Basement membrane dynamics; Single-cell omics; Mechanotransduction

## 1 Introduction

The outermost layer of the skin is primarily composed of the interfollicular epidermis, which is the stratified epithelium located between pilosebaceous units (Banjac et al., 2023). The epidermis serves as a crucial interface, establishing a formidable barrier that separates the internal milieu of the body from the external environment and protects organisms from dehydration, physical trauma, chemical erosion, and pathogen invasion (Schmuth et al., 2024; Zhong et al., 2024). To comprehend the barrier function of the epidermis, it is essential to understand how it evolves from its initial state during development to its mature form. The epidermis initially develops from a monolayer of epithelial cells originating from the ectoderm

(Smart, 1970). During embryonic and postnatal development, stem cells produce two equally proliferative daughter cells through symmetric division to accommodate growth, a process that continues throughout life with a decreasing proliferation rate after birth (Dekoninck et al., 2020). Around embryonic day 9.5 (E9.5), upon induction by p63, progenitor cells begin to downregulate the expression of keratin-8 (K8) and upregulate the expression of K5 and K14 (Koster and Roop, 2004), giving rise to two cells with different fates by asymmetric division perpendicular to the basement membrane, one of which is committed to differentiation. The process of differentiation is irreversible and gradually leads to the formation of the stratum spinosum, stratum granulosum, and stratum corneum, establishing the epidermal barrier with tighter cell connections to withstand post-embryonic environmental conditions (Hardman et al., 1998).

In the human epidermis, it typically takes 14 to 18 d for a newly produced cell to transition into a cornified cell (Christophers and Laurence, 1976). This cycle of cell renewal is primarily controlled by stem cells located in the basal epidermis. However, in adulthood, most progenitor cells produce offspring through

✉ Yang YANG, [fccyangyang430@zju.edu.cn](mailto:fccyangyang430@zju.edu.cn)

Dengwen LI, [dwli@nankai.edu.cn](mailto:dwli@nankai.edu.cn)

Yang YANG, <https://orcid.org/0000-0001-7432-4735>

Dengwen LI, <https://orcid.org/0000-0002-4573-7191>

Kai LIU, <https://orcid.org/0000-0001-9811-1275>

Kaidi REN, <https://orcid.org/0000-0002-2011-6743>

Received Feb. 23, 2025; Revision accepted Apr. 22, 2025;  
Crosschecked Apr. 1, 2026; Published online Apr. 15, 2026

© Zhejiang University Press 2026

asymmetric division, half of which preserve the progenitor's proliferative potential while the other half, regulated by internal and external environmental factors, are committed to differentiation (Gola and Fuchs, 2021; Bansaccal et al., 2023; Eyermann et al., 2024; Roig-Rosello et al., 2024). The traditional view is that fast-cycling progenitor cells originate from slow-cycling stem cells in the epidermis (Mascré et al., 2012). However, recent studies have proposed an alternative perspective, suggesting that there are two independent cell populations with different cell cycle kinetics, both of which contain stem cells (Sada et al., 2016). In either case, this process of differentiation is irreversible as cells gradually migrate upward and replace shed stratum corneum cells, thereby maintaining the barrier function of the skin (Rompolas et al., 2016).

The disruption of balance between proliferation and differentiation can result in structural damage to the epidermis and the formation of tumors (Blanpain and Simons, 2013). A fundamental aspect of skin homeostasis is understanding how basal stem cells achieve dynamic equilibrium between symmetric divisions for proliferation and asymmetric divisions for differentiation. The mechanisms by which stem cell pools remain stable and how daughter cells differentiate to replace shed cells are critical but not fully understood aspects of epidermis renewal.


## 2 Development and structure of the epidermis

The stratified epidermis of the skin is derived from the monolayer epithelium that surrounds the embryonic ectoderm (Matsui, 2023; Holthaus and Eckhart, 2024). Prior to the establishment of a fully functional barrier, a transitional structure called periderm is formed. A subset of cuboidal ectodermal cells expresses K17; these cells detach from the basement membrane, and migrate over the basal layer to form a temporary barrier structure (Visscher et al., 2022; Jacob et al., 2023). Unlike the human single-layer periderm, the mouse periderm consists of two layers, an outer layer that is presumed to be absorptive and an inner layer packed with specific granules. The periderm disintegrates due to loss of contact with terminally-differentiated keratinocytes and is replaced by stratified epithelium (Panteleyev, 2022). At around E8.5, p63 expression initiates the process of epidermal stratification. During this phase, progenitor cells undergo the

downregulation of simple keratin intermediate filaments such as K8 and begin to express early differentiation markers K5 and K14 (Koster and Roop, 2004; Baba et al., 2020). At E13.5, epidermal stratification initiates with the emergence of intermediate cells expressing K1/K10. During this phase, robust proliferative activity is observed, and mitotic cells are detectable even within suprabasal layers. At E15.5, a fully differentiated stratum spinosum is established (Fuchs and Green, 1980; Liu et al., 2013; Damen et al., 2021). As epidermal development progresses, spinous cells undergo differentiation to become granular cells that express loricrin and pro-filaggrin at E16.5 (Mehrel et al., 1990; Joost et al., 2016). Ultimately, the formation of a rigid scaffold through the cross-linking of proteins and the covalent attachment of lipids on their outer surface results in the development of the cornified cell envelope, thereby establishing the barrier function of the skin at E18.5 (Liu et al., 2013; Rinnerthaler et al., 2015). Prior to birth, a fully developed epidermis comprising the stratum basale, stratum spinosum, stratum granulosum, and stratum corneum has already been established (Yin et al., 2023) (Fig. 1). Notably, the current stratification method for epidermal layers relies primarily on intermediate filament protein profiling. However, under specific pathological conditions, particularly in cases of abnormal cellular thickening, it remains challenging to discern whether this phenomenon results from the hyperproliferation of basal cells that mechanically displace a subset of cells into the suprabasal compartment or from microenvironmental alterations in the suprabasal niche that enable vertically divided daughter cells (presumably destined for differentiation) to retain partial basal progenitor properties, as evidenced by persistent K5/K14 immunoreactivity. This diagnostic ambiguity necessitates the integration of complementary detection modalities to accurately characterize aberrant cell populations. Such multidimensional analysis could significantly enhance diagnostic specificity and therapeutic targeting in epidermal pathologies.

## 3 Contribution of the basement membrane to epidermis renewal

The basement membrane is a highly organized extracellular matrix network located at the epithelial–mesenchymal interface of tissue. It comprises two




**Fig. 1** Development and structure of the epidermis. At embryonic day 9.5 (E9.5), the periderm cells detach from the basement membrane to form a temporary barrier structure, and are eventually shed alongside terminally differentiated corneocytes. By E13.5, a subset of basal progenitor cells undergoes planar-to-apical division reorientation, initiating terminal differentiation programs that drive the emergence of keratin-1 (K1)/K10-positive intermediate cells within suprabasal compartments. The formation of the stratum spinosum occurs at E15.5, followed by the stratum granulosum at E16.5, and finally the stratum corneum at E18.5.

independent networks of polymerized laminin and cross-linked collagens (Piperigkou et al., 2018), providing structural support for cells and influencing their fate (Bergman et al., 1992; Koster and Roop, 2007). Between the epidermis and dermis, the basement membrane provides adhesion and controls cell fate via integrin-mediated focal adhesions and hemidesmosomes (Rousselle et al., 2022). The integrin- and laminin-dependent contact on the basement membrane during cell division appears to contribute to polarity formation and influence the proliferation/differentiation decisions of basal keratinocytes (Haake et al., 2024; Villeneuve et al., 2024). Integrins are thought to cross-talk with receptors such as epidermal growth factor receptor (EGFR), triggering the signaling cascade of mitogen-activated protein kinase (MAPK), Janus kinase/signal transducer and activator of transcription (JAK/STAT), and phosphatidylinositol-3 kinase (PI3K), which in turn promotes the maintenance of the stem cell population in basal cells (Simpson et al., 2011; te Molder et al., 2021; Arnaud et al., 2023). With aging, the loss of hemidesmosomes from the basement membrane triggers mechanical signals that alter its structure and reduce the anchoring of basal keratinocytes—one of the main reasons that ultimately forces them to differentiate (Khalilgharibi and Mao, 2021; Roig-Rosello et al., 2024). The basement membrane

contains a variety of growth factors, morphogens, and other regulatory macromolecules that play a role in orchestrating the various processes necessary for the proper functioning of keratinocytes (Pozzi et al., 2017; Randles et al., 2017). Additionally, as the epidermis extends into the dermis and embeds between dermal papillae, the unique structure of the basement membrane reinforces connectivity between the dermis and epidermis, imposing mechanical constraints on keratinocytes that may affect their fate (Lawlor and Kaur, 2015) (Fig. 2).

The basement membrane, functioning as a biomechanical scaffold that preserves tissue architecture and transduces essential mechanotransduction cues, simultaneously orchestrates stem cell differentiation and maintains tissue-specific progenitor niches through the spatial presentation of growth factors (e.g., fibroblast growth factors (FGFs) and transforming growth factor- $\beta$  (TGF- $\beta$ )). This dual regulatory network endows the basement membrane with spatiotemporal regulatory precision, enabling robust defense against homeostatic imbalances caused by extrinsic environmental perturbations or intrinsic cellular dysregulation. Advances in understanding the composition of basement membrane and its roles in epidermis renewal are driving innovations in wound healing and disease treatment. In recent years, several studies in



**Fig. 2** Contribution of the basement membrane to epidermis renewal. The basement membrane is a highly organized extracellular matrix network located at the epithelial–mesenchymal interface of tissue, which comprises two independent networks of polymerized laminins and crosslinked collagens. Hemidesmosomes connect to the basement membrane through  $\alpha 6 \beta 4$  integrins and bullous pemphigoid antigen 180 (BP180) and are tethered to intermediate filaments by plectin and BP230. Focal adhesions are mediated through the interaction between  $\alpha 3 \beta 1$  integrin and laminin 511, and are linked to the actin cytoskeleton through Talin and Kindlin. Integrins are thought to crosstalk with receptors such as epidermal growth factor receptor (EGFR) to induce the proliferation of cells via the signaling cascade of mitogen-activated protein kinase (MAPK) and phosphatidylinositol-3 kinase (PI3K). As cells undergo stratification, the decreased density of integrins allows for differentiation. JAK/STAT: Janus kinase/signal transducer and activator of transcription.

this field have been conducted. For example, the topical application of recombinant type VII collagen promoted wound healing (Wang et al., 2013); the development of decellularized matrix-based bioinks incorporating basement membrane components (collagen IV and laminin) enhanced artificial skin mechanics and cytocompatibility (Kim et al., 2020); and base-editing technologies correcting basement membrane-associated genes in progeria models recovered regenerative capacity (Koblan et al., 2021). However, epidermis renewal involves complex basement membrane dynamics that serve not only as a physical scaffold but also as an information processor directing stem cell fate. While tissue engineering has progressed from component replication to dynamic biomimicry, cross-scale regulation (molecular-cellular-tissue) remains a challenging task. Programmable basement

membrane reconstruction will be enabled by the future integration of synthetic biology, single-cell mechanophenotyping (e.g., atomic force microscopy), and artificial intelligence (AI)-driven scaffold design.

#### 4 Cell division during epidermis renewal

The skin epidermis generates approximately  $10^7$  new cells every day to replace continuously shed stratum corneum cells (Banjac et al., 2023). The core of maintaining a stem cell pool lies in balancing self-renewal and differentiation to generate a sufficient number of specialized cells to execute basic tissue functions. To accommodate rapid growth during embryonic development, epidermal basal cells need to undergo symmetric divisions to generate a large number

of daughter cells with equivalent proliferative potential. Even from E13.5 to E15.5, suprabasal layer cells can proliferate symmetrically to expand the cell population (Damen et al., 2021). From this point onward, certain cells are stimulated by signals to undergo asymmetric divisions in order to produce differentiated cells (Lechler and Fuchs, 2005; Williams et al., 2011). There is also evidence indicating that some basal cells participate in epidermis renewal by directly delaminating from the basement membrane rather than undergoing division (Damen et al., 2021). Upon reaching adulthood, it is no longer necessary for the epidermis to have a rapid proliferation rate but instead it transitions toward a balance that maintains the number of cells with proliferative potential and those destined for differentiation (Morrison and Kimble, 2006; Gola and Fuchs, 2021). Cell division plays a critical role in determining the fate of hair follicles and interfollicle epidermal stem cells (Niessen et al., 2013), as well as in regulating hair follicle formation and epidermis hyperproliferation (Ouspenskaia et al., 2016; Seldin et al., 2016). The pattern of cell division is closely associated with the orientation of the mitotic spindle apparatus (Bergstralh et al., 2017; Seldin and Macara, 2017). It is important to note that “oriented cell division” does not necessarily mean “asymmetric” or “symmetric” cell division, with the former referring to the alignment of directions along a particular axis during division and the latter referring to whether the distribution of materials within the cell and the fate of the daughter cells are equal (Williams and Fuchs, 2013). Even when cells divide symmetrically, the daughter cells may have different fates under different environmental stimuli. In the epidermis, cell divisions oriented perpendicular to the basement membrane typically result in asymmetric cell fate allocation under homeostatic conditions. This outcome is determined by both the heterogeneous distribution of intrinsic signaling molecules between daughter cells and the biomechanical heterogeneity of the microenvironment, as discussed in detail below. Epidermal stem cells and progenitors regulate their number and stemness during homeostasis in two main ways: (1) In wound healing, most daughter cells from symmetrical divisions parallel to the basal membrane still maintain their proliferation ability due to basement membrane contact, increasing the stem cell pool; (2) During normal renewal, one daughter cell from asymmetric divisions


perpendicular to the basal membrane remains adherent, maintaining stem cell numbers. There is a “self-renewal checkpoint” that restricts the self-renewal of stem cells in response to genotoxic stress (Matsumura et al., 2016). It has been shown that stress-induced collagen type XVII  $\alpha 1$  chain (COL17A1) proteolysis plays a role in mediating the checkpoint function of stem cells, leading to the natural selection of eligible stem cells for organ homeostasis over their lifespan (Liu et al., 2019). Furthermore, self-renewal can also be promoted by neighboring differentiation events (Mesa et al., 2018). Aberrant proliferation kinetics in basal layer cells, such as excessive mitotic activity, may lead to pathological hyperplasia or carcinogenesis, while insufficient proliferation could compromise epidermal barrier integrity. Maintaining this homeostatic equilibrium requires a sophisticated regulatory network, the elucidation of which holds critical implications for understanding disease pathogenesis and developing therapeutic interventions.

## 5 Spindle orientation during cell division

In the basal epidermis, approximately 60%–80% of the cells undergo asymmetric division to produce one basal cell and one differentiated cell (Rompolas et al., 2016). The polarity of asymmetric division in epidermal basal cells is determined by both external and internal factors. External factors, such as  $\beta 1$  integrins,  $\beta 4$  integrins, and the  $\beta 1$ -integrin-binding protein Kindlin-1 on the basement membrane, play significant roles in cell division polarity (Weaver et al., 2002; Ussar et al., 2006). The intrinsic molecular mechanism of asymmetric division has been highly conserved during evolution (Betschinger and Knoblich, 2004). It has been observed that the basal cells do not establish a division axis before mitosis, and the spindles are randomly distributed even in the prophase until the direction of division is finally determined by the movement of centrosomes in the prometaphase and late metaphase (Poulson and Lechler, 2010). Spindle orientation involves two steps mediated by the apical polarity complex and the spindle position complex. During junctional maturation, Merlin associates directly with  $\alpha$ -catenin and links it to partitioning defective 3 (Par3), thereby providing an essential link between the adhesion junction and Par3. It is also possible that

Par3 localizes to the apical cortex in the same way as it does during asymmetric division (Gladden et al., 2010). The apical polarity complex composed of Par3, atypical protein kinase C (aPKC), and Par6 locates on the cortical crescents and then recruits the leucine-glycine-asparagine (Leu-Gly-Asn) repeat-enriched protein (LGN)-G $\alpha_{13}$ -nuclear mitotic apparatus protein (NuMA)-dynein-dynactin complex to the apical cortex through Inscuteable (Insc) (Poulson and Lechler, 2010; Williams et al., 2011; Niessen et al., 2013; Ali et al., 2016; Seldin et al., 2016; Ma et al., 2022; Yan et al., 2022). Cell division cycle protein 42 (CDC42) may also play a role in this process (Prado-Mantilla and Lechler, 2023). In *Drosophila*, neuroblasts lacking *Insc* fail to properly orient their spindle position complex. Conversely, in the neuroectoderm, the over-expression of *Insc* can reorient the spindle (Kraut et al., 1996), demonstrating that *Insc* is both necessary and sufficient for the initiation of asymmetric division. This notion is further supported by the upregulation

of *Insc* at E13.5 coinciding with the stratification of the epidermis (Poulson and Lechler, 2010). During mitosis, Insc assists in recruiting LGN to the cell cortex through glycosylphosphatidylinositol (GPI)-linked G $\alpha_{13}$ , which binds LGN C-terminal GoLoco motifs. The stability of this complex structure is enhanced when all three components are present (Williams et al., 2011). Recent structural studies on LGN indicate its inability to bind both Insc and NuMA simultaneously, with a higher affinity for Insc than for NuMA (Zhu et al., 2011). This discovery is consistent with the model proposing that Par3-Insc and G $\alpha_{13}$  facilitate the apical recruitment of LGN during early mitosis, thereby promoting its transfer to NuMA. LGN-NuMA plays a crucial role in redirecting the mitotic spindle (Williams et al., 2014), and a variety of proteins collaborate to finalize the process of mitosis (Zhong et al., 2022; Yang et al., 2023; Yu et al., 2023) (Fig. 3). Interestingly, in *K14-Cre<sup>+</sup>*, *Sas-4<sup>fl</sup>*, and *p53<sup>fl</sup>* conditional double-knockout mice, the proliferation and differentiation of



**Fig. 3 Spindle orientation and cell differentiation commitment.** During prometaphase, the apical polarity complex composed of atypical protein kinase C (aPKC), partitioning defective 3 (Par3), and Par6 localizes on the cortical crescents. This may be facilitated by the interaction of this complex with  $\alpha$ -catenin via Merlin. Subsequently, the apical complex recruits the leucine-glycine-asparagine (Leu-Gly-Asn) repeat-enriched protein (LGN)-G $\alpha_{13}$ -nuclear mitotic apparatus protein (NuMA)-dynein-dynactin complex to the apical cortex, leading to spindle rotation. The  $\beta$ 1 integrins,  $\beta$ 4 integrins, and  $\beta$ 1-integrin-binding protein Kindlin-1 located on the basement membrane also play vital roles in establishing and maintaining cell division polarity. After the establishment of cell division polarity, certain signaling molecules, such as Notch and epidermal growth factor receptor (EGFR), become asymmetrically distributed, thereby determining the fate of daughter cells. Insc: Inscuteable.

progenitor cells in the late epidermal development stage are not coordinated with the direction of division. This suggests that, aside from spindle orientation, there may be other mechanisms regulating cell differentiation at this stage (Damen et al., 2021). When stimulated by internal and external signals, cells begin to differentiate and replace the shed cells through asymmetric division.

Despite that our knowledge of cell division in the epidermis has significantly improved in recent years, the mechanisms by which cells determine their specific division orientation with spatiotemporal precision remain unclear. Elucidating the molecular basis of this process holds significant value in understanding epidermal development and homeostasis maintenance. As mentioned above, a critical question persists: does the basement membrane dictate stem cell division patterns by mechanical signaling changes that subsequently influence daughter cell fate, or do stem cells sense the local cellular density and autonomously regulate their division orientation and utilize the basement membrane to determine progeny cell destiny? Determining which component plays the predominant regulatory role in this process, or whether it results from coordinated interactions between both elements, crucially requires clarifying the underlying molecular mechanisms; resolving these questions can substantially improve our comprehension of epithelial tissue dynamics.

## 6 Cellular lineage and cell differentiation commitment

The epidermis at different locations may exhibit distinct patterns of cell lineages. Classical models of adult epidermal homeostasis suggest that cells with infrequent division are long-lived stem cells that give rise to rapidly dividing short-lived progenitor cells (Clayton et al., 2007; Sada and Tumber, 2013). Consistent with this model, the adult mouse skin epidermis has two types of cells with proliferative potential: slow-cycling stem cells and more rapidly cycling committed progenitor cells, each performing different functions during homeostasis and interfollicular epidermis repair (Shin and Peterson, 2013; Aragona et al., 2020; Lotti et al., 2022). The basal layer of the mouse ear and paw epidermis comprises a single equipotent stem cell population, where daughter cells are generated

with either an independent fate in the ear or a positively correlated fate in the paw, indicating non-hierarchical or asymmetric division (Rompolas et al., 2016). However, in the mouse tail epidermis, these two populations, identified by the two markers distal-less homeobox 1 (*Dlx1*) and solute carrier family 1 member 3 (*Slc1a3*), respectively, inhabit distinct territories and exhibit different rates of proliferation and upward cellular transport (Sada et al., 2016). Another study indicated that, in the mouse tail epidermis, cells of the basal layer house a distinct slow-cycling epidermal stem cell population distinguished by thymocyte cell surface antigen 1 (*Thy1*) expression, which coexists with transient amplifying progenitors (Koren et al., 2022). This dual population theory appears to be more appropriate than the single stem cell theory for explaining the differential proliferation rate within the scale and interscale during adult mouse tail epidermis homeostasis (Dekoninck et al., 2020).

Although epidermal stem cells display regional heterogeneity in molecular signatures and subpopulation compositions across anatomical sites, their progeny differentiation processes often share common features. Cell fate asymmetry originates from mitotic polarity establishment in the mother cell during cytokinesis, driving the asymmetric segregation of fate-specific signaling complexes between daughter cells. Daughter cells containing EGFR exhibit increased cycling activity (Xu et al., 2024), and the overexpression of EGFR in the basal layer contributes to epidermal hyperproliferation and delays differentiation (Arnaud et al., 2023). In contrast, Notch signaling has an opposing function to that of EGFR. Notch represses proliferation and induces terminal differentiation (Zhou et al., 2022). EGFR overrides Notch at the basal layer, inhibiting Notch activity. In the suprabasal layer where EGFR is absent, Notch promotes differentiation (Poulson and Lechler, 2012) (Fig. 3). Nevertheless, once daughter cells commit to differentiation from stem cells or progenitor cells, there is a general consensus on the model for how cells migrate from the basal layer of the epidermis to the outermost layer. The epidermal differentiation unit (EDU), a three-dimensional (3D) organization of columns modified by the epidermal proliferation unit (Ghazizadeh and Taichman, 2001), is defined by the perimeter of the most external terminally differentiated cells. Approximately 90% of the cells in the unit align into a vertical column and move upward to recycle the same space occupied by preceding cells,


while a minority of cells inhabiting the basal and spinous layers can switch to neighboring EDUs. Additionally, the inhibition of Wnt signaling promotes differentiation in interfollicular epidermis and sebaceous gland lineages (Satta et al., 2024). When cells arrive in the granular layer, their fate becomes vertically fixed. Following departure from the basal layer, the granular layer serves as a buffer zone, spatially coordinating the transition to cornified tissue (Rompolas et al., 2016) (Fig. 4).

The evolution of epidermis renewal models—from the classical unipotent stem-progenitor paradigm to the recently proposed dual stem cell paradigm—has significantly advanced our understanding of epidermal homeostasis. However, this progression raises the critical questions of (1) whether there exists a definitive demarcation between stem and progenitor cells and whether distinct biomarkers or functional criteria could reliably discriminate these populations; (2) beyond the tail epidermis demonstrating dual stem cell populations, whether other regions harbor heterogeneous stem cell subpopulations whose identification has been hindered by technical limitations in single-cell omics resolution; and (3) given that the murine

tail's developmental expansion requires orthogonal growth axes maintained by discrete stem cell subsets, whether analogous functional heterogeneity emerges during human morphogenesis to coordinate differential growth rates across body regions. These considerations suggest an intriguing research frontier in developmental biology.

## 7 Conclusions

The skin's ability to maintain structural integrity through continuous epidermis renewal involves complex interactions between basal stem cells, microenvironmental cues, and molecular signaling networks. While recent advances have elucidated the mechanisms governing stem cell division patterns and basement membrane dynamics, key questions remain regarding the spatiotemporal regulation of cellular heterogeneity and fate determination. Future research should prioritize mapping functional stem cell subpopulations across anatomical regions using spatial multi-omics, particularly the investigation of how mechanical forces and matrix remodeling influence



**Fig. 4 Cellular lineage and cell differentiation commitment.** (a) The cellular lineage hierarchy consists of long-lived stem cells that give rise to short-lived progenitors, which in turn generate differentiated cells via asymmetric division. (b) Utilizing the green fluorescent protein-tagged histone 2B (H2B-GFP) pulse-chase system, two territories (low copy repeat (LCR) and non-LCR) have been identified in the mouse tail epidermis. These territories are composed of distinct stem cells exhibiting different patterns of proliferation, differentiation, and upward cellular transport. It is currently unknown whether these stem cells have the capacity to give rise to progenitor cells. (c) Basal stem cells stochastically commit to differentiation and transition through the suprabasal layers predominantly via existing epidermal differentiation units. A small percentage of cells switch to another unit to complete this process.

division axis orientation during aging or injury. Concurrently, deciphering feedback mechanisms that adjust symmetric/asymmetric division ratios via mechanosensitive pathways (e.g., integrin–EGFR crosstalk) could revolutionize therapeutic strategies. Clinically, these insights enable the application of targeted approaches: modulating Notch signaling for psoriatic hyperplasia, bioengineering laminin-rich scaffolds for chronic wounds, or preserving COL17A1-mediated stem cell competition in aging skin. Technological integration proves that live imaging of 3D organoids by AI modeling can predict division outcomes under varying biomechanical stresses, while base editing offers the precision correction of basement membrane gene defects. Emerging synthetic matrices with tunable stiffness gradients may recreate niche-specific signals for in vitro epidermal expansion. Ultimately, converging single-cell resolution datasets with dynamic micro-environmental profiling will be able to unravel the four-dimensional control of epidermal homeostasis, bridging molecular mechanisms with clinical translation. This system-level understanding promises to transform regenerative dermatology through personalized interventions that restore tissue architecture in genetic disorders, chronic ulcers, and degenerative conditions, while informing broader paradigms of epithelial tissue engineering.

### Acknowledgments

This work was supported by the National Key R&D Program of China (No. 2021YFA1101001) and the National Natural Science Foundation of China (Nos. 82204389 and 82470403).

### Author contributions

Kai LIU and Dengwen LI conceptualized the review and wrote the original draft. Kai LIU created the figures. Dengwen LI and Yang YANG supervised the project and provided critical revisions. Yang YANG acquired funding, provided administrative support, and edited the manuscript. Kaidi REN contributed to the literature search and manuscript writing. All the authors have read and approved the final version of the manuscript.

### Compliance with ethics guidelines

Kai LIU, Kaidi REN, Dengwen LI, and Yang YANG declare that they have no conflicts of interest.

This review does not contain any studies with human or animal subjects performed by any of the authors.

### Declaration on the use of generative AI tools

During the preparation of this work, the authors used ChatGPT in order to check for grammatical errors. After using

this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

### References

- Ali NJA, Dias Gomes M, Bauer R, et al., 2016. Essential role of polarity protein Par3 for epidermal homeostasis through regulation of barrier function, keratinocyte differentiation, and stem cell maintenance. *J Invest Dermatol*, 136(12): 2406-2416.  
<https://doi.org/10.1016/j.jid.2016.07.011>
- Aragona M, Sifrim A, Malfait M, et al., 2020. Mechanisms of stretch-mediated skin expansion at single-cell resolution. *Nature*, 584(7820):268-273.  
<https://doi.org/10.1038/s41586-020-2555-7>
- Arnaud T, Rodrigues-Lima F, Viguier M, et al., 2023. Interplay between EGFR, E-cadherin, and PTP1B in epidermal homeostasis. *Tissue Barriers*, 11(3):2104085.  
<https://doi.org/10.1080/21688370.2022.2104085>
- Baba K, Sasaki K, Morita M, et al., 2020. Cell jamming, stratification and p63 expression in cultivated human corneal epithelial cell sheets. *Sci Rep*, 10:9282.  
<https://doi.org/10.1038/s41598-020-64394-6>
- Banjac I, Maimets M, Jensen KB, 2023. Maintenance of high-turnover tissues during and beyond homeostasis. *Cell Stem Cell*, 30(4):348-361.  
<https://doi.org/10.1016/j.stem.2023.03.008>
- Bansaccal N, Vieugue P, Sarate R, et al., 2023. The extracellular matrix dictates regional competence for tumour initiation. *Nature*, 623(7988):828-835.  
<https://doi.org/10.1038/s41586-023-06740-y>
- Bergman KA, Meis JF, Horrevorts AM, et al., 1992. Acute renal failure in a neonate due to pelviureteric candidal bezoars successfully treated with long-term systemic fluconazole. *Acta Paediatr*, 81(9):709-711.  
<https://doi.org/10.1111/j.1651-2227.1992.tb12342.x>
- Bergstrahl DT, Dawney NS, St Johnston D, 2017. Spindle orientation: a question of complex positioning. *Development*, 144(7):1137-1145.  
<https://doi.org/10.1242/dev.140764>
- Betschinger J, Knoblich JA, 2004. Dare to be different: asymmetric cell division in *Drosophila*, *C. elegans* and vertebrates. *Curr Biol*, 14(16):R674-R685.  
<https://doi.org/10.1016/j.cub.2004.08.017>
- Blanpain C, Simons BD, 2013. Unravelling stem cell dynamics by lineage tracing. *Nat Rev Mol Cell Biol*, 14(8): 489-502.  
<https://doi.org/10.1038/nrm3625>
- Christophers E, Laurence EB, 1976. Kinetics and structural organisation of the epidermis. *Curr Probl Dermatol*, 6: 87-106.  
<https://doi.org/10.1159/000399007>
- Clayton E, Doupé DP, Klein AM, et al., 2007. A single type of progenitor cell maintains normal epidermis. *Nature*, 446(7132):185-189.  
<https://doi.org/10.1038/nature05574>
- Damen M, Wirtz L, Soroka E, et al., 2021. High proliferation and delamination during skin epidermal stratification. *Nat Commun*, 12:3227.

- <https://doi.org/10.1038/s41467-021-23386-4>
- Dekoninck S, Hannezo E, Sifrim A, et al., 2020. Defining the design principles of skin epidermis postnatal growth. *Cell*, 181(3):604-620.e22.  
<https://doi.org/10.1016/j.cell.2020.03.015>
- Eyermann CE, Chen X, Somuncu OS, et al., 2024.  $\Delta$ Np63 regulates homeostasis, stemness, and suppression of inflammation in the adult epidermis. *J Invest Dermatol*, 144(1):73-83.e10.  
<https://doi.org/10.1016/j.jid.2023.07.005>
- Fuchs E, Green H, 1980. Changes in keratin gene expression during terminal differentiation of the keratinocyte. *Cell*, 19(4):1033-1042.  
[https://doi.org/10.1016/0092-8674\(80\)90094-x](https://doi.org/10.1016/0092-8674(80)90094-x)
- Ghazizadeh S, Taichman LB, 2001. Multiple classes of stem cells in cutaneous epithelium: a lineage analysis of adult mouse skin. *EMBO J*, 20(6):1215-1222.  
<https://doi.org/10.1093/emboj/20.6.1215>
- Gladden AB, Hebert AM, Schneeberger EE, et al., 2010. The NF2 tumor suppressor, Merlin, regulates epidermal development through the establishment of a junctional polarity complex. *Dev Cell*, 19(5):727-739.  
<https://doi.org/10.1016/j.devcel.2010.10.008>
- Gola A, Fuchs E, 2021. Environmental control of lineage plasticity and stem cell memory. *Curr Opin Cell Biol*, 69: 88-95.  
<https://doi.org/10.1016/j.ccb.2020.12.015>
- Haake SM, Rios BL, Pozzi A, et al., 2024. Integrating integrins with the hallmarks of cancer. *Matrix Biol*, 130:20-35.  
<https://doi.org/10.1016/j.matbio.2024.04.003>
- Hardman MJ, Sisi P, Banbury DN, et al., 1998. Patterned acquisition of skin barrier function during development. *Development*, 125(8):1541-1552.  
<https://doi.org/10.1242/dev.125.8.1541>
- Holthaus KB, Eckhart L, 2024. Development-associated genes of the epidermal differentiation complex (EDC). *J Dev Biol*, 12(1):4.  
<https://doi.org/10.3390/jdb12010004>
- Jacob T, Annusver K, Czarnewski P, et al., 2023. Molecular and spatial landmarks of early mouse skin development. *Dev Cell*, 58(20):2140-2162.e5.  
<https://doi.org/10.1016/j.devcel.2023.07.015>
- Joost S, Zeisel A, Jacob T, et al., 2016. Single-cell transcriptomics reveals that differentiation and spatial signatures shape epidermal and hair follicle heterogeneity. *Cell Syst*, 3(3):221-237.e9.  
<https://doi.org/10.1016/j.cels.2016.08.010>
- Khalilgharibi N, Mao YL, 2021. To form and function: on the role of basement membrane mechanics in tissue development, homeostasis and disease. *Open Biol*, 11(2):200360.  
<https://doi.org/10.1098/rsob.200360>
- Kim MK, Jeong W, Lee SM, et al., 2020. Decellularized extracellular matrix-based bio-ink with enhanced 3D printability and mechanical properties. *Biofabrication*, 12(2): 025003.  
<https://doi.org/10.1088/1758-5090/ab5d80>
- Koblan LW, Erdos MR, Wilson C, et al., 2021. In vivo base editing rescues Hutchinson-Gilford progeria syndrome in mice. *Nature*, 589(7843):608-614.  
<https://doi.org/10.1038/s41586-020-03086-7>
- Koren E, Feldman A, Yusupova M, et al., 2022. Thy1 marks a distinct population of slow-cycling stem cells in the mouse epidermis. *Nat Commun*, 13:4628.  
<https://doi.org/10.1038/s41467-022-31629-1>
- Koster MI, Roop DR, 2004. The role of p63 in development and differentiation of the epidermis: Tanioku Kihei memorial lecture. *J Dermatol Sci*, 34(1):3-9.  
<https://doi.org/10.1016/j.jdermsci.2003.10.003>
- Koster MI, Roop DR, 2007. Mechanisms regulating epithelial stratification. *Annu Rev Cell Dev Biol*, 23(1):93-113.  
<https://doi.org/10.1146/annurev.cellbio.23.090506.123357>
- Kraut R, Chia W, Jan LY, et al., 1996. Role of *inscuteable* in orienting asymmetric cell divisions in *Drosophila*. *Nature*, 383(6595):50-55.  
<https://doi.org/10.1038/383050a0>
- Lawlor KT, Kaur P, 2015. Dermal contributions to human interfollicular epidermal architecture and self-renewal. *Int J Mol Sci*, 16(12):28098-28107.  
<https://doi.org/10.3390/ijms161226078>
- Lechler T, Fuchs E, 2005. Asymmetric cell divisions promote stratification and differentiation of mammalian skin. *Nature*, 437(7056):275-280.  
<https://doi.org/10.1038/nature03922>
- Liu N, Matsumura H, Kato T, et al., 2019. Stem cell competition orchestrates skin homeostasis and ageing. *Nature*, 568(7752):344-350.  
<https://doi.org/10.1038/s41586-019-1085-7>
- Liu S, Zhang HS, Duan EK, 2013. Epidermal development in mammals: key regulators, signals from beneath, and stem cells. *Int J Mol Sci*, 14(6):10869-10895.  
<https://doi.org/10.3390/ijms140610869>
- Lotti R, Palazzo E, Quadri M, et al., 2022. Isolation of an “early” transit amplifying keratinocyte population in human epidermis: a role for the low affinity neurotrophin receptor CD271. *Stem Cells*, 40(12):1149-1161.  
<https://doi.org/10.1093/stmcls/sxac060>
- Ma HX, Qi FF, Ji L, et al., 2022. NuMA forms condensates through phase separation to drive spindle pole assembly. *J Mol Cell Biol*, 14(1):mjab081.  
<https://doi.org/10.1093/jmcb/mjab081>
- Mascre G, Dekoninck S, Drogat B, et al., 2012. Distinct contribution of stem and progenitor cells to epidermal maintenance. *Nature*, 489(7415):257-262.  
<https://doi.org/10.1038/nature11393>
- Matsui T, 2023. Epidermal barrier development via corneoptosis: a unique form of cell death in stratum granulosum cells. *J Dev Biol*, 11(4):43.  
<https://doi.org/10.3390/jdb11040043>
- Matsumura H, Mohri Y, Binh NT, et al., 2016. Hair follicle aging is driven by transepidermal elimination of stem cells via COL17A1 proteolysis. *Science*, 351(6273):aad4395.  
<https://doi.org/10.1126/science.aad4395>
- Mehrel T, Hohl D, Rothnagel JA, et al., 1990. Identification of a major keratinocyte cell envelope protein, loricrin. *Cell*, 61(6):1103-1112.  
[https://doi.org/10.1016/0092-8674\(90\)90073-n](https://doi.org/10.1016/0092-8674(90)90073-n)

- Mesa KR, Kawaguchi K, Cockburn K, et al., 2018. Homeostatic epidermal stem cell self-renewal is driven by local differentiation. *Cell Stem Cell*, 23(5):677-686.e4. <https://doi.org/10.1016/j.stem.2018.09.005>
- Morrison SJ, Kimble J, 2006. Asymmetric and symmetric stem-cell divisions in development and cancer. *Nature*, 441(7097):1068-1074. <https://doi.org/10.1038/nature04956>
- Niessen MT, Scott J, Zielinski JG, et al., 2013. aPKC $\lambda$  controls epidermal homeostasis and stem cell fate through regulation of division orientation. *J Cell Biol*, 202(6):887-900. <https://doi.org/10.1083/jcb.201307001>
- Ouspenskaia T, Matos I, Mertz AF, et al., 2016. WNT-SHH antagonism specifies and expands stem cells prior to niche formation. *Cell*, 164(1-2):156-169. <https://doi.org/10.1016/j.cell.2015.11.058>
- Panteleyev AA, 2022. Dual nature of mouse periderm structure, function and fate. *Exp Dermatol*, 31(7):1102-1108. <https://doi.org/10.1111/exd.14608>
- Piperigkou Z, Götte M, Theocharis AD, et al., 2018. Insights into the key roles of epigenetics in matrix macromolecules-associated wound healing. *Adv Drug Deliv Rev*, 129:16-36. <https://doi.org/10.1016/j.addr.2017.10.008>
- Poulson ND, Lechler T, 2010. Robust control of mitotic spindle orientation in the developing epidermis. *J Cell Biol*, 191(5):915-922. <https://doi.org/10.1083/jcb.201008001>
- Poulson ND, Lechler T, 2012. Asymmetric cell divisions in the epidermis. *Int Rev Cell Mol Biol*, 295:199-232. <https://doi.org/10.1016/B978-0-12-394306-4.00012-5>
- Pozzi A, Yurchenco PD, Iozzo RV, 2017. The nature and biology of basement membranes. *Matrix Biol*, 57-58:1-11. <https://doi.org/10.1016/j.matbio.2016.12.009>
- Prado-Mantilla A, Lechler T, 2023. Polarity in skin development and cancer. *Curr Top Dev Biol*, 154:317-336. <https://doi.org/10.1016/bs.ctdb.2023.02.003>
- Randles MJ, Humphries MJ, Lennon R, 2017. Proteomic definitions of basement membrane composition in health and disease. *Matrix Biol*, 57-58:12-28. <https://doi.org/10.1016/j.matbio.2016.08.006>
- Rinnerthaler M, Bischof J, Streubel MK, et al., 2015. Oxidative stress in aging human skin. *Biomolecules*, 5(2):545-589. <https://doi.org/10.3390/biom5020545>
- Roig-Rosello E, Dayan G, Bovio S, et al., 2024. Dermal stiffness governs the topography of the epidermis and the underlying basement membrane in young and old human skin. *Aging Cell*, 23(4):e14096. <https://doi.org/10.1111/ace1.14096>
- Rompolas P, Mesa KR, Kawaguchi K, et al., 2016. Spatiotemporal coordination of stem cell commitment during epidermal homeostasis. *Science*, 352(6292):1471-1474. <https://doi.org/10.1126/science.aaf7012>
- Rousselle P, Laigle C, Rousselet G, 2022. The basement membrane in epidermal polarity, stemness, and regeneration. *Am J Physiol Cell Physiol*, 323(6):C1807-C1822. <https://doi.org/10.1152/ajpcell.00069.2022>
- Sada A, Tumber T, 2013. New insights into mechanisms of stem cell daughter fate determination in regenerative tissues. *Int Rev Cell Mol Biol*, 300:1-50. <https://doi.org/10.1016/B978-0-12-405210-9.00001-1>
- Sada A, Jacob F, Leung E, et al., 2016. Defining the cellular lineage hierarchy in the interfollicular epidermis of adult skin. *Nat Cell Biol*, 18(6):619-631. <https://doi.org/10.1038/ncb3359>
- Satta JP, Lan Q, Taketo MM, et al., 2024. Stabilization of epithelial  $\beta$ -catenin compromises mammary cell fate acquisition and branching morphogenesis. *J Invest Dermatol*, 144(6):1223-1237.e10. <https://doi.org/10.1016/j.jid.2023.11.018>
- Schmuth M, Eckmann S, Moosbrugger-Martinez V, et al., 2024. Skin barrier in atopic dermatitis. *J Invest Dermatol*, 144(5):989-1000.e1. <https://doi.org/10.1016/j.jid.2024.03.006>
- Seldin L, Macara I, 2017. Epithelial spindle orientation diversities and uncertainties: recent developments and lingering questions. *F1000Res*, 6:984. <https://doi.org/10.12688/f1000research.11370.1>
- Seldin L, Muroyama A, Lechler T, 2016. NuMA-microtubule interactions are critical for spindle orientation and the morphogenesis of diverse epidermal structures. *eLife*, 5:e12504. <https://doi.org/10.7554/eLife.12504>
- Shin L, Peterson DA, 2013. Human mesenchymal stem cell grafts enhance normal and impaired wound healing by recruiting existing endogenous tissue stem/progenitor cells. *Stem Cells Transl Med*, 2(1):33-42. <https://doi.org/10.5966/sctm.2012-0041>
- Simpson CL, Patel DM, Green KJ, 2011. Deconstructing the skin: cytoarchitectural determinants of epidermal morphogenesis. *Nat Rev Mol Cell Biol*, 12(9):565-580. <https://doi.org/10.1038/nrm3175>
- Smart IHM, 1970. Variation in the plane of cell cleavage during the process of stratification in the mouse epidermis. *Br J Dermatol*, 82(3):276-282. <https://doi.org/10.1111/j.1365-2133.1970.tb12437.x>
- te Molder L, de Pereda JM, Sonnenberg A, 2021. Regulation of hemidesmosome dynamics and cell signaling by integrin  $\alpha 6 \beta 4$ . *J Cell Sci*, 134(18):jcs259004. <https://doi.org/10.1242/jcs.259004>
- Ussar S, Wang HV, Linder S, et al., 2006. The Kindlins: subcellular localization and expression during murine development. *Exp Cell Res*, 312(16):3142-3151. <https://doi.org/10.1016/j.yexcr.2006.06.030>
- Villeneuve C, Hashmi A, Ylivinkka I, et al., 2024. Mechanical forces across compartments coordinate cell shape and fate transitions to generate tissue architecture. *Nat Cell Biol*, 26(2):207-218. <https://doi.org/10.1038/s41556-023-01332-4>
- Visscher MO, Carr AN, Narendran V, 2022. Epidermal immunity and function: origin in neonatal skin. *Front Mol Biosci*, 9:894496. <https://doi.org/10.3389/fmolb.2022.894496>
- Wang XY, Ghasri P, Amir M, et al., 2013. Topical application

- of recombinant type VII collagen incorporates into the dermal-epidermal junction and promotes wound closure. *Mol Ther*, 21(7):1335-1344.  
<https://doi.org/10.1038/mt.2013.87>
- Weaver VM, Lelièvre S, Lakins JN, et al., 2002.  $\beta$ 4 integrin-dependent formation of polarized three-dimensional architecture confers resistance to apoptosis in normal and malignant mammary epithelium. *Cancer Cell*, 2(3):205-216.  
[https://doi.org/10.1016/s1535-6108\(02\)00125-3](https://doi.org/10.1016/s1535-6108(02)00125-3)
- Williams SE, Fuchs E, 2013. Oriented divisions, fate decisions. *Curr Opin Cell Biol*, 25(6):749-758.  
<https://doi.org/10.1016/j.ccb.2013.08.003>
- Williams SE, Beronja S, Pasolli HA, et al., 2011. Asymmetric cell divisions promote Notch-dependent epidermal differentiation. *Nature*, 470(7334):353-358.  
<https://doi.org/10.1038/nature09793>
- Williams SE, Ratliff LA, Postiglione MP, et al., 2014. Par3-mInsc and  $G\alpha_3$  cooperate to promote oriented epidermal cell divisions through LGN. *Nat Cell Biol*, 16(8):758-769.  
<https://doi.org/10.1038/ncb3001>
- Xu P, Yang LL, Lai S, et al., 2024. Effects of EGFR-TKI on epidermal melanin unit integrity: therapeutic implications for hypopigmented skin disorders. *Pigment Cell Melanoma Res*, 37(4):514-529.  
<https://doi.org/10.1111/pcmr.13171>
- Yan JJ, Zhang K, Guo TM, et al., 2022. Par3 regulates the asymmetric division of basal stem cells in psoriasis via the Par3/mInsc/LGN signaling axis. *Cell Immunol*, 373:104496.  
<https://doi.org/10.1016/j.cellimm.2022.104496>
- Yang S, Luo YG, Yang ML, et al., 2023. Src inhibition induces mitotic arrest associated with chromosomal passenger complex. *Cell Tissue Res*, 392(3):733-743.  
<https://doi.org/10.1007/s00441-023-03765-7>
- Yin HX, Hu MZ, Li DW, 2023. Regulation of epidermal stratification and development by basal keratinocytes. *J Cell Physiol*, 238(4):742-748.  
<https://doi.org/10.1002/jcp.30978>
- Yu F, Yang S, Ni H, et al., 2023. O-GlcNAcylation regulates centrosome behavior and cell polarity to reduce pulmonary fibrosis and maintain the epithelial phenotype. *Adv Sci*, 10(36):e2303545.  
<https://doi.org/10.1002/advs.202303545>
- Zhong F, Wang F, Yang HF, 2024. Composition and structure analysis of different depths in the stratum corneum using confocal Raman microscopy combined with two-dimensional correlation spectroscopy. *Talanta*, 270:125559.  
<https://doi.org/10.1016/j.talanta.2023.125559>
- Zhong T, Wu XF, Xie W, et al., 2022. ENKD1 promotes epidermal stratification by regulating spindle orientation in basal keratinocytes. *Cell Death Differ*, 29(9):1719-1729.  
<https://doi.org/10.1038/s41418-022-00958-5>
- Zhou BH, Lin WL, Long YL, et al., 2022. Notch signaling pathway: architecture, disease, and therapeutics. *Signal Transduct Target Ther*, 7:95.  
<https://doi.org/10.1038/s41392-022-00934-y>
- Zhu JW, Wen WY, Zheng Z, et al., 2011. LGN/mInsc and LGN/NuMA complex structures suggest distinct functions in asymmetric cell division for the Par3/mInsc/LGN and Gai/LGN/NuMA pathways. *Mol Cell*, 43(3):418-431.  
<https://doi.org/10.1016/j.molcel.2011.07.011>