



## Transgenic barley with overexpressed *PTrx* increases aluminum resistance in roots during germination\*

Qiao-yun LI<sup>§1</sup>, Hong-bin NIU<sup>§1</sup>, Jun YIN<sup>†‡1</sup>, Hong-bo SHAO<sup>†‡2,3</sup>,  
 Ji-shan NIU<sup>1</sup>, Jiang-ping REN<sup>1</sup>, Yong-chun LI<sup>1</sup>, Xiang WANG<sup>1</sup>

<sup>1</sup>National Engineering Research Center for Wheat, Henan Agricultural University, Zhengzhou 450002, China)

<sup>2</sup>Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China)

<sup>3</sup>Institute of Life Sciences, Qingdao University of Science and Technology (QUST), Qingdao 266042, China)

<sup>†</sup>E-mail: xmzxyj@126.com; shaohongbochu@126.com

Received Feb. 10, 2010; Revision accepted Apr. 5, 2010; Crosschecked Oct. 15, 2010

**Abstract:** A transgenic barley line (LSY-11-1-1) with overexpressed *Phalaris coerulescens* thioredoxin gene (*PTrx*) was employed to measure the growth, protein oxidation, cell viability, and antioxidase activity in barley roots during germination on the presence of 2 mmol/L AlCl<sub>3</sub> on filter paper. The results show that (1) compared with the non-transgenic barley, LSY-11-1-1 had enhanced root growth, although both were seriously inhibited after AlCl<sub>3</sub> treatment; (2) the degree of protein oxidation and loss of cell viability in roots of LSY-11-1-1 were much less than those in roots of non-transgenic barley, as reflected by lower contents of protein carbonyl and Evans blue uptakes in LSY-11-1-1; (3) activities of catalase (CAT), glutathione peroxidase (GPX), ascorbate peroxidase (APX), and glutathione reductase (GR) in LSY-11-1-1 root tips were generally higher than those in non-transgenic barley root tips, although these antioxidase activities gave a rise to different degrees in both LSY-11-1-1 and non-transgenic barley under aluminum stress. These results indicate that overexpressing *PTrx* could efficiently protect barley roots from oxidative injury by increasing antioxidase activity, thereby quenching ROS caused by AlCl<sub>3</sub> during germination. These properties raise the possibility that transgenic barley with overexpressed *PTrx* may be used to reduce the aluminum toxicity in acid soils.

**Key words:** *Phalaris coerulescens* thioredoxin gene (*PTrx*), Aluminum, Transgenic barley, Oxidative stress  
 doi:10.1631/jzus.B1000048 Document code: A CLC number: Q943.2

### 1 Introduction

Numerous environmental stresses induce the formation of reactive oxygen species (ROS) in plant cells (Boscolo *et al.*, 2003). Excessive metallic ions

are a well-known example of environmental stress, and aluminum is a particularly important ion because it is involved in a number of physiological processes, such as increasing antioxidase activity (Cakmak and Horst, 1991; Yamamoto *et al.*, 2001), and inducing the expression of various genes related to oxidative stress (Richards *et al.*, 1998; Maron *et al.*, 2008; Goodwin and Sutter, 2009). Corresponding with this, there is a complex scavenging system to cope with excessive production of ROS in plant cells, including low molecular weight compounds such as ascorbic acid and glutathione, and various antioxidases (Vieira Dos Santos and Rey, 2006) such as guaiacol peroxidase (POD), catalase (CAT), superoxide dismutase

<sup>‡</sup> Corresponding authors

<sup>§</sup> The two authors contributed equally to this work

\* Project supported by the National Natural Science Foundation of China (No. 30871530), the Science and Technology Transformation Plan of Henan Province (No. 0636000005), One Hundred-Talent Plan of Chinese Academy of Sciences (CAS), CAS-local Government Cooperative Project, the CAS/SAFEA International Partnership Program for Creative Research Teams, and CAS Young Scientists Fellowship (No. 2009Y2B211), China

© Zhejiang University and Springer-Verlag Berlin Heidelberg 2010

(SOD), glutathione peroxidase (GPX), and glutathione reductase (GR) (Mittler, 2002). The ROS scavenging system efficiently reduces ROS under normal conditions, but complete reduction might not occur in stress environments, and the result may be a state of oxidative stress causing oxidative damage of biomolecules, such as lipids, proteins, and DNA, or even causing cell death (Delisle *et al.*, 2001; Pan *et al.*, 2001; Boscolo *et al.*, 2003; Shao *et al.*, 2007a; 2007b; 2008; 2009).

Thioredoxin (Trx) is a family of small, 12 kDa proteins, and can be found in all plants. The active site (Trp-Cys-Gly(Pro)-Pro-Cys) of these proteins contains a redox-active disulfide group (Lemaire *et al.*, 2003; Wong *et al.*, 2004). The exact function of Trx is still largely unknown. There are more and more reports, however, on the participation of plant Trx in the response to oxidative stress (Rey *et al.*, 2005; Vieira Dos Santos and Rey, 2006; Shao *et al.*, 2007a; 2007b; 2008; 2009).

The development of molecular biology has provided new insights into plant Trx functions in the oxidative stress response. ROS induces the expression of plant *Trx* genes. The amount of *Arabidopsis Trxh5* expression, for example, increases remarkably under oxidative stress conditions (Laloi *et al.*, 2004), and methyl viologen induces an increase in *Trxh* transcript abundance in rice seedlings (Tsukamoto *et al.*, 2005; Vieira Dos Santos and Rey, 2006). As well, target proteins of plant Trx are involved in antioxidative mechanisms. One of the 2-Cys peroxiredoxins (Prxs) and B-type methionine sulfoxide reductase (MsrB), for example, have been identified as targets of CDSP32 (the chloroplastic drought-induced stress protein, composed of two Trx modules) (Rey *et al.*, 2005). Several plant GPXs display peroxidase activity preferentially in the presence of Trx (Vieira Dos Santos and Rey, 2006). In addition, there are some Trx-regulated enzymes, such as CAT, SOD, and germin-like proteins (Balmer *et al.*, 2004). Finally, Trx may protect plant from oxidative damage as antioxidants *in vivo*. The sensitivity and lipid peroxidation levels of transformed plants lacking CDSP32 are higher, for example, than those of wild-type plants upon photo-oxidative stress exposure (Broin and Rey, 2003).

These results indicate that plant Trx fulfils an important function in protecting the plant from oxi-

dativ damage, and raises the possibility that transgenic barley with overexpressed *PTrx* will improve tolerance of aluminum stress. The *PTrx* gene belongs to the thioredoxin family because it shares nearly 94% sequence identity with *Trxh*, and its expression product has the same bio-functions and active site as *Trxh* (Li *et al.*, 1995). In the present study, we measured the change of the growth, protein oxidation, cell viability, and antioxidase activity in roots of germinating barley seeds treated with 2 mmol/L  $AlCl_3$  on filter paper. The main aim was to determine whether the transgenic barley with overexpressed *PTrx* activates protective responses to aluminum stress through increasing antioxidase activity during germination.

## 2 Materials and methods

### 2.1 Materials

Non-transgenic barley variety (*Hordeum vulgare* L. cv. LSY) and its third generation transgenic lines (LSY-11-1-1) transferred with exogenous *PTrx* (isolated from *Phalaris coerulea*, mRNA No. GB: AF159388) by particle bombardment, were used in this study.

### 2.2 Polymerase chain reaction (PCR) analysis

Genomic DNA was extracted from leaf tissue according to cetyltrimethylammonium bromide (CTAB) method described by Wang and Fang (1998). Specific primers were designed according to the coding sequence of *PTrx* gene. The sequences of the primers were P1 (5'-TTCTGTGCCAGCCATGCT TAT-3') to target the endosperm-specific  $\alpha$ -gliadin promoter region and P2 (5'-GTCCCAGTCCTCT TTGGTAGTTATG-3') to target the *PTrx* sequence. PCR products with loading dye were separated on a 1.5% (w/v) agarose gel by electrophoresis and scanned using a gel scanner JEDA801E (Jieda Science and Technology, Jiangsu, China).

### 2.3 Reverse transcriptase PCR (RT-PCR) analysis and sulfhydryl content in seeds

Seed samples of transgenic and non-transgenic lines were prepared and total RNA was isolated from the samples using the RNeasy kit (Sangon, Shanghai, China) according to the manufacturer's instructions.

RT-PCR analysis was used to determine transcript levels both in transgenic and non-transgenic lines according to the method of Li *et al.* (2009). Specific primers P3 (5'-CTCCCGATCCCAGGGCCTTCA-3') and P4 (5'-GGCGGAAAAGACACGGAAACTG-3') for *PTrx* were used in the RT-PCR analysis. The reaction included an initial 3 min of denaturation at 94 °C, followed by 27 cycles of 30 s at 94 °C, 30 s at 59 °C, and 40 s at 72 °C, a final extension of 10 min at 72 °C. As a control, a 415-bp PCR fragment of the barley constitutively-expressed actin gene *Hvpro1* (AK248710) (Sato *et al.*, 2009) was amplified from the above samples. The primers for *Hvpro1* were HvPr-1 (5'-CAGCATTGTAGGAAGGCCACG-3') and HvPr-2 (5'-CCACCACTGAGAACAACATTACCG-3'). The PCR parameters for amplifying *Hvpro1* fragments were the same as those for *PTrx*. PCR products were separated and photographed as described in PCR analysis. Two bands, 415 and 285 bp, were detected for the *Hvpro1* and *PTrx* genes, respectively.

The sulfhydryl content was measured by the method of Ellman (1959). Flours (0.15 g) were homogenized in 10 ml of Tris-glycine buffer (0.08 mol/L, pH 8.0) containing 3 mmol/L ethylene diamine tetraacetic acid (EDTA), and then 4 ml denaturation reagent (8 mol/L urea) and 0.1 ml Ellman's reagent [Tris-glycine buffer containing 0.4% (w/v) 2,2'-dinitro-5,5'-dithiodibenzoate (DTNB)] were added in 1 ml samples. Thionitrobenzoate, released by sulfhydryl groups of proteins reacted with DTNB, was measured at 412 nm, and the results were expressed as  $\mu\text{mol}$  sulfhydryl/g dry weight (DW).

#### 2.4 Germination conditions

Germination and growth conditions were conducted according to the method of Tamás *et al.* (2006). Seeds were surface sterilized with 12%  $\text{H}_2\text{O}_2$  for 10 min and then rinsed three times with sterilized distilled water. After 4 h of incubation in 8.2 mmol/L  $\text{CaCl}_2$  solution (pH 4.0, untreated) or in 6.2 mmol/L  $\text{CaCl}_2$  containing 2 mmol/L  $\text{AlCl}_3$  solution (pH 4.0, Al-treated) at 25 °C in darkness, the seeds were germinated between two layers of filter paper fully moistened with the same solutions, and the germinating seeds were transferred on the freshly moistened filter papers with appropriate solutions every 20 h. Root growth was measured by the length of the main roots.

#### 2.5 Protein oxidation and cell viability

The degree of protein oxidation was measured by the reaction of carbonyls with 2,4-dinitrophenylhydrazine (DNPH), according to the method of Levine *et al.* (1990). Fresh root tips (0.2 g) were homogenized in 2 ml phosphate buffer solution (25 mmol/L, pH 7.0) containing 0.1 mmol/L EDTA and 1 mmol/L phenylmethanesulfonyl fluoride (PMSF), and then centrifuged at  $15000\times g$  for 15 min. After the supernatant reaction with DNPH, the carbonyl content was calculated by absorbance of aliphatic hydrazones [extinction coefficient: 22.1 L/(mmol·cm)] at 374 nm and expressed as nmol carbonyl/mg protein.

The cell viability loss was investigated through Evans blue staining method as described by Yamamoto *et al.* (2001) and Boscolo *et al.* (2003), with minor modifications. Fresh roots were washed three times with deionized water and stained with 0.25% (w/v) Evans blue aqueous solution for 15 min. After rinsing with deionized water for 30 min, 20 root tips (5 mm) were excised and the accumulated Evans blue was released by dipping the root sections in 3 ml of *N,N*-dimethylformamide reagent for 1 h at 37 °C, and then centrifuged at  $3000\times g$  for 5 min. The optical density (OD) of the supernatant was determined spectrophotometrically at 600 nm.

Soluble protein concentrations were measured using the method of Bradford (1976).

#### 2.6 Enzyme activity

Enzyme activity was measured spectrophotometrically, and temperature was controlled to  $(25\pm 0.5)$  °C.

CAT activity was determined as outlined by Cakmak and Horst (1991), and the activity was analyzed by measuring the degradation of  $\text{H}_2\text{O}_2$ , which was monitored at 240 nm. One unit of CAT decomposed 1  $\mu\text{mol}$ /L of  $\text{H}_2\text{O}_2$  [extinction coefficient: 39.4 L/(mmol·cm)] per minute. GR activity was assayed following the procedure of Schaedle and Bassham (1977), measuring the decrease of optical density at 340 nm, with the extinction coefficient of 6.22 L/(mmol·cm). One unit of activity is defined as the amount of enzyme that oxidizes 1  $\mu\text{mol}$  of nicotinamide adenine dinucleotide phosphate (NADPH) per minute. GPX activity was determined following the method of Livingstone *et al.* (1992). The assay was

monitored at 340 nm and one unit of activity is defined as the amount of enzyme that oxidizes 1  $\mu\text{mol}$  of NADPH [extinction coefficient: 6.22 L/(mmol·cm)] per minute. APX activity was measured according to Asada (1984) and the activity was recorded by the decrease of optical density at 290 nm, taking 2.8 L/(mmol·cm) as the extinction coefficient, and one unit of activity is defined as the amount of enzyme that oxidizes 1  $\mu\text{mol}$  of ascorbate per minute.


### 2.7 Statistical analysis

Each experiment was repeated at least five times. Data were expressed as mean $\pm$ standard error (SE). Variance analysis between different treatments was carried out with the SPSS 11.5 program using Duncan's multiple range test at  $P<0.01$ .

## 3 Results

### 3.1 PCR analysis

The presence of exogenous *PTrx* gene in transgenic barley was examined with PCR analysis. DNA from leaves of the transgenic barley lines (LSY-11-1-1) was amplified by PCR reaction using primers specific for exogenous *PTrx*. The transgenic plants exhibited the polymorphic band 810 bp in length, the same size as the positive control (plasmid), but non-transgenic plant DNA was negative (Fig. 1). The results showed that exogenous *PTrx* gene had been transferred in barley genome.




**Fig. 1** PCR analysis of transgenic barley lines (LSY-11-1-1) 1–13: transgenic plants of LSY-11-1-1 (the next generation from T<sub>5</sub> in Figs. 2 and 3); CK: non-transgenic plants; P: positive control (plasmid); M: marker

### 3.2 RT-PCR analysis and sulfhydryl content

To investigate the expression of exogenous *PTrx* gene in LSY-11-1-1, the template cDNAs from transgenic and non-transgenic barley seeds 15 d after anthesis were assayed by RT-PCR and the sulfhydryl


content in the mature seeds was measured. The specific band of 215 bp was exhibited in transgenic seeds, while non-transgenic seeds were negative (Fig. 2). These results indicate that exogenous *PTrx* gene can be transcribed and expressed normally in transgenic barley seeds.



**Fig. 2** RT-PCR analysis of *PTrx* expression in seeds of transgenic barley 15 d after anthesis

CK: non-transgenic plants; T<sub>1</sub>–T<sub>5</sub>: seeds of different transgenic lines of the third generation (T<sub>5</sub> is the plants of LSY-11-1-1)

The sulfhydryl content in seeds of transgenic barley was generally higher than that in non-transgenic (Fig. 3,  $P<0.01$ ), and, in LSY-11-1-1, was 1.3 times that of non-transgenic barley.




**Fig. 3** Sulfhydryl content in seeds of transgenic and non-transgenic barley seeds

Data are expressed as mean $\pm$ SE ( $n=5$ ); Bars (i.e. means) with different letters are significantly different ( $P<0.01$ ). CK: non-transgenic; T<sub>1</sub>–T<sub>5</sub>: seeds of different transgenic lines of the third generation (T<sub>5</sub> is the seeds of LSY-11-1-1)

### 3.3 Effect of Al-treatment on root growth

As seen in Fig. 4, the root growth of germinating barley seeds was obviously inhibited between 72 and 96 h after the onset of seed imbibition treated by 2 mmol/L AlCl<sub>3</sub>. Although a continuous decrease of the root growth of both transgenic and non-transgenic germinating barley seeds was observed, the root length of LSY-11-1-1 was obviously longer than that of non-transgenic barley.



**Fig. 4** Root length as a function of time of transgenic barley (LSY-11-1-1) and non-transgenic barley under  $\text{Al}^{3+}$  stress during germination

Data are expressed as mean $\pm$ SE ( $n=5$ )

### 3.4 Protein oxidation and cell viability


Carbonyl group content in the transgenic and non-transgenic barley root tips were presented in Fig. 5a. The result showed that protein oxidation of LSY-11-1-1 was clearly lower than that of non-transgenic barley. The carbonyl content of LSY-11-1-1 was decreased, for example, by 37.0%, 20.8%, and 23.3% after 2 mmol/L  $\text{AlCl}_3$  stress for 72, 84, and 96 h, respectively, compared with the results of non-transgenic barley.

The Evans blue uptakes in root tips were significantly increased after  $\text{AlCl}_3$  treatment (Fig. 5b). The non-transgenic barley showed the highest value after 96 h treatment, and it was 2.9-fold higher than that of the untreated sample. Compared to the non-transgenic barley, the Evans blue uptakes in LSY-11-1-1 were significantly lower ( $P<0.01$ ), only 66.0%, 68.5%, and 56.1% that of non-transgenic barley after 48, 72, and 96 h treatments, respectively.

### 3.5 Effect of Al-treatment on antioxidase activity

The changes in antioxidase activities are shown in Table 1. Our experimental results demonstrated that the activities of CAT, GPX, GR, and APX were significantly increased after aluminum stress, except that of GR at 48 h ( $P<0.01$ ).

Compared to non-transgenic barley, the activities of these antioxidases were enhanced significantly in LSY-11-1-1 root tips ( $P<0.01$ ). In the presence of 2 mmol/L  $\text{Al}^{3+}$  treatment, the CAT activity of LSY-11-1-1 was 11.8%, 18.9%, 32.6%, and 17.6% higher than that of non-transgenic barley at 60, 72, 84, and 96 h after treatment, respectively. *PTrx* overexpressing



**Fig. 5** Contents of protein carbonyls (a) and Evans blue uptakes ( $\text{OD}_{600}$ ) (b) as a function of time in transgenic (LSY-11-1-1) and non-transgenic barley roots under  $\text{Al}^{3+}$  stress during germination

Data are expressed as mean $\pm$ SE ( $n=5$ ); Bars (i.e., means) with different letters are significantly different ( $P<0.01$ )

induced similar activity changes of GPX, the maximal value of GPX in LSY-11-1-1 was 96.7 U/mg protein at the activity peak, while it was 76.8 U/mg protein in non-transgenic barley. Compared with non-transgenic barley, the GPX activity of LSY-11-1-1 was significantly increased by 14.6%, 40.1%, and 30.9% after 72, 84, and 96 h treatments, respectively. APX activity of LSY-11-1-1 was 7.9%, 13.9%, 10.2%, and 11.3% greater than that of non-transgenic barley at 60, 72, 84, and 96 h after treatment, respectively. Although the GR activity was not significantly different between LSY-11-1-1 and non-transgenic barley from 48 to 72 h after  $\text{Al}^{3+}$  stress, it was significantly higher in LSY-11-1-1 at the other time sampled. The GR activity of LSY-11-1-1 was 1.2 and 1.1 times of that of non-transgenic barley at 84 and 96 h, respectively.

## 4 Discussion

We transferred exogenous *PTrx* gene into malting barley variety and obtained transgenic plants

**Table 1** CAT, GPX, APX, and GR activities as a function of time in barley roots under Al<sup>3+</sup> stress during germination

Time (h)	Group	CAT activity (U/mg protein)	GPX activity (U/mg protein)	APX activity (mU/mg protein)	GR activity (mU/mg protein)
48	Non-transgenic	25.65±0.66 <sup>a</sup>	41.46±1.19 <sup>a</sup>	398.77±3.38 <sup>a</sup>	151.44±0.83 <sup>a</sup>
	LSY-11-1-1	24.47±0.68 <sup>a</sup>	39.93±0.81 <sup>a</sup>	418.66±9.84 <sup>ab</sup>	145.48±1.83 <sup>a</sup>
	Al-non-transgenic	50.69±1.23 <sup>b</sup>	68.14±1.39 <sup>b</sup>	455.27±6.89 <sup>b</sup>	148.62±1.38 <sup>a</sup>
	Al-LSY-11-1-1	50.04±0.74 <sup>b</sup>	70.63±0.65 <sup>b</sup>	453.01±9.55 <sup>b</sup>	150.58±1.92 <sup>a</sup>
60	Non-transgenic	35.04±0.75 <sup>a</sup>	39.99±1.32 <sup>a</sup>	340.17±4.11 <sup>a</sup>	130.69±1.72 <sup>a</sup>
	LSY-11-1-1	38.96±0.93 <sup>a</sup>	39.11±0.90 <sup>a</sup>	321.33±2.40 <sup>a</sup>	139.93±1.09 <sup>b</sup>
	Al-non-transgenic	47.56±0.92 <sup>b</sup>	76.77±1.91 <sup>b</sup>	377.33±5.24 <sup>b</sup>	203.44±1.99 <sup>c</sup>
	Al-LSY-11-1-1	53.19±1.69 <sup>c</sup>	77.85±1.01 <sup>b</sup>	407.33±4.67 <sup>c</sup>	209.59±2.11 <sup>c</sup>
72	Non-transgenic	30.48±1.46 <sup>a</sup>	36.88±1.38 <sup>a</sup>	381.67±5.24 <sup>b</sup>	134.60±2.56 <sup>a</sup>
	LSY-11-1-1	32.59±1.80 <sup>a</sup>	41.85±1.06 <sup>a</sup>	344.00±3.21 <sup>a</sup>	123.79±1.63 <sup>a</sup>
	Al-non-transgenic	46.30±1.10 <sup>b</sup>	56.96±1.48 <sup>b</sup>	493.65±5.04 <sup>c</sup>	198.72±1.29 <sup>b</sup>
	Al-LSY-11-1-1	55.07±1.03 <sup>c</sup>	65.26±0.63 <sup>c</sup>	562.66±3.38 <sup>d</sup>	208.18±2.39 <sup>b</sup>
84	Non-transgenic	32.18±1.21 <sup>a</sup>	44.00±1.74 <sup>a</sup>	452.78±5.59 <sup>b</sup>	161.14±1.71 <sup>a</sup>
	LSY-11-1-1	31.38±1.14 <sup>a</sup>	44.30±2.09 <sup>a</sup>	402.66±6.84 <sup>a</sup>	153.59±2.00 <sup>a</sup>
	Al-non-transgenic	55.00±1.61 <sup>b</sup>	68.99±0.71 <sup>b</sup>	693.66±8.84 <sup>c</sup>	239.11±1.74 <sup>b</sup>
	Al-LSY-11-1-1	72.92±1.72 <sup>c</sup>	96.66±2.17 <sup>c</sup>	764.43±10.49 <sup>d</sup>	275.01±2.08 <sup>c</sup>
96	Non-transgenic	38.11±1.49 <sup>a</sup>	46.29±0.99 <sup>a</sup>	410.00±6.43 <sup>a</sup>	136.63±0.38 <sup>a</sup>
	LSY-11-1-1	34.40±1.41 <sup>a</sup>	39.79±0.76 <sup>a</sup>	441.63±4.94 <sup>a</sup>	140.01±1.07 <sup>a</sup>
	Al-non-transgenic	52.24±1.26 <sup>b</sup>	70.87±1.15 <sup>c</sup>	677.25±3.76 <sup>b</sup>	234.40±2.41 <sup>b</sup>
	Al-LSY-11-1-1	61.44±2.03 <sup>c</sup>	92.78±1.45 <sup>d</sup>	753.66±18.22 <sup>c</sup>	269.41±1.23 <sup>c</sup>

Data are expressed as mean±SE (n=5). Means followed by different letters are significantly different ( $P<0.01$ )

by the detection of PCR analysis. In order to confirm the expression of *PTrx*, the specific band of 215 bp in the transgenic line was lifted by RT-PCR, while the non-transgenic barley was negative. This revealed a positive result, indicating that exogenous *PTrx* gene had been successfully transferred and expressed in transgenic barley. The active site Trp-Cys-Gly(Pro)-Pro-Cys of Trx contains a highly reactive dithiol, which may reduce disulfide bridges of many proteins by thiol-disulfide interchange (Lemaire *et al.*, 2003), so overexpressed *PTrx* will enhance the reducing ability in the transgenic barley seeds. Our experimental data demonstrate that the sulfhydryl content in the seeds of transgenic barley is higher than that in the non-transgenic barley, which also explains the *PTrx* expression.

Root growth of both transgenic barley and non-transgenic barley was inhibited seriously in the presence of Al<sup>3+</sup>, consistent with the results reported by Kochian (1995) and Boscolo *et al.* (2003), but the transgenic barley showed a stronger root growth than non-transgenic barley on filter paper with AlCl<sub>3</sub> treatment during germination, as reflected by the greater root length. These results demonstrate that the




phytotoxic effects of Al<sup>3+</sup> on root growth are less severe in transgenic barley root growth than in non-transgenic barley root growth.

Oxidative stress is possibly an important component of the plant's reaction to aluminum toxicity (Pan *et al.*, 2001). Several studies have shown that aluminum treatment causes excessive ROS formation (Cakmak and Horst, 1991; Boscolo *et al.*, 2003). In our experiment, the increased activities of SOD, POD, CAT, APX, GPX, and GR induced by aluminum treatment suggest that Al<sup>3+</sup> induces ROS excessive production.

The result of excessive ROS formation may cause biomolecule oxidation or even cell death (Cakmak and Horst, 1991; Delisle *et al.*, 2001; Pan *et al.*, 2001; Yamamoto *et al.*, 2001). We investigated protein oxidation through a determination of the amount of carbonyl, an indicator that describes protein oxidation degree caused by ROS (Levine *et al.*, 1990; Boscolo *et al.*, 2003), and cell death through changes of Evans blue uptakes, a parameter of loss of cell viability, another parameter of oxidative stress damage in plant (Yamamoto *et al.*, 2001; Boscolo *et al.*, 2003). The lower protein carbonyl contents and

Evans blue uptakes in roots of LSY-11-1-1 indicate that excessive ROS, caused by AlCl<sub>3</sub> stress, was eliminated more efficiently in the transgenic barley roots than in the non-transgenic barley roots during germination, so the degree of the damage caused by AlCl<sub>3</sub> stress in transgenic plants was less than that in non-transgenic plants, which may be due to its effective antioxidase system.

In general, the major ROS-scavenging enzyme pathways of plants include SOD [Eq. (1)], CAT [Eq. (2)], the ascorbate-glutathion cycle [Eq. (3)], and GPX cycle [Eq. (4)] (Mittler, 2002).



In our experiment, AlCl<sub>3</sub> treatment induced a significant increase in SOD, POD, CAT, GPX, APX, and GR activities, although the activities of SOD and POD were similar changes in both LSY-11-1-1 and non-transgenic barley (data not shown). It is apparent that all the ROS-scavenging enzyme pathways of plants were indispensable for ROS detoxification in roots during barley seed germination in the presence of 2 mmol/L AlCl<sub>3</sub>, but the CAT pathway, ascorbate glutathion cycle, and GPX cycle played a more important role on the progress of quenching ROS caused by Al<sup>3+</sup> stress in transgenic barley, and the increased antioxidase activity contributed to a more effective reduction in the oxidative damage in roots during germination.

It is suggested that the redox state of proteins, including enzymes and regulatory components, is controlled by cellular redox agents, the paramount one among which is Trx (Wong *et al.*, 2004). Several antioxidases, protecting plant cells from oxidative stress, are potential targets of Trx, such as CAT, SOD (Cu-Zn), and GR (Jung *et al.*, 2002), almost all of the which contain conserved cysteine, e.g., the

numbers of conserved cysteine of CAT, GPX, POD, SOD (Cu-Zn), APX, 2-Cys Prxs, and 1-Cys Prxs are 5, 1, 8, 2, 2, 1, and 1, respectively (Wong *et al.*, 2004). Trx might be involved in modulating redox-dependent signaling cascades, as shown by Schenk *et al.* (1994), or reduced Trx transfers its electrons to a target disulfide (Meyer *et al.*, 1999; Wong *et al.*, 2004), as a result providing reducing equivalents for regulating activity and increasing the antioxidase activity (Wong *et al.*, 2004). Moreover, the thioredoxin gene and several genes encoding Cys-rich proteins were previously shown to be induced by aluminum stress (Maron *et al.*, 2008; Goodwin and Sutter, 2009), suggesting that thioredoxins could play similar role as antioxidants.

Aluminum is the most abundant metal in the earth's crust, and aluminum toxicity is a very important factor limiting crop growth in acidic soil (Kochian, 1995). Thus, it is necessary to enhance the aluminum endurance of plants. Increased activities of antioxidases endow the transgenic barley with more effective elimination of excessive ROS induced by aluminum stress. This decreases the damage of biomolecules by oxidation and contributes to the integrity of membrane structure. In turn, aluminum is prevented from exerting its full physiological toxicity in the roots, as reflected by the fact that the roots of transgenic barley germinating seeds showed better growth than those of non-transgenic barley. It is possible that, in the future, *PTrx*-overexpressed barley can grow in aluminum-polluted environments without decreasing production because of the enhanced aluminum resistance.

## References

- Asada, K., 1984. Chloroplasts: formation of active oxygen and its scavenging. *Methods Enzymol.*, **105**:422-429. [doi:10.1016/S0076-6879(84)05059-X]
- Balmer, Y., Vensel, W.H., Tanaka, C.K., Hurkman, W.J., Gelhaye, E., Rouhier, N., Jacquot, J.P., Manieri, W., Schürmann, P., Droux, M., *et al.*, 2004. Thioredoxin links redox to the regulation of fundamental processes of plant mitochondria. *PNAS*, **101**(8):2642-2647. [doi:10.1073/pnas.0308583101]
- Boscolo, P.R.S., Menossi, M., Jorge, R.A., 2003. Aluminum-induced oxidative stress in maize. *Phytochemistry*, **62**(2): 181-189. [doi:10.1016/S0031-9422(02)00491-0]
- Bradford, M.M., 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*,

- 72(1-2):248-254. [doi:10.1016/0003-2697(76)90527-3]
- Broin, M., Rey, P., 2003. Potato plants lacking the CDSP32 plastidic thioredoxin exhibit over-oxidation of the BAS1 2-Cys peroxiredoxin and increased lipid peroxidation in thylakoids under photooxidative stress. *Plant Physiol.*, **132**(3):1335-1343. [doi:10.1104/pp.103.021626]
- Cakmak, I., Horst, W.J., 1991. Effect of aluminum on lipid peroxidation, superoxide dismutase, catalase, and peroxidases activities in root tips of soybean (*Glycine max*). *Physiol. Plantarum*, **83**(3):463-468. [doi:10.1034/j.1399-3054.1991.830320.x]
- Delisle, G., Champoux, M., Houde, M., 2001. Characterization of oxalate oxidase and cell death in Al-sensitive and tolerant wheat roots. *Plant Cell Physiol.*, **42**(3):324-333. [doi:10.1093/pcp/pce041]
- Ellman, G.D., 1959. Tissue sulfhydryl groups. *Arch. Biochem. Biophys.*, **82**(1):70-77. [doi:10.1016/0003-9861(59)90090-6]
- Goodwin, S.B., Sutter, T.R., 2009. Microarray analysis of *Arabidopsis* genome response to aluminum stress. *Biol. Plantarum*, **53**(1):85-99. [doi:10.1007/s10535-009-0012-4]
- Jung, B.G., Lee, K.O., Lee, S.S., Chi, Y.H., Jang, H.H., Kang, S.S., Lee, K., Lim, D., Yoon, S.H., Yun, D.J., et al., 2002. A Chinese cabbage cDNA with high sequence identity to phospholipid hydroperoxide glutathione peroxidases encodes a novel isoform of thioredoxin-dependent peroxidase. *J. Biol. Chem.*, **277**(15):12572-12578. [doi:10.1074/jbc.M110791200]
- Kochian, L.V., 1995. Cellular mechanisms of aluminum toxicity and resistance in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, **46**(1):237-260. [doi:10.1146/annurev.pp.46.060195.001321]
- Laloi, C., Mestres-Ortega, D., Marco, Y., Meyer, Y., Reichheld, J.P., 2004. The *Arabidopsis* cytosolic thioredoxin h5 gene induction by oxidative stress and its W-box-mediated response to pathogen elicitor. *Plant Physiol.*, **134**(3):1006-1016. [doi:10.1104/pp.103.035782]
- Lemaire, S.D., Collin, V., Keryer, E., Issakidis-Bourguet, E., Lavergne, D., 2003. *Chlamydomonas reinhardtii*: a model organism for the study of the thioredoxin family. *Plant Physiol. Biochem.*, **41**(6-7):513-521. [doi:10.1016/S0981-9428(03)00079-2]
- Levine, R.L., Garland, D., Olive, C.N., Amici, A., Climent, I., Lenz, A.G., Ahn, B.W., Shaltiel, S., Stadtman, E.R., 1990. Determination of carbonyl content in oxidatively modified proteins. *Methods Enzymol.*, **186**:464-478. [doi:10.1016/0076-6879(90)86141-H]
- Li, X.M., Nield, J., Hayman, D., Langridge, P., 1995. Thioredoxin activity in the C terminus of Phalaris S protein. *Plant J.*, **8**(1):133-138. [doi:10.1046/j.1365-313X.1995.08010133.x]
- Li, Y.C., Ren, J.P., Cho, M.J., Zhou, S.M., Kim, Y.B., Guo, H.X., Wong, J.H., Niu, H.B., Kim, H.K., Morigasaki, S., et al., 2009. The level of expression of thioredoxin is linked to fundamental properties and applications of wheat seeds. *Mol. Plant*, **2**(3):430-441. [doi:10.1093/mp/ssp025]
- Livingstone, D.R., Lips, F., Garcia Martinez, P., Pipe, P.K., 1992. Antioxidant enzymes in the digestive gland of the common mussel *Mytilus edulis*. *Mar. Biol.*, **112**(2):265-276. [doi:10.1007/BF00702471]
- Maron, L.G., Kirst, M., Mao, C., Milner, M.J., Menossi, M., Kochian, L.V., 2008. Transcriptional profiling of aluminum toxicity and tolerance responses in maize roots. *New Phytol.*, **179**(1):116-128. [doi:10.1111/j.1469-8137.2008.02440.x]
- Meyer, Y., Verdoucq, L., Vignols, F., 1999. Plant thioredoxins and glutaredoxins: identity and putative roles. *Trends Plant Sci.*, **4**(10):388-394. [doi:10.1016/S1360-1385(99)01475-2]
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.*, **7**(9):405-410. [doi:10.1016/S1360-1385(02)02312-9]
- Pan, J.W., Zhu, M.Y., Chen, H., 2001. Aluminum-induced cell death in root-tip cells of barley. *Environ. Exp. Bot.*, **46**(1):71-79. [doi:10.1016/S0098-8472(01)00083-1]
- Rey, P., Cuiné, S., Eymery, F., Garin, J., Court, M., Jacquot, J.P., Rouhier, N., Broin, M., 2005. Analysis of the proteins targeted by CDSP32, a plastidic thioredoxin participating in oxidative stress responses. *Plant J.*, **41**(1):31-42. [doi:10.1111/j.1365-313X.2004.02271.x]
- Richards, K.D., Schott, E.J., Sharma, Y.K., Davis, K.R., Gardner, R.C., 1998. Aluminum induces oxidative stress genes in *Arabidopsis thaliana*. *Plant Physiol.*, **116**(1):409-418. [doi:10.1104/pp.116.1.409]
- Sato, K., Shin-I, T., Seki, M., Shinozaki, K., Yoshida, H., Takeda, K., Yamazaki, Y., Conte, M., Kohara, Y., 2009. Development of 5006 full-length cDNAs in barley: a tool for accessing cereal genomics resources. *DNA Res.*, **16**(2):81-89. [doi:10.1093/dnares/dsn034]
- Schaedle, M., Bassham, J.A., 1977. Chloroplast glutathione reductase. *Plant Physiol.*, **59**(5):1011-1012. [doi:10.1104/pp.59.5.1011]
- Schenk, H., Klein, M., Erdbrugger, W., Droge, W., Schulze-Osthoff, K., 1994. Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF- $\kappa$ B and AP-1. *PNAS*, **91**(5):1672-1676.
- Shao, H.B., Jiang, S.Y., Li, F.M., Chu, L.Y., Shao, M.A., Li, F., 2007a. Some advances in plant stress physiology and their implications in the systems biology era. *Colloids Surf. B Biointerfaces*, **54**(1):33-36. [doi:10.1016/j.colsurfb.2006.05.011]
- Shao, H.B., Chu, L.Y., Lu, Z.H., Kang, C.M., 2007b. Primary antioxidant free radical scavenging and redox signaling pathways in higher plant cells. *Int. J. Biol. Sci.*, **4**(1):8-14.
- Shao, H.B., Chu, L.Y., Shao, M.A., 2008. Calcium as a versatile plant signal transducer under soil water stress. *Bioessays*, **30**(7):634-641. [doi:10.1002/bies.20770]
- Shao, H.B., Jaleel, C.A., Manivannan, P., Panneerselvam, R., Shao, M.A., 2009. Understanding water deficit stress-induced changes in the basic metabolism of higher plants—biotechnologically and sustainably improving agriculture and the environment in arid regions of the



- globe. *Crit. Rev. Biotechnol.*, **29**(2):131-151. [doi:10.1080/07388550902869792]
- Tamás, L., Budíková, S., Šimonovičová, M., Huttova, J., Siroka, B., Mistrik, I., 2006. Rapid and simple method for Al-toxicity analysis in emerging barley roots during germination. *Biol. Plantarum*, **50**(1):87-93. [doi:10.1007/s10535-005-0079-5]
- Tsukamoto, S., Morita, S., Hirano, E., Yokoi, H., Masumura, T., Tanaka, K., 2005. A novel *cis*-element that is responsive to oxidative stress regulates three antioxidant defense genes in rice. *Plant Physiol.*, **137**(1):317-327. [doi:10.1104/pp.104.045658]
- Vieira Dos Santos, C., Rey, P., 2006. Plant thioredoxins are key actors in the oxidative stress response. *Trends Plant Sci.*, **11**(7):329-334. [doi:10.1016/j.tplants.2006.05.005]
- Wang, G.L., Fang, H.J., 1998. Principle and Technique of Plant Genetic Engineering. Science Press, Beijing (in Chinese).
- Wong, J.H., Cai, N., Balmer, Y., Tanaka, C.K., Vensel, W.H., Hurkman, W.J., Buchanan, B.B., 2004. Thioredoxin targets of developing wheat seeds identified by complementary proteomic approaches. *Phytochemistry*, **65**(11):1629-1640. [doi:10.1016/j.phytochem.2004.05.010]
- Yamamoto, Y., Kobayashi, Y., Matsumoto, H., 2001. Lipid peroxidation is an early symptom triggered by aluminum, but not the primary cause of elongation inhibition in pea roots. *Plant Physiol.*, **125**(1):199-208. [doi:10.1104/pp.125.1.199]

## Journals of Zhejiang University-SCIENCE (A/B/C)

# Latest trends and developments

These journals are among the best of China's University Journals. Here's why:

- *JZUS (A/B/C)* have developed rapidly in specialized scientific and technological areas.  
*JZUS-A (Applied Physics & Engineering)* split from *JZUS* and launched in 2005  
*JZUS-B (Biomedicine & Biotechnology)* split from *JZUS* and launched in 2005  
*JZUS-C (Computers & Electronics)* split from *JZUS-A* and launched in 2010
- We are the first in China to completely put into practice the international peer review system in order to ensure the journals' high quality (more than 7600 referees from over 60 countries, <http://www.zju.edu.cn/jzus/reviewer.php>)
- We are the first in China to pay increased attention to Research Ethics Approval of submitted papers, and the first to join CrossCheck to fight against plagiarism
- Comprehensive geographical representation (the international authorship pool enlarging every day, contributions from outside of China accounting for more than 46% of papers)
- Since the start of an international cooperation with Springer in 2006, through SpringerLink, *JZUS*'s usage rate (download) is among the tops of all of Springer's 82 co-published Chinese journals
- *JZUS*'s citation frequency has increased rapidly since 2004, on account of DOI and Online First implementation (average of more than 60 citations a month for each of *JZUS-A* & *JZUS-B* in 2009)
- *JZUS-B* is the first university journal to receive a grant from the National Natural Science Foundation of China (2009-2010)