



Variations in cadmium and nitrate co-accumulation among water spinach genotypes and implications for screening safe genotypes for human consumption*

Lin TANG¹, Wei-jun LUO¹, Zhen-li HE², Hanumanth Kumar GURAJALA¹,
 Yasir HAMID¹, Kiran Yasmin KHAN¹, Xiao-e YANG^{†‡1}

¹Ministry of Education Key Laboratory of Environmental Remediation and Ecosystem Health,
 College of Environmental and Resource Sciences, Zhejiang University, Hangzhou 310058, China

²Indian River Research and Education Center, Institute of Food and Agricultural Sciences,
 University of Florida, Fort Pierce, Florida 34945, USA

[†]E-mail: xeyang@zju.edu.cn

Received Jan. 12, 2017; Revision accepted Apr. 9, 2017; Crosschecked Jan. 27, 2018

Abstract: Vegetables are important constituents of the human diet. Heavy metals and nitrate are among the major contaminants of vegetables. Consumption of vegetables and fruits with accumulated heavy metals and nitrate has the potential to damage different body organs leading to unwanted effects. Breeding vegetables with low heavy metal and nitrate contaminants is a cost-effective approach. We investigated 38 water spinach genotypes for low Cd and nitrate co-accumulation. Four genotypes, i.e. *JXDY*, *GZQL*, *XGDB*, and *B888*, were found to have low co-accumulation of Cd (<0.71 mg/kg dry weight) and nitrate (<3100 mg/kg fresh weight) in the edible parts when grown in soils with moderate contamination of both Cd (1.10 mg/kg) and nitrate (235.2 mg/kg). These genotypes should be appropriate with minimized risk to humans who consume them. The Cd levels in the edible parts of water spinach were positively correlated with the concentration of Pb or Zn, but Cd, Pb, or Zn was negatively correlated with P concentration. These results indicate that these three heavy metals may be absorbed into the plant in similar proportions or in combination, minimizing the influx to aerial parts. Increasing P fertilizer application rates appears to prevent heavy metal and nitrate translocation to shoot tissues and the edible parts of water spinach on co-contaminated soils.

Key words: Genotypic difference; Heavy metal; Nitrate; Soil pollution; Water spinach
<https://doi.org/10.1631/jzus.B1700017>

CLC number: X53

1 Introduction

Cadmium (Cd) is one of the most toxic and mobile heavy metals that affect human health through

the food chain. Agricultural practices, intensive industrial activity, and urban expansion have accelerated the release of Cd into soil, water, and air (Lane et al., 2015). About 27860000 m² of agricultural soils in China are polluted with Cd (Liu et al., 2015) and Cd contamination has become one of the most important barriers to agricultural sustainability in China (Zhang et al., 2002). Cd in agricultural soils is taken up by plants and then enters humans and animals through the food chain (Kirkham, 2006). Phytoremediation is the best strategy to reduce the risk of Cd entry into food chain.

[‡] Corresponding author

* Project supported by the Key Projects from Ministry of Science and Technology of China (No. 2016YFD0800805), the Zhejiang Provincial Science and Technology Bureau (Nos. 2015C02011-3 and 2015C03020-2), and the Fundamental Research Funds for the Central University, China

ORCID: Lin TANG, <https://orcid.org/0000-0002-4540-4987>

© Zhejiang University and Springer-Verlag GmbH Germany, part of Springer Nature 2018

Fertilizer application is one of the key factors in improving crop productivity. However, excessive use of chemical fertilizers, especially nitrogen (N) fertilizers, results in severe environmental problems (Hakeem et al., 2013). China is one of the largest consumers of N fertilizers in the world, but the average N use efficiency is low (only about 35%) (Wang et al., 2014), which results in waste of resources and environmental contamination, and also poses a serious hazard to human health (Xu et al., 2012; Chen et al., 2014). Nitrate is potentially carcinogenic and may increase the incidence of gastric, bladder, and oesophageal cancers (Gulis et al., 2002). However, nitrate can decrease blood pressure, thus reducing the risk of cardiovascular disease, myocardial infarction, and stroke. Daily consumption of nitrate-rich vegetables is associated with beneficial effects for patients with gastric ulcer, renal failure, and metabolic syndrome (Habermeyer et al., 2015).

Vegetables are nutritious and are assumed to be safe to consume; people are unaware that some parts of the vegetable may be contaminated with heavy metals and are a major source of human exposure to Cd and nitrate (Tang et al., 2016; Fan et al., 2017). Water spinach (*Ipomoea aquatica* Forsk.), an important leafy vegetable in eastern and southern Asia, can absorb Cd and nitrate naturally into their vacuoles (Wang et al., 2009; Xin et al., 2010). Therefore, evaluation of water spinach genotypes for low Cd and nitrate accumulation when grown on contaminated soils has become a research priority in minimizing human exposure to these co-contaminants.

Accumulation of Cd by plants varies with species and genotype. New strategies have been applied to breeding and screening heavy metal low accumulator vegetable genotypes, such as Chinese cabbage (*Brassica chinensis* L.) (Liu et al., 2010; Wang X et al., 2015), pakchoi (*Brassica rapa* L. ssp. *chinensis*) (Chen et al., 2012), soybean (*Glycine max* Merr.) (Arao et al., 2003; Sugiyama et al., 2011), welsh onion (*Allium fistulosum* L.) (Li et al., 2012), and sweet potato (*Ipomoea batatas* (L.) Lam.) (Huang et al., 2015). Nitrate concentration in high-accumulation genotypes was several times greater than that in low-accumulation genotypes, such as lettuce (*Lactuca sativa* L.) (Escobar-Gutiérrez et al., 2002; Burns et al., 2011a), leaf mustard (*Brassica juncea* (L.) Czern.) (Sharma et al., 2010), and taro (*Colocasia esculenta* (L.) Schott) (Kristl et al., 2016). Vegetable genotypes with

a lower nitrate concentration need to be identified for agricultural production and improved human health.

Soils in southern China are often polluted by multiple contaminants. Among them Cd and nitrate contaminations are attributed to irrigation practices, use of low-grade organic fertilizers, and heavy application of N fertilizers, all of which can affect growth, metal tolerance, and metal accumulation in water spinach. Screening different vegetable genotypes with low absorption of Cd and nitrate provides an opportunity to safeguard human consumption.

Previous studies were limited to pot and hydroponics where correlations between different contaminants and nutrients were not observed. This study aimed to investigate co-accumulative remediation capacity of Cd and nitrate in field conditions among 38 genotypes of water spinach.

2 Materials and methods

2.1 Soil characterization

The experiment was performed in greenhouses (Chunyi farm) located at 30°23'37" N and 120°2'13" E, Hangzhou, Zhejiang Province, China with an average temperature of 30 °C throughout the day. The soil had been moderately contaminated by Cd and nitrate during several decades of intensive vegetable production (Tang et al., 2016). The depth of soil sampling for analysis was 10–15 cm, and four soil samples were collected to represent the area. The preliminary soil assessment and initial concentrations of metals in the soil (Table 1) were determined according to the previous methods (Li, 2000; Bao, 2008; Tang et al., 2016).

2.2 Sample collection and cultivation

Thirty-eight genotypes of water spinach (*I. aquatica* F.) were obtained from a local seed market in Hangzhou, China. Their genus, origin, and characteristics are listed in Table 2. Split plot design was used with three replicates for major plots and genotypes for the minor plots. Each minor plot was 10 m². Approximately 900 seeds of each genotype were soaked overnight in aerated deionized water at (23±0.3) °C, disinfected with 0.7 g/L NaClO for 30 min, drained and sown in the field in June 2014. Planting density and field management were the same as in conventional farming practice.

Table 1 Physicochemical properties of soil in the field experiment

pH	Organic matter (g/kg)	Total N (g/kg)	Total P (g/kg)	Total K (g/kg)	Available N (g/kg)	Available P (g/kg)
5.39±0.05	23.07±0.69	1.22±0.03	0.60±0.07	1.69±0.14	97.35±6.74	77.59±10.11
Available K (g/kg)	Total Cd (g/kg)	Total Pb (g/kg)	DTPA Cd (g/kg)	DTPA Pb (g/kg)	Nitrate-N (g/kg)	
296.87±5.06	1.10±0.18	32.66±0.33	0.23±0.01	0.96±0.12	235.21±6.64	

DTPA: diethylene triamine pentaacetic acid. Reprinted from Tang et al. (2016), Copyright 2016, with permission from Elsevier

Table 2 Tested genotypes of water spinach

Serial No.	Accession name	Origin*	Horticultural characteristics	
			Petioles	Blades
1	<i>JXDY</i>	Jiangxi	Green	Extra-big
2	<i>JADY</i>	Jiangxi	Green	Big
3	<i>GDLY</i>	Jiangxi	White	Small
4	<i>GZTB</i>	Guangdong	White	Extra-big
5	<i>G501</i>	Guangdong	Light yellow	Small
6	<i>GZQL</i>	Hong Kong	Green	Small
7	<i>XGDB</i>	Fujian	White	Big
8	<i>G268</i>	Fujian	Green	Small
9	<i>TWBD</i>	Jiangxi	White	Big
10	<i>TWLY</i>	Thailand	Green	Small
11	<i>TWBL</i>	Guangdong	White	Small
12	<i>TWYX</i>	Taiwan	Green	Small
13	<i>TWZY</i>	Indonesia	Green	Small
14	<i>T311</i>	Taiwan	White	Small
15	<i>TAIG</i>	Thailand	Green	Small
16	<i>TGLY</i>	Thailand	Green	Small
17	<i>T221</i>	Guangxi	Green	Small
18	<i>TGJY</i>	Thailand	Green	Small
19	<i>TGLL</i>	Jiangsu	Green	Small
20	<i>YXBL</i>	Guangdong	White	Small
21	<i>X606</i>	Guangdong	Light green	Small
22	<i>Y601</i>	Beijing	Green	Small
23	<i>CHFE</i>	Hebei	Green	Extra-big
24	<i>QCUI</i>	Tianjin	Green	Extra-big
25	<i>XIAO</i>	Jiangsu	Green	Small
26	<i>LZHU</i>	Thailand	Green	Small
27	<i>GDZY</i>	Thailand	Green	Small
28	<i>GLCQ</i>	Jiangxi	Green	Small
29	<i>DAYE</i>	Jiangxi	Green	Extra-big
30	<i>BQBA</i>	Fujian	Green	Small
31	<i>CBDY</i>	Jiangxi	White	Extra-big
32	<i>LIUY</i>	Jiangxi	Green	Small
33	<i>BGDY</i>	Hubei	White	Big
34	<i>LYQG</i>	Fujian	Green	Small
35	<i>HSZY</i>	Shanghai	Green	Small
36	<i>B888</i>	Jiangxi	White	Small
37	<i>DQGU</i>	Jiangxi	Light green	Small
38	<i>QGZY</i>	Hebei	Green	Small

* Region (province or city of China) or country (Thailand, Indonesia)

2.3 Fresh weight and biomass determination

The experiment was performed according to the previous method (Tang et al., 2017). All genotypes were grown in green house for four weeks without application of fertilizer. Plants excised at 0.5 cm above the base were used for fresh weight (FW) and biomass determination. Mature and immature leaves (leaf blades and petioles) were separated and FWs were estimated. After rinsing with tap water, ten representative plants selected from each genotype were combined together to make a composite sample. Subsamples were dried in an oven at 105 °C for 30 min, and then at 65 °C until a constant weight was attained and the biomass of shoots was then recorded.

2.4 Heavy metal analysis and nitrate determination of plant samples

The concentrations of Cd and other metals (K, Ca, Mg, Fe, Zn, Cu, Pb, Mn, and Se) of plant samples, which were digested with HNO₃ and HClO₄ (5:1, v/v), were determined using inductively coupled plasma mass spectrometer (ICP-MS, 7500a, Aligent, USA), according to the previous method (Bao, 2008). Fresh samples were placed in 10 ml deionized water and heated in a boiling water bath for 30 min. Then the nitrate concentration in the water extract was determined using an ultraviolet spectrophotometer (Lambda 350V-vis, Perkin Elmer, Singapore), according to salicylic acid colorimetric method described by Li (2000). Analyses of other mineral elements (N and P) and nutritional indices (chlorophyll, protein, vitamin C, and cellulose) were performed according to the pervious methods (Tang et al., 2016).

2.5 Statistical analysis

The statistical evaluation was represented as the mean±standard error (SE) of four replicates. The differences between 38 genotypes of water spinach

were estimated by the least significant difference (LSD) method and the correlation was analyzed by Bivariate analysis (SPSS 20.0) and graphical representation was generated by Origin Pro 8.0.

3 Results

3.1 Fresh weight and biomass yields

FWs and biomass of 38 water spinach genotypes were determined (Figs. 1 and 2) after growth for four weeks. FW ranged from 16.62 g/plant (*LIUY*) to

10.22 g/plant (*BQBA*), approximately a 1.6-fold difference between the highest and the lowest levels with a mean value of 13.30 g/plant (Fig. 1). Biomass values ranged from 1.24 g/plant (*LIUY*) to 0.74 g/plant (*BQBA*), approximately a 1.7-fold difference with a mean value of 0.98 g/plant (Fig. 2).

3.2 Cd and nitrate uptake

Cd uptake ranged from 2.75 mg/kg dry weight (DW) (*GDZY*) to 0.07 mg/kg DW (*TWBL*), a difference of more than 39-fold with a mean value of 1.15 mg/kg DW (Fig. 3). Nitrate concentrations




Fig. 1 Shoot fresh weights of 38 water spinach genotypes grown in co-contaminated soils
 Bars represent standard error of the mean with four replicates. Data analysis was performed using LSD method




Fig. 2 Shoot biomass of 38 water spinach genotypes grown in co-contaminated soils
 Bars represent standard error of the mean with four replicates. Data analysis was performed using LSD method

among all genotypes ranged from 10983 mg/kg FW (*LIUY*) to 1809 mg/kg FW (*JXDY*), a 6-fold difference with a mean value of 5177 mg/kg FW (Fig. 4).

3.3 Correlations between Cd, nitrate, and other elements

No correlation was observed between Cd and nitrate concentrations (Fig. 5a). However, there was a significant positive correlation between Cd, Pb, and Zn

concentrations (Fig. 6) and a significant negative correlation between Cd or Pb and P (Figs. 7a and 7b), and between Zn and P (Fig. 7c). There was a positive correlation between nitrate and chlorophyll concentrations (Fig. 8b), and a significant positive correlation between nitrate and vitamin C (Fig. 8a). However, no significant correlation between Cd, nitrate and biomass yield in water spinach (Figs. 5b and 5c) was observed.




Fig. 3 Cd concentration of 38 water spinach genotypes grown in co-contaminated soils

Bars represent standard error of the mean with four replicates. Data analysis was performed using LSD method




Fig. 4 Nitrate concentration in 38 water spinach genotypes grown in co-contaminated soils

Bars represent standard error of the mean with four replicates. Data analysis was performed using LSD method




Fig. 5 Correlation coefficients between Cd, nitrate concentrations, and plant biomass of water spinach genotypes in Cd and nitrate contaminated soils

(a) Nitrate vs. Cd; (b) Plant biomass vs. Cd; (c) Plant biomass vs. nitrate




Fig. 6 Correlation coefficients between Cd, Pb, and Zn concentrations in water spinach genotypes in Cd and nitrate contaminated soils

(a) Pb vs. Cd; (b) Zn vs. Cd; (c) Zn vs. Pb. * Significance at $P<0.05$; ** Significance at $P<0.01$




Fig. 7 Correlation coefficients between P concentrations and Cd, Pb, and Zn accumulation levels in water spinach genotypes in Cd and nitrate contaminated soil

(a) Cd vs. P; (b) Pb vs. P; (c) Zn vs. P. * Significance at $P<0.05$; ** Significance at $P<0.01$




Fig. 8 Correlation coefficients between nitrate concentrations and vitamin C, chlorophyll accumulation levels in water spinach genotypes in Cd and nitrate contaminated soil

(a) Vitamin C vs. nitrate; (b) Chlorophyll vs. nitrate.
* Significance at $P < 0.05$; ** Significance at $P < 0.01$

4 Discussion

Water spinach is a common leafy vegetable which provides rich nutrients to the diet. All the selected water spinach showed successive growth in Cd and nitrate co-contaminated soil. The FW or biomass yield is close to the mean value. Water spinach tolerates high Cd and nitrate stress relatively well. The results were consistent with previous studies (Wang et al., 2007). Since there is no visible symptom of toxicity when water spinach is grown in Cd and nitrate co-contaminated soils, the potential health risk may be high from inadvertently consuming contaminated water spinach. The National Food Safety Standard of China GB 2762-2012 (Ministry of Health of the People's Republic of China, 2012) sets the safe limit for Cd contamination of fresh leafy vegetables at 0.05 mg/kg FW. Since the water content of water spinach is 93%, it can be calculated that the safe

consumption level of Cd in water spinach shoot is 0.71 mg/kg DW. Given this standard, 17 water spinach genotypes tested in our study can be safe for consumption. The maximum permissible concentration (MPC) of nitrate is 3100 mg/kg FW (Zhou et al., 2000), which means that only five water spinach genotypes can be safe for consumption. Using the combined standards for Cd and nitrate, four genotypes were recognized as safe, i.e. *JXDY*, *GZQL*, *XGDB*, and *B888* (Table 3), which are low Cd and nitrate co-accumulators and suitable to grow in slightly or moderately contaminated soils without any risk to human health.

Table 3 Cd and nitrate concentrations in the shoots of four safe water spinach genotypes

Genotypes	Cd (mg/kg DW)	Nitrate (mg/kg FW)
<i>JXDY</i>	0.50±0.08	1809.6±96.4
<i>GZQL</i>	0.11±0.02	2154.9±236.2
<i>XGDB</i>	0.41±0.04	2723.7±443.6
<i>B888</i>	0.57±0.03	2549.3±668.0

Data are expressed as mean±standard error with four replicates

Previous studies showed that low doses of heavy metals may promote plant hormone secretion and regulate plant growth and development (Liu et al., 2010). However, the distinct tolerance mechanisms of water spinach genotypes to Cd are not fully understood. Variations in Cd accumulation among genotypes may be related to uptake, transfer, and bioaccumulation of Cd in shoots (Uraguchi et al., 2009) and roots (Lux et al., 2011) and are influenced by several environmental factors such as light, cold, and humidity (Cheng et al., 2006; Li et al., 2015), including properties of the rhizosphere (Arao et al., 2009; Zheng and Zhang, 2011).

The accumulation of Cd in plants varies greatly not only among plant species but also among genotypes or cultivars within the same species (Yu et al., 2006; Liu et al., 2007; Wang et al., 2007; Martin et al., 2012). The results of the present study showed that different genotypes of water spinach had different Cd retention abilities, resulting in different Cd accumulations. Genotype-dependent Cd accumulation of water spinach largely depends on bioprocesses occurring in shoots and roots (Xin et al., 2013a). Some evidence indicates that Cd subcellular distribution may be associated with Cd tolerance and detoxification in

plants. Distribution of Cd in plants is influenced by cross-membrane transport systems, the existence of intracellular binding sites, vacuole sequestration, xylem and phloem transport, and root retention (Grant et al., 2008). In most cases heavy metals are retained in plant root cell wall, restricting their translocation to shoot and minimizing plant damage (Wang Y et al., 2015). Similar results were reported by Xin et al. (2013a, 2013b) and Huang et al. (2016). Retention of Cd in the cell wall is the effective and most important mechanism for Cd detoxification in all water spinach tissues, especially in young leaves (Xin et al., 2013b), though this may vary between genotypes (Huang et al., 2016). The presence of thicker phloem and outer cortex cell walls in the low Cd cultivars may explain why low Cd cultivar roots were able to retain more Cd, thus reducing Cd translocation to shoots (Xin et al., 2013b).

Nitrate is taken up by plants from fertilizer, but this differs depending on the genotype and environment (Burns et al., 2011b). Nitrate concentration in plants depends mainly on environmental, genetic, and nutritional factors (Anjana et al., 2009). When plants are provided with excess nitrate, only a small portion taken up by roots may be immediately translocated to and assimilated by shoots, while the majority of the absorbed nitrate is stored in vacuoles in both roots and shoots (Luo et al., 2006). Nitrate accumulation in plant tissues acts not only as a temporary store, but also as a replacement osmoticum for other plant solutes for maintaining turgor and driving leaf expansion (Burns et al., 2010). Excessive N accumulation in plant tissues is attributed to the imbalance between uptake and assimilation (Cárdenas-Navarro et al., 1999). As a result, the uptake of nitrate may reduce osmotic pressure that limits the storage of organic solutes in vacuole (Wojciechowska and Kolton, 2014).

Although there is strong evidence for the genotypical effects on Cd or nitrate accumulation in vegetable plants (Arao et al., 2003; Burns et al., 2011a; Sugiyama et al., 2011; Wang X et al., 2015; Kristl et al., 2016), less is known about the higher specificity. The present study indicated that no correlation occurred between plant Cd, nitrate, and biomass, suggesting that yield, Cd and nitrate levels may be independent of each other. This suggests that these three traits could be improved separately or in combination for high yield with low concentrations of Cd and nitrate.

Heavy metals, such as Cd, Pb, and Zn, are phytotoxic when present in excessive amounts and can interfere with photosynthetic and respiratory activities, mineral nutrition, enzymatic activity, membrane functions, and hormone balance (Clijsters and van Assche, 1985). The significant positive correlation between Cd or Pb and Zn in water spinach genotypes may indicate that they have similar uptake mechanisms. Similar results were reported in previous studies (Wu and Zhang, 2002; Liu et al., 2003; Dong et al., 2006; Li et al., 2015). These heavy metals are probably transported by similar transporters in the form of compounds or chelate complexes, and the mobilizing function of root exudates is effective not only for Cd but also for Pb and Zn (Kabata-Pendias, 2011). Cohen et al. (1998) reported that *IRT1* may facilitate the transport of heavy metals in the form of divalent cations such as Cd^{2+} , Pb^{2+} , and Zn^{2+} . Almost all Cd hyper-accumulation plants could accumulate high concentrations of Pb and Zn (He et al., 2002). All of these suggest that these three metals may have similar transport mechanisms, although the interaction of Cd with some nutrients in soil is not fully understood.

P is a macronutrient and has an important role in plant physiology including metabolism. When P is deficient, plant growth and crop yield are reduced, as P is an essential element for the synthesis of nucleic acids, phospholipids, and adenosine triphosphate (ATP) (Yin et al., 2016). Our results indicated that there was a significant negative correlation between Cd, Pb, or Zn and P concentration in water spinach plants (Fig. 7). Similar results have also been reported (Keller and Römer, 2001; Zhang et al., 2002; Dheri et al., 2007). Application of P-containing materials influences the bioavailability of heavy metals such as Cd, Pb, and Zn in soil (Qiu et al., 2011). Jiang et al. (2007) reported that increased P in soil resulted in substantial precipitation of heavy metal-P complexes in the cell wall and vacuoles in maize (*Zea mays* L.), and a similar effect was reported in strawberry (*Fragaria ananassa* D.) (Nuzhath et al., 2013). P-heavy metal interactions reduce the availability of heavy metals in soil and limit their mobility in plants (Clemens, 2006). The supply of adequate and balanced mineral nutrients to crops has the potential to improve plant tolerance, growth, development, and productivity under stress environments (Mitchell et al., 2000; Astolfi et al.,

2004). However, the proportion of macronutrient fertilizers in agriculture is often out of balance in China. Our present study points out that increasing the P supply to heavy metal (Cd, Pb, or Zn)-contaminated soils reduces Cd, Pb, or Zn concentration in water spinach.

Chlorophyll concentration is influenced by environmental factors and N fertilization (Barickman and Kopsell, 2016) and the form and ratio of N in plants (Borowski and Michalek, 2008). Previous research demonstrated that higher ratios of NO_3^- -N to NH_4^+ -N positively influenced chlorophyll concentrations in the leaf tissue of kale (*Brassica oleracea* L. var. *acephala*) (Kopsell et al., 2007) and leaf lettuce (*L. sativa* L.) (Stagnari et al., 2015). Chlorophyll concentration in low nitrate accumulation genotypes oilseed rape (*Brassica napus* L.) was reported to be significantly lower than that in high nitrate accumulation genotypes (Han et al., 2016). In the present study, there was a positive correlation between nitrate and chlorophyll concentration (Fig. 8b), which was consistent with previous studies. There was a significant positive correlation between nitrate and vitamin C in water spinach genotypes (Fig. 8a). Similar results in spinach (*Spinacia oleracea* L.) were previously reported by Conesa et al. (2009) and Koh et al. (2012). The levels of vitamin C and nitrate in leafy vegetables are two key indices for evaluating their nutritional quality (Konstantopoulou et al., 2010), and vitamin C content in leafy vegetables is mainly controlled by the form and ratio of N (Sørensen et al., 1994). The significant positive correlation between nitrate and vitamin C concentration in water spinach genotypes suggests that it would be difficult to reduce nitrate and increase vitamin C concentration simultaneously, so it is necessary to improve these two traits separately.

5 Conclusions

We have identified four genotypes of water spinach, *JXDY*, *GZQL*, *XGDB*, and *B888*, as low co-accumulators for Cd and nitrate, indicating that they should be the preferred genotypes for growing in slightly or moderately contaminated soils, minimizing risk to human health. It should be possible to combine high yields with low concentrations of Cd and nitrate as these variables are independent of each

other. Increasing P fertilizer rates appears to increase tolerance in water spinach to Cd, Pb, and Zn toxicity and nitrate concentrations. Our experiment has shown that screening vegetable genotypes for those which do not exceed allowable levels of a contaminant is a cost-effective strategy for minimizing the risk of contaminants to human health via the food chain.

Compliance with ethics guidelines

Lin TANG, Wei-jun LUO, Zhen-li HE, Hanumanth Kumar GURAJALA, Yasir HAMID, Kiran Yasmin KHAN, and Xiao-e YANG declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

- Anjana, Umar S, Iqbal M, 2009. Factors responsible for nitrate accumulation: a review. In: Lichtfouse E, Navarrete M, Debaeke P, et al. (Eds.), Sustainable Agriculture. Springer, Dordrecht, p.533-549.
https://doi.org/10.1007/978-90-481-2666-8_33
- Arao T, Ae N, Sugiyama M, et al., 2003. Genotypic differences in cadmium uptake and distribution in soybeans. *Plant Soil*, 251(2):247-253.
<https://doi.org/10.1023/A:1023079819086>
- Arao T, Kawasaki A, Baba K, et al., 2009. Effects of water management on cadmium and arsenic accumulation and dimethylarsinic acid concentrations in Japanese rice. *Environ Sci Technol*, 43(24):9361-9367.
<https://doi.org/10.1021/es9022738>
- Astolfi S, Zuchi S, Passera C, 2004. Role of sulphur availability on cadmium-induced changes of nitrogen and sulphur metabolism in maize (*Zea mays* L.) leaves. *J Plant Physiol*, 161(7):795-802.
<https://doi.org/10.1016/j.jplph.2003.11.005>
- Bao SD, 2008. Soil agricultural Chemistry Analysis Method, 3rd Ed. China Agriculture Press, Beijing, China (in Chinese).
- Barickman TC, Kopsell DA, 2016. Nitrogen form and ratio impact Swiss chard (*Beta vulgaris* subsp. *cicla*) shoot tissue carotenoid and chlorophyll concentrations. *Sci Hortic*, 204:99-105.
<https://doi.org/10.1016/j.scienta.2016.04.007>
- Borowski E, Michalek S, 2008. The effect of nitrogen form and air temperature during foliar fertilization on gas exchange, the yield and nutritive value of spinach (*Spinacia oleracea* L.). *Folia Hortic*, 20(2):17-27.
<https://doi.org/10.2478/fhort-2013-0110>
- Burns IG, Zhang KF, Turner MK, et al., 2010. Iso-osmotic regulation of nitrate accumulation in lettuce. *J Plant Nutr*, 34(2):283-313.
<https://doi.org/10.1080/01904167.2011.533328>
- Burns IG, Zhang KF, Turner MK, et al., 2011a. Screening for genotype and environment effects on nitrate accumulation in 24 species of young lettuce. *J Sci Food Agric*, 91(3):553-562.
<https://doi.org/10.1002/jsfa.4220>

- Burns IG, Zhang KF, Turner MK, et al., 2011b. Genotype and environment effects on nitrate accumulation in a diversity set of lettuce accessions at commercial maturity: the influence of nitrate uptake and assimilation, osmotic interactions and shoot weight and development. *J Sci Food Agric*, 91(12):2217-2233.
<https://doi.org/10.1002/jsfa.4442>
- Cárdenas-Navarro R, Adamowicz S, Robin P, 1999. Nitrate accumulation in plants: a role for water. *J Exp Bot*, 50(334):613-624.
<https://doi.org/10.1093/jexbot/50.334.613>
- Chen XP, Cui ZL, Fan MS, et al., 2014. Producing more grain with lower environmental costs. *Nature*, 514(7523):486-489.
<https://doi.org/10.1038/nature13609>
- Chen Y, Li TQ, Han X, et al., 2012. Cadmium accumulation in different pakchoi cultivars and screening for pollution-safe cultivars. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)*, 13(6):494-502.
<https://doi.org/10.1631/jzus.B1100356>
- Cheng WD, Zhang GP, Yao HG, et al., 2006. Genotypic and environmental variation in cadmium, chromium, arsenic, nickel, and lead concentrations in rice grains. *J Zhejiang Univ-Sci B*, 7(7):565-571.
<https://doi.org/10.1631/jzus.2006.B0565>
- Clemens S, 2006. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie*, 88(11):1707-1719.
<https://doi.org/10.1016/j.biochi.2006.07.003>
- Clijsters H, van Assche F, 1985. Inhibition of photosynthesis by heavy metal. *Phytosyn Res*, 7(1):31-40.
<https://doi.org/10.1007/BF00032920>
- Cohen CK, Fox TC, Garvin DF, et al., 1998. The role of iron-deficiency stress responses in stimulating heavy-metal transport in plants. *Plant Physiol*, 116(3):1063-1072.
<https://doi.org/10.1104/pp.116.3.1063>
- Conesa E, Ninirola D, Vicente MJ, et al., 2009. The influence of nitrate/ammonium ratio on yield quality and nitrate, oxalate and vitamin C content of baby leaf spinach and bladder campion plants grown in a floating system. *Acta Hort*, 843:269-273.
- Dheri GS, Brar MS, Malhi SS, 2007. Influence of phosphorus application on growth and cadmium uptake of spinach in two cadmium-contaminated soils. *J Plant Nutr Soil Sci*, 170(4):495-499.
<https://doi.org/10.1002/jpln.200625051>
- Dong J, Wu FB, Zhang GP, 2006. Influence of cadmium on antioxidant capacity and four microelement concentrations in tomato seedlings (*Lycopersicon esculentum*). *Chemosphere*, 64(10):1659-1666.
<https://doi.org/10.1016/j.chemosphere.2006.01.030>
- Escobar-Gutiérrez AJ, Burns IG, Lee A, et al., 2002. Screening lettuce cultivars for low nitrate content during summer and winter production. *J Hort Sci Biotechnol*, 77(2):232-237.
<https://doi.org/10.1080/14620316.2002.11511485>
- Fan SK, Zhu J, Tian WH, et al., 2017. Effects of split applications of nitrogen fertilizers on the Cd level and nutritional quality of Chinese cabbage. *J Zhejiang Univ-Sci B (Biomed & Biotechnol)*, 18(10):897-905.
<https://doi.org/10.1631/jzus.B1600272>
- Grant CA, Clarke JM, Duguid S, et al., 2008. Selection and breeding of plant cultivars to minimize cadmium accumulation. *Sci Total Environ*, 390(2-3):301-310.
<https://doi.org/10.1016/j.scitotenv.2007.10.038>
- Gulis G, Czompolyova M, Cerhan JR, 2002. An ecologic study of nitrate in municipal drinking water and cancer incidence in Trnava District, Slovakia. *Environ Res*, 88(3):182-187.
<https://doi.org/10.1006/enrs.2002.4331>
- Habermeyer M, Roth A, Guth S, et al., 2015. Nitrate and nitrite in the diet: how to assess their benefit and risk for human health. *Mol Nutr Food Res*, 59(1):106-128.
<https://doi.org/10.1002/mnfr.201400286>
- Hakeem KR, Mir BA, Qureshi MI, et al., 2013. Physiological studies and proteomic analysis for differentially expressed proteins and their possible role in the root of N-efficient rice (*Oryza sativa* L.). *Mol Breeding*, 32(4):785-798.
<https://doi.org/10.1007/s11032-013-9906-0>
- Han YL, Song HX, Liao Q, et al., 2016. Nitrogen use efficiency is mediated by vacuolar nitrate sequestration capacity in roots of *Brassica napus*. *Plant Physiol*, 170(3):1684-1698.
<https://doi.org/10.1104/pp.15.01377>
- He B, Yang XE, Ni WZ, et al., 2002. *Sedum alfredii*: a new lead-accumulating ecotype. *Acta Bot Sin*, 44(11):1365-1370 (in Chinese).
- Huang BF, Xin JL, Dai HW, et al., 2015. Identification of low-Cd cultivars of sweet potato (*Ipomoea batatas* (L.) Lam.) after growing on Cd-contaminated soil: uptake and partitioning to the edible roots. *Environ Sci Pollut Res*, 22(15):11813-11821.
<https://doi.org/10.1007/s11356-015-4449-z>
- Huang YY, Shen C, Chen JX, et al., 2016. Comparative transcriptome analysis of two *Ipomoea aquatica* Forsk. cultivars targeted to explore possible mechanism of genotype-dependent accumulation of cadmium. *J Agric Food Chem*, 64(25):5241-5250.
<https://doi.org/10.1021/acs.jafc.6b01267>
- Jiang HM, Yang JC, Zhang JF, 2007. Effects of external phosphorus on the cell ultrastructure and the chlorophyll content of maize under cadmium and zinc stress. *Environ Pollut*, 147(3):750-756.
<https://doi.org/10.1016/j.envpol.2006.09.006>
- Kabata-Pendias A, 2011. Trace Elements in Soils and Plants, 4th Ed. CRC Press Inc., Boca Raton, USA.
- Keller H, Römer W, 2001. Cu, Zn, and Cd acquisition by two spinach cultivars depending on P nutrition and root exudation. *J Plant Nutr Soil Sci*, 164(3):335-342 (in German).
[https://doi.org/10.1002/1522-2624\(200106\)164:3<335::aid-jpln335>3.0.co;2-c](https://doi.org/10.1002/1522-2624(200106)164:3<335::aid-jpln335>3.0.co;2-c)
- Kirkham MB, 2006. Cadmium in plants on polluted soils: effects of soil factors, hyperaccumulation, and amendments. *Geoderma*, 137(1-2):19-32.
<https://doi.org/10.1016/j.geoderma.2006.08.024>
- Koh E, Charoenprasert S, Mitchell AE, 2012. Effect of organic

- and conventional cropping systems on ascorbic acid, vitamin C, flavonoids, nitrate, and oxalate in 27 varieties of spinach (*Spinacia oleracea* L.). *J Agric Food Chem*, 60(12):3144-3150.
<https://doi.org/10.1021/jf300051f>
- Konstantopoulou E, Kapotis G, Salachas G, et al., 2010. Nutritional quality of greenhouse lettuce at harvest and after storage in relation to N application and cultivation season. *Sci Hortic*, 125(2):93.e1-93.e5.
<https://doi.org/10.1016/j.scienta.2010.03.003>
- Kopsell DA, Kopsell DE, Curran-Celentano J, et al., 2007. Carotenoid pigments in kale are influenced by nitrogen concentration and form. *J Sci Food Agric*, 87(5):900-907.
<https://doi.org/10.1002/jsfa.2807>
- Kristl J, Ivancic A, Mergedus A, et al., 2016. Variation of nitrate content among randomly selected taro (*Colocasia esculenta* (L.) Schott) genotypes and the distribution of nitrate within a corm. *J Food Compos Anal*, 47:76-81.
<https://doi.org/10.1016/j.jfca.2016.01.007>
- Lane EA, Canty MJ, More SJ, 2015. Cadmium exposure and consequence for the health and productivity of farmed ruminants. *Res Vet Sci*, 101:132-139.
<https://doi.org/10.1016/j.rvsc.2015.06.004>
- Li HS, 2000. Principle and Technology of Plant Physiological and Biochemical Experiment. Higher Education Press, Beijing, China (in Chinese).
- Li N, Kang Y, Pan WJ, et al., 2015. Concentration and translocation of heavy metals in vegetables and risk assessment of human exposure to bioaccessible heavy metals in soil near a waste-incinerator site, South China. *Sci Total Environ*, 521-522:144-151.
<https://doi.org/10.1016/j.scitotenv.2015.03.081>
- Li XH, Zhou QX, Wei SH, et al., 2012. Identification of cadmium excluding welsh onion (*Allium fistulosum* L.) cultivars and their mechanisms of low cadmium accumulation. *Environ Sci Pollut Res*, 19(5):1773-1780.
<https://doi.org/10.1007/s11356-011-0692-0>
- Liu F, Liu XN, Ding C, et al., 2015. The dynamic simulation of rice growth parameters under cadmium stress with the assimilation of multi-period spectral indices and crop model. *Field Crops Res*, 183:225-234.
<https://doi.org/10.1016/j.fcr.2015.08.004>
- Liu JG, Li KQ, Xu JK, et al., 2003. Interaction of Cd and five mineral nutrients for uptake and accumulation in different rice cultivars and genotypes. *Field Crop Res*, 83(3):271-281.
[https://doi.org/10.1016/S0378-4290\(03\)00077-7](https://doi.org/10.1016/S0378-4290(03)00077-7)
- Liu JG, Qian M, Cai GL, et al., 2007. Uptake and translocation of Cd in different rice cultivars and the relation with Cd accumulation in rice grain. *J Hazard Mater*, 143(1-2):443-447.
<https://doi.org/10.1016/j.jhazmat.2006.09.057>
- Liu WT, Zhou QX, An J, et al., 2010. Variations in cadmium accumulation among Chinese cabbage cultivars and screening for Cd-safe cultivars. *J Hazard Mater*, 173(1-3):737-743.
<https://doi.org/10.1016/j.jhazmat.2009.08.147>
- Luo JK, Sun SB, Jia LJ, et al., 2006. The mechanism of nitrate accumulation in pakchoi [*Brassica campestris* L.ssp. *Chinensis* (L.)]. *Plant Soil*, 282(1-2):291-300.
<https://doi.org/10.1007/s11104-005-6094-7>
- Lux A, Martinka M, Vaculik M, et al., 2011. Root responses to cadmium in the rhizosphere: a review. *J Exp Bot*, 62(1):21-37.
<https://doi.org/10.1093/jxb/erq281>
- Martin SR, Llugany M, Barceló J, et al., 2012. Cadmium exclusion a key factor in differential Cd-resistance in *Thlaspi arvense* ecotypes. *Biol Plant*, 56(4):729-734.
<https://doi.org/10.1007/s10535-012-0056-8>
- Ministry of Health of the People's Republic of China, 2012. GB 2762-2012: National Food Standard. Maximum Levels of Contaminants in Foods.
- Mitchell LG, Grant CA, Racz GJ, 2000. Effect of nitrogen application on concentration of cadmium and nutrient ions in soil solution and in durum wheat. *Can J Soil Sci*, 80(1):107-115.
<https://doi.org/10.4141/S98-085>
- Nuzahath A, Abdukadir A, Dilmur M, 2013. Effect of phosphorus on chemical forms and physiological properties of cadmium in *Fragaria ananassa* D. *J Soil Sci*, 44(6):1460-1464 (in Chinese).
- Qiu Q, Wang Y, Yang Z, et al., 2011. Effects of phosphorus supplied in soil on subcellular distribution and chemical forms of cadmium in two Chinese flowering cabbage (*Brassica parachinensis* L.) cultivars differing in cadmium accumulation. *Food Chem Toxicol*, 49(9):2260-2267.
<https://doi.org/10.1016/j.fct.2011.06.024>
- Sharma A, Sainger M, Dwivedi S, et al., 2010. Genotypic variation in *Brassica juncea* (L.) Czern. cultivars in growth, nitrate assimilation, antioxidant responses and phytoremediation potential during cadmium stress. *J Environ Biol*, 31(5):773-780.
- Sørensen JN, Johansen AS, Poulsen N, 1994. Influence of growth conditions on the value of crisphead lettuce. 1. Marketable and nutritional quality as affected by nitrogen supply, cultivar and plant age. *Plant Foods Hum Nutr*, 46(1):1-11.
<https://doi.org/10.1007/BF01088455>
- Stagnari F, Galièni A, Pisante M, 2015. Shading and nitrogen management affect quality, safety and yield of greenhouse-grown leaf lettuce. *Sci Hortic*, 192:70-79.
<https://doi.org/10.1016/j.scienta.2015.05.003>
- Sugiyama M, Ae N, Hajika M, 2011. Developing of a simple method for screening soybean seedling cadmium accumulation to select soybean genotypes with low seed cadmium. *Plant Soil*, 341(1-2):413-422.
<https://doi.org/10.1007/s11104-010-0654-1>
- Tang L, Luo WJ, Tian SK, et al., 2016. Genotypic differences in cadmium and nitrate co-accumulation among the Chinese cabbage genotypes under field conditions. *Sci Hortic*, 201:92-100.
<https://doi.org/10.1016/j.scienta.2016.01.040>
- Tang L, Luo WJ, Chen WK, et al., 2017. Field crops (*Ipomoea aquatica* Forsk. and *Brassica chinensis* L.) for phytoremediation of cadmium and nitrate co-contaminated soils

- via rotation with *Sedum alfredii* Hance. *Environ Sci Pollut Res*, 24(23):19293-19305.
<https://doi.org/10.1007/s11356-017-9146-7>
- Uraguchi S, Mori S, Kuramata M, et al., 2009. Root-to-shoot Cd translocation via the xylem is the major process determining shoot and grain cadmium accumulation in rice. *J Exp Bot*, 60(9):2677-2688.
<https://doi.org/10.1093/jxb/erp119>
- Wang GL, Ding GD, Li L, et al., 2014. Identification and characterization of improved nitrogen efficiency in interspecific hybridized new-type *Brassica napus*. *Ann Bot*, 114(3):549-559.
<https://doi.org/10.1093/aob/mcu135>
- Wang JL, Fang W, Yang ZY, et al., 2007. Inter- and intraspecific variations of cadmium accumulation of 13 leafy vegetable species in a greenhouse experiment. *J Agric Food Chem*, 55(22):9118-9123.
<https://doi.org/10.1021/jf0716432>
- Wang JL, Yuan JG, Yang ZY, et al., 2009. Variation in cadmium accumulation among 30 cultivars and cadmium subcellular distribution in 2 selected cultivars of water spinach (*Ipomoea aquatica* Forsk.). *J Agric Food Chem*, 57(19):8942-8949.
<https://doi.org/10.1021/jf900812s>
- Wang X, Shi Y, Chen X, et al., 2015. Screening of Cd-safe genotypes of Chinese cabbage in field condition and Cd accumulation in relation to organic acids in two typical genotypes under long-term Cd stress. *Environ Sci Pollut Res*, 22(21):16590-16599.
<https://doi.org/10.1007/s11356-015-4838-3>
- Wang Y, Shen H, Xu L, et al., 2015. Transport, ultrastructural localization and distribution of chemical forms of lead in radish (*Raphanus sativus* L.). *Front Plant Sci*, 6:293.
<https://doi.org/10.3389/fpls.2015.00293>
- Wojciechowska R, Kołton A, 2014. Comparison of the ability of fifteen onion (*Allium cepa* L.) cultivars to accumulate nitrates. *Acta Agrobot*, 67(1):27-32.
<https://doi.org/10.5586/aa.2014.006>
- Wu FB, Zhang G, 2002. Genotypic differences in effect of Cd on growth and mineral concentrations in barley seedlings. *Bull Environ Contam Toxicol*, 69(2):219-227.
<https://doi.org/10.1007/s00128-002-0050-5>
- Xin JL, Huang BF, Yang ZY, et al., 2010. Responses of different water spinach cultivars and their hybrid to Cd, Pb and Cd-Pb exposures. *J Hazard Mater*, 175(1-3):468-476.
<https://doi.org/10.1016/j.jhazmat.2009.10.029>
- Xin JL, Huang BF, Yang JZ, et al., 2013a. Role of roots in cadmium accumulation of two water spinach cultivars: reciprocal grafting and histochemical experiments. *Plant Soil*, 366(1-2):425-432.
<https://doi.org/10.1007/s11104-012-1439-5>
- Xin JL, Huang BF, Yang ZY, et al., 2013b. Comparison of cadmium subcellular distribution in different organs of two water spinach (*Ipomoea aquatica* Forsk.) cultivars. *Plant Soil*, 372(1-2):431-444.
<https://doi.org/10.1007/s11104-013-1729-6>
- Xu GH, Fan XR, Miller AJ, 2012. Plant nitrogen assimilation and use efficiency. *Annu Rev Plant Biol*, 63:153-182.
<https://doi.org/10.1146/annurev-arplant-042811-105532>
- Yin AG, Yang ZY, Ebbs S, et al., 2016. Effects of phosphorus on chemical forms of Cd in plants of four spinach (*Spinacia oleracea* L.) cultivars differing in Cd accumulation. *Environ Sci Pollut Res*, 23(6):5753-5762.
<https://doi.org/10.1007/s11356-015-5813-8>
- Yu H, Wang JL, Fang W, et al., 2006. Cadmium accumulation in different rice cultivars and screening for pollution-safe cultivars of rice. *Sci Total Environ*, 370(2-3):302-309.
<https://doi.org/10.1016/j.scitotenv.2006.06.013>
- Zhang GP, Fukami M, Sekimoto H, 2002. Influence of cadmium on mineral concentrations and yield components in wheat genotypes differing in Cd tolerance at seedling stage. *Field Crop Res*, 77(2-3):93-98.
[https://doi.org/10.1016/S0378-4290\(02\)00061-8](https://doi.org/10.1016/S0378-4290(02)00061-8)
- Zheng SN, Zhang MK, 2011. Effect of moisture regime on the redistribution of heavy metals in paddy soil. *J Environ Sci*, 23(3):434-443.
[https://doi.org/10.1016/S1001-0742\(10\)60428-7](https://doi.org/10.1016/S1001-0742(10)60428-7)
- Zhou ZY, Wang MJ, Wang JS, 2000. Nitrate and nitrite contamination in vegetables in China. *Food Rev Int*, 16(1):61-76.
<https://doi.org/10.1081/fri-100100282>

中文概要

题目: 基于筛选安全的共低积累基因型评价空心菜对镉和硝酸盐积累的变异

目的: 筛选镉-硝酸盐共低积累空心菜基因型, 并研究降低空心菜重金属含量, 提高营养品质的农艺措施。

创新点: 首次筛选得到镉-硝酸盐共低积累空心菜基因型, 并研究空心菜可食部污染物、矿质元素和营养指标之间的相关性, 提出进一步降低空心菜可食部镉和硝酸盐含量的农艺措施。

方法: 共 38 个空心菜基因型收集于世界各地, 种植在连作了 7 年的中度镉-硝酸盐复合污染土壤上 (Cd 1.10 mg/kg, NO₃⁻ 235.2 mg/kg), 4 周后收获。用 HNO₃-HClO₄ (体积比 5:1) 消煮, 电感耦合等离子体质谱仪 (ICP-MS) 测定各种金属元素, 水杨酸-硫酸比色法测定硝酸盐含量, 钒钼黄比色法测定磷含量, 2,6-二氯喹啉滴定法测定维生素 C 含量, 乙醇-丙酮 (体积比 2:1) 比色法测定叶绿素含量。

结论: 本试验筛选得到镉-硝酸盐共低积累空心菜基因型 4 个 (Cd < 0.71 mg/kg DW, NO₃⁻ < 3100 mg/kg FW), 分别是 *JXDY*、*GZQL*、*XGDB* 和 *B888*, 可以在中轻度镉-硝酸盐复合污染土壤上安全生产。空心菜地上部镉与铅、锌含量呈正相关, 而这 3 种元素均与磷量呈负相关。这些结果表明镉、铅和锌通过相同的途径被空心菜吸收, 可以同时被治理。增加磷肥供应率可以抑制复合污染土壤中的镉和硝酸盐向空心菜可食部的转移。

关键词: 基因型变异; 重金属; 硝酸盐; 土壤污染; 空心菜