



## Review

<https://doi.org/10.1631/jzus.B2400406>

# Emerging role of the metabolite succinate in bone-related diseases

Zuping WU\*, Qiaoli DAI\*, Ying WANG, Na WU, Chenyu WANG, Jiejun SHI✉

*Stomatology Hospital, School of Stomatology, Zhejiang University School of Medicine, Clinical Research Center for Oral Diseases of Zhejiang Province, Key Laboratory of Oral Biomedical Research of Zhejiang Province, Cancer Center of Zhejiang University, Hangzhou 310016*

**Abstract:** Bone-related diseases, including osteoporosis (OP), osteoarthritis (OA), rheumatoid arthritis (RA), fractures, and periodontitis, significantly impact human health. Succinate, primarily known as a metabolic intermediate in the tricarboxylic acid (TCA) cycle, has emerged as a regulator of cellular functions beyond its metabolic role. Under stress, succinate accumulates in mitochondria and acts as a signaling molecule, modulating cellular processes. Notably, succinate activates angiogenesis and inflammation by stabilizing hypoxia-inducible factor 1 alpha (HIF-1 $\alpha$ ). Moreover, it influences various pathophysiological processes by interacting with the succinate receptor (SUCNR-1), thereby impacting immune response, inflammation, cancer metastasis, and bone homeostasis. The multifaceted roles of succinate as a signaling molecule vary depending on its cellular location and concentration. Recent metabolomic analyses have revealed elevated succinate levels in bone-related diseases, indicating its potential association with these conditions. The objective of this review is to elucidate the impact of succinate on different bone-related diseases and to discuss potential therapeutic targets and drug molecules based on its mechanisms of action.

**Key words:** Succinate; Osteoarthritis; Rheumatoid arthritis; Osteoporosis; Periodontitis

## 1 Introduction

Various bone-related diseases pose a significant threat to human health due to their high incidence and disability rates; they severely limit patients' mobility and decrease their quality of life. The occurrence and development of these diseases are often associated with factors such as bone homeostasis, inflammation, and immunity. Succinate is a metabolic intermediate of the tricarboxylic acid (TCA) cycle, which can be oxidized to fumarate by means of succinate dehydrogenase (SDH) catalysis, which is crucial for adenosine triphosphate (ATP) production in mitochondria. However, recent studies have increasingly revealed regulatory roles of succinate beyond its activity as a metabolic intermediate. Under physiological or pathological stress, disruptions in the TCA cycle may lead to accumulation of succinate in mitochondria and its secretion into the cytoplasm as a signaling molecule to regulate cellular functions. In particular, by inhibiting prolyl hydroxylase (PHD) activity and stabilizing the expression of hypoxia-inducible factor 1 alpha (HIF-1 $\alpha$ ), succinate activates downstream signaling pathways, promoting angiogenesis and inflammation. Additionally, succinate can be secreted into the extracellular space and mediate a variety of pathophysiological processes by interacting with succinate receptor 1 (SUCNR-1). These processes include immune response, inflammation, cancer metastasis, and bone homeostasis. Succinate also plays a crucial role in energy metabolism by inducing protein

---

✉ Jiejun SHI, [sjiejun@zju.edu.cn](mailto:sjiejun@zju.edu.cn)

\* The two authors contributed equally to this work

Jiejun SHI, <https://orcid.org/0000-0001-6722-2909>

Zuping WU, <https://orcid.org/0000-0003-3661-8593>

Qiaoli DAI, <https://orcid.org/0009-0005-3273-525x>

post-translational modifications (PTMs) through succinylation, thereby regulating the activity of various metabolic enzymes. It has been found to be involved in the development of many diseases, such as ischemia/reperfusion injury, myocardial infarction, liver fibrosis, obesity, hypertension, neuroendocrine tumors, and hepatocellular carcinoma (Rubic et al., 2008; Hakak et al., 2009; Hogberg et al., 2011; Pell et al., 2016). Metabolites are the ultimate functional products that integrate external stimuli with intracellular signals (Peng et al., 2015). In recent years, elevated levels of succinate have been observed in metabolomic analyses of bone-related diseases such as osteoporosis (OP), osteoarthritis (OA), rheumatoid arthritis (RA), and periodontitis, suggesting an association between succinate and bone-related diseases.

A comprehensive understanding of the role of succinate in bone-related diseases is crucial for effectively targeting succinate and related factors as therapeutic interventions to influence the progression of bone-related diseases. This article elucidates the potential roles of succinate in the bone-related diseases listed above. Furthermore, based on its mechanisms of action, we discuss potential therapeutic targets and related drug molecules.

## 2 Source and structure of succinate and its receptors

### 2.1 Source of succinate

#### 2.1.1 Succinate is generated by somatic mitochondria or microbial metabolism

Succinate is typically produced in somatic mitochondria as a metabolic intermediate of the TCA cycle, and its levels generally remain stable. However, recent discoveries have identified succinate as a response molecule under stress conditions such as ischemia, hypoxia, bacterial infection, and hyperglycemia (Hohl et al., 1987; Palmieri, 2013; Valls-Lacalle et al., 2016; Wang et al., 2019). Under these conditions, succinate accumulates in mitochondria through various pathways. In addition to somatic mitochondria, succinate can also be produced through microbial metabolism. Succinate is a key metabolic product of the gut microbiota (Connors et al., 2019), which produces it by metabolizing polysaccharides and amino acids (Fernández-Veledo and Vendrell, 2019). Within the gut microbiota, the genus *Bacteroides* are the main producers of succinate. They utilize specialized carbohydrate-active enzymes to break down dietary fibers and complex polysaccharides into monosaccharide, producing intermediates such as pyruvate and oxaloacetate, which are then converted into succinate (De Vadder and Mithieux, 2018). Moreover, periodontal pathogens have also been found to produce succinate. *Porphyromonas gingivalis* can metabolize asparagine into succinate through fumarate, as an intermediate which plays an important role in its energy production and pathogenicity (Shah and Williams, 1987; Takahashi et al., 2000). Likewise, *Prevotella intermedia* utilizes glucose to produce succinate with the aid of bicarbonate (Takahashi and Yamada, 2000). Interestingly, in both gut and oral microbiota, succinate cross-feeding occurs (Fernández-Veledo and Vendrell, 2019; Kin et al., 2020), in which one microorganism produces succinate and another uses it for energy, optimizing metabolic efficiency within the microbial community.

#### 2.1.2 Four pathways of succinate accumulation under stress conditions

##### 2.1.2.1 Activity of SDH in the TCA cycle is inhibited and succinate cannot be converted to fumarate

SDH catalyzes the oxidation of succinate to fumarate, which is an important step in maintaining the normal flow of the TCA cycle. Alterations in SDH activity can impede this oxidation process, leading to succinate accumulation. Mutations in the SDH gene have been identified in various cancers (Bardella et al., 2011; Wu and Zhao, 2013; Jiang and Yan, 2017; Zeng et al., 2023), and are proven to contribute to tumorigenesis (Bardella et al., 2011). The accumulation of succinate under altered SDH activity is a potential mechanism (Dalla Pozza et al., 2020), which is associated with promoting tumor-cell survival, migration, and angiogenesis (Zhang and Lang, 2023). Additionally, SDH function relies on the oxidized forms of FAD and NAD<sup>+</sup> as cofactors. Under

hypoxic conditions, these cofactors exist in reduced form, leading to SDH dysfunction and thus elevated levels of succinate (Tannahill et al., 2013). Furthermore, oxidative stress during hyperglycemia generates a significant amount of reactive oxygen species (ROS). This can also reduce SDH activity (Guo et al., 2017).

#### 2.1.2.2 Reverse activity of SDH is activated, converting fumarate to succinate

It is known that mitochondrial ROS-induced oxidative damage is a key factor in the pathogenesis of ischemia-reperfusion injury. Studies have found that succinate is a significant driving factor in ROS production during reperfusion. In contrast to normal conditions, in which succinate mainly comes from the TCA cycle's canonical activity ( $\alpha$ -ketoglutarate  $\rightarrow$  succinyl-CoA  $\rightarrow$  succinate), succinate is primarily generated through the reverse activity of SDH under ischemic conditions. The study by Chouchani et al. first used computer simulations to reveal that under ischemic conditions, reversal of SDH is the optimal solution for succinate production. Subsequently, intravenous infusion of dimethyl malonate (a competitive inhibitor of SDH, DMM) resulted in a reduction of succinate accumulation by more than half under ischemic conditions (Chouchani et al., 2014a). In hypoxic synovial fibroblasts, the use of DMM was also observed to decrease succinate levels (Li et al., 2018). During ischemia or hypoxia, SDH's ability to oxidize succinate to fumarate is impaired due to a lack of sufficient electron acceptors such as oxygen and FAD. Fumarate can accumulate through the malate-aspartate shuttle (MAS) and the purine nucleotide cycle (PNC) during ischemia (Tomitsuka et al., 2010), and is converted back to succinate by activating the reverse activity of SDH (Pellerin and Magistretti, 1994).

#### 2.1.2.3 Glutamine (Gln)-dependent anaplerosis

The accumulation of succinate in macrophages stimulated by lipopolysaccharide (LPS) has been observed in many studies. LPS-stimulated macrophages undergo metabolic reprogramming, shifting from oxidative phosphorylation to glycolysis, which leads to decreased activity of the TCA cycle. Under these conditions, Gln-dependent anaplerosis becomes the primary source of succinate production (Tannahill, et al., 2013). It is known that Gln enters the cytoplasm through membrane transport protein. In LPS-stimulated macrophages, levels of the Gln transporter SLC3A2 significantly increase after 24 hours (Tannahill, et al., 2013). The Gln that enters the cytoplasm through this transporter is converted into glutamate (Glu) by glutaminase, and further transformed into  $\alpha$ -ketoglutarate ( $\alpha$ -KG) by glutamate dehydrogenase, replenishing the TCA cycle and thus promoting succinate production (Lee et al., 2023).

#### 2.1.2.4 Gamma-aminobutyric acid (GABA) shunt pathway

Besides the Gln-dependent anaplerosis, the GABA shunt pathway is also a major source of succinate accumulation under LPS stimulation. Tannahill et al. found through gene-expression analysis and metabolomics that LPS promotes the expression of GABA transporters (SLC6A13, SLC6A12) and generation of GABA (Tannahill, et al., 2013). Moreover, GABA-transaminase inhibitors inhibit succinate production, indicating that the GABA shunt pathway also contributes to succinate accumulation (Li, et al., 2018). In the cytosol, GABA is decarboxylated from Glu by glutamate decarboxylase (GAD). Subsequently, it enters the mitochondria via GABA transporters (SLC6A13, SLC6A12) and reacts with  $\alpha$ -KG under the action of GABA transaminase (GABA-T) to generate succinic acid semialdehyde, which is then oxidized to succinate.

The four pathways of succinate accumulation under stress conditions are shown in **Fig. 1**.

**Fig. 1 Four pathways of succinate accumulation under stress conditions.**

(1) Inhibited SDH pathway: When the activity of SDH in the TCA cycle is inhibited, succinate cannot be converted to fumarate and accumulates. (2) Reversed SDH pathway: Fumarate can accumulate through the malate-aspartate shuttle (MAS) and the purine nucleotide cycle (PNC) during ischemia, and is converted back to succinate by activating the reverse activity of SDH. (3) Gln-dependent anaplerosis: In LPS-stimulated macrophages, the Gln that enters the cytoplasm through Gln transporters is converted into Glu by glutaminase, and further transformed into  $\alpha$ -KG by glutamate dehydrogenase, replenishing the TCA cycle and thus promoting the production of succinate. (4) GABA shunt pathway: In the cytosol, GABA is decarboxylated from Glu by GAD. Subsequently, GABA enters the mitochondria via GABA transporters and reacts with  $\alpha$ -KG under the action of GABA-T to generate succinic acid semialdehyde, which is then oxidized to succinate. (Image created with BioRender.com, with permission).

## 2.2 Expression and structure of SUCNR1

Under stress conditions, accumulated succinate in mitochondria is secreted into the cytoplasm as a signaling molecule to regulate local cell metabolism to adapt to stressful environments or promote disease progression. However, continued research has uncovered the fact that succinate can also act as an extracellular signal component to mediate a variety of pathophysiological processes by binding to the succinate receptor SUCNR1.

SUCNR1 is widely expressed in human cells (Wu, 2023), with the highest expression observed in adipose tissue, followed by liver, kidney, pancreas, spleen, and small intestine tissues (Guo et al., 2020b). Additionally, SUCNR1 is expressed in immune and hematopoietic lineage cells, especially in mesenchymal stem cells (MSCs), monocytes, and macrophages (Uhlen et al., 2015; Guo, et al., 2017). SUCNR1 is closely associated with angiogenesis, bone homeostasis, and immune activation. Studies have also confirmed that there is SUCNR1 expression in human gingival fibroblasts, providing a new perspective on the pathogenesis of periodontitis.

SUCNR1 (or GPR91) is a member of the G protein-coupled receptor (GPCR) family, the largest group of membrane proteins in the human genome. Human SUCNR1 encodes a protein of 334 amino acids, which shares high homology among humans, rats, and mice. The SUCNR1 sequence primarily differs at the C-terminus, being 12 amino acids shorter in rodents (Ariza et al., 2012). SUCNR1 has typical disulfide bonds and two N-glycosylation sites in the extracellular region (Gilissen et al., 2016). In addition to succinate, the ligands of SUCNR1 include maleate and methylmalonate, but their activation potency is much lower than that of succinate

(Ariza, et al., 2012). Antagonists of 2c and 4c can bind to remote sites of SUCNR1 and act as allosteric antagonists (Bhuniya et al., 2011). The selectivity of SUCNR1 for succinate is mediated by four positively charged key residues in the binding cavity enclosed by the seven transmembrane domains of the receptor, namely Arg99, His103, Arg252, and Arg281 (He et al., 2004).

## 2.3 Mechanisms of action and related signaling pathways of succinate

### 2.3.1 Transmembrane transport of succinate

Succinate is typically confined to mitochondria and is generated by the TCA cycle. However, under stress, accumulated succinate in mitochondria can be secreted into the cytoplasm or extracellular space to exert its effects. Succinate has different effects in different cellular locations, making its transport between different locations crucial. Transport of succinate between mitochondria, cytoplasm, and extracellular space relies on various transport channels. Under physiological pH conditions, succinate is a charged molecule and cannot diffuse across membranes. With the accumulation of succinate in mitochondria, it can be exported to the cytoplasm via the dicarboxylic acid translocator SLC25A10 on the inner mitochondrial membrane and the voltage-dependent anion channel (VDAC/porin) on the outer mitochondrial membrane (Selak et al., 2005). During ischemia-reperfusion and intense exercise, succinate efflux is mainly mediated by monocarboxylate transporter 1 (MCT1) (Reddy et al., 2020; Prag et al., 2021). Organic anion transporters (OATs) can also transport succinate from the intracellular space to the extracellular space, but they have low affinity for succinate (Kaufhold et al., 2011). The plasma membrane-bound Na-dependent dicarboxylic acid transporter NaDC3 (SLC13A3 gene) is an important pathway for uptake of extracellular succinate (Zhunussova et al., 2015). As a member of the SLC13 family, NaDC3 has a broad transport spectrum for dicarboxylic and tricarboxylic acids, with succinate being one of its highest-affinity substrates. The transport process of NaDC3 is dependent on sodium ions (Na<sup>+</sup>). When one molecule of succinate binds to three Na<sup>+</sup> at the substrate binding site of NaDC3, the “rocking bundle” mechanism induces a structural change in NaDC3, thereby facilitating the uptake of succinate into the cell (Pajor, 2014). Lithium ions can interfere with this process (Schlessinger et al., 2014).

### 2.3.2 Succinate-related signaling pathways

In addition to serving as an intermediate in metabolism, succinate can also be secreted into the cytoplasm and extracellular space as a signaling molecule to regulate cellular responses. The primary mechanisms involved are the succinate-HIF-1 $\alpha$  axis, succinylation, and the succinate-SUCNR1 axis (**Fig. 2**).

#### 2.3.2.1 Intracellular signaling pathways of succinate

As an oxygen sensor of cells, HIF-1 $\alpha$  is normally activated by hypoxia, thereby initiating the transcription of a series of hypoxia-inducible genes to adapt to the hypoxic environment. It plays an important role in angiogenesis, glycolysis, cancer-cell invasion and metastasis, as well as the expression of inflammatory genes. It is known that hypoxia induces succinate accumulation and that accumulated succinate in mitochondria is transported to the cytoplasm to stabilize HIF-1 $\alpha$  in response to hypoxia (Agani et al., 2000). Exogenous addition of succinate can bypass hypoxia and activate HIF-1 $\alpha$ -related pathways under normoxic conditions, which is known as “pseudohypoxia.” Thus, succinate has been discovered by some scientists to act as a superior pseudo-hypoxia agent, exerting excellent effects in angiogenesis (Löffler et al., 2023). As a substrate of PHD,  $\alpha$ -KG produces succinate as a byproduct during the hydroxylation of HIF-1 $\alpha$ . Therefore, high concentrations of succinate directly inhibit PHD activity, hindering degradation of HIF-1 $\alpha$  and stabilizing it. ROS can impair PHD activity by oxidizing the key PHD cofactor Fe<sup>2+</sup> to Fe<sup>3+</sup>. Thus, succinate can also indirectly stabilize HIF-1 $\alpha$  by inducing ROS production (Mao et al., 2020). Tannahill's study showed that intracellular succinate accumulation enhances the expression of interleukin 1- $\beta$  (IL-1 $\beta$ ) by stabilizing HIF-1 $\alpha$  in macrophages (Tannahill, et al., 2013). Succinate can also promote vascular endothelial growth-factor (VEGF) expression

through HIF-1 $\alpha$ , thereby regulating angiogenesis.

As a post-translational modification, succinylation of lysine is closely related to metabolic regulation. Succinylation of lysine can activate SDH, thereby contributing to the self-regulation of succinate levels in mitochondria (Park et al., 2013). Lysine succinylation participates in the regulation of various metabolic signaling pathways by the desuccinylase SIRT5 and is widespread in mitochondrial energy metabolism. Studies have found that LPS stimulation increases protein succinylation by promoting succinate accumulation and inhibiting the expression and activity of SIRT5 (Tannahill, et al., 2013).

### 2.3.2.2 Succinate-SUCNR1 axis

Extracellular succinate acts on SUCNR1 on the cell membrane through autocrine or paracrine mechanisms, mediating ligand-receptor signaling. For instance, macrophages can migrate and generate cytokines through the autocrine action of succinate. Succinate can also exert its biological effects by binding to SUCNR1 on other cells through paracrine mechanisms. In human skeletal muscle, SUCNR1 mRNA is not expressed in muscle fibers but is highly expressed in subsets of macrophages. Therefore, in the adaptive response of skeletal muscle to exercise, succinate and SUCNR1 may induce a transition from fast to slow muscle fibers through a paracrine mechanism involving M2-like macrophages (Abdelmoez et al., 2023). Additionally, in mice with high blood-sugar levels, succinate that accumulates abnormally in bone-marrow stromal cells (BMSCs) exerts its effects by binding to SUCNR1 in osteoclast precursor cells (Guo, et al., 2017). The cascade reactions mediated by SUCNR1 are similar to the Gi and Gq signaling cascades, and lead to accumulation of inositol trisphosphate (IP3), calcium mobilization, downregulation of 3',5'-cyclic adenosine monophosphate (cAMP), and phosphorylation of extracellular signal-regulated kinase (ERK) (He, et al., 2004). Activation of SUCNR1 by succinate can also promote calcium-dependent nitric oxide (NO) and prostaglandin E2 (PGE2) production. The succinate-SUCNR1 axis plays a significant role in immune-inflammatory responses (Zhang and Lang, 2023) and contributes to the development of various inflammatory diseases by enhancing inflammation. These illnesses include diabetic retinopathy, hypertension, ulcerative colitis, and RA (He, et al., 2004; Sadagopan et al., 2007; Toma et al., 2008). Activation of the succinate-SUCNR1 pathway induces M1 macrophages to release pro-inflammatory cytokines and chemokines, thereby aggravating inflammation (Van Diepen et al., 2017; Murphy and O'Neill, 2018). Succinate can also promote dendritic-cell migration and activity of antigen-presenting cells by triggering an immune response with SUCNR1. Interestingly, in a mouse model of multiple sclerosis (MS), succinate secreted by MS tissues interacted with SUCNR1 expressed on transplanted neural stem cells (NSCs), inducing secretion of PEG2 by NSCs and clearance of extracellular succinate, and thereby exerting anti-inflammatory effects (Peruzzotti-Jametti et al., 2018). Simultaneously, the succinate-SUCNR1 axis can control inflammation in the adipose tissue of healthy lean individuals by activating the anti-inflammatory phenotype (M2 type) of macrophages (Keiran et al., 2019). It is worth noting that the succinate-HIF-1 $\alpha$  axis and the succinate-SUCNR1 axis are not entirely independent. Extracellular succinate can increase HIF-1 $\alpha$  expression through the SUCNR-1-PI3K/Akt signaling pathway (Mao, et al., 2020).

### 2.3.3 Role of succinate in bone and cartilage metabolism

The succinate-SUCNR1 axis has been found to stimulate osteoclastogenesis via the NF- $\kappa$ B pathway, thus mediating the bone-resorption process. However, in osteogenesis, succinate appears to exhibit more complex effects. It can inhibit the activity and osteogenic differentiation of MSCs and other osteoprogenitor cells via the HIF-1 $\alpha$  pathway. In vitro experiments, though, have also shown that specific concentrations of succinate can promote the migration and osteogenic differentiation of human mesenchymal stem cells (hMSCs). Furthermore, exogenous addition of succinate has been found to play a positive role in angiogenesis by promoting the proliferation and migration of endothelial cells without harming osteoblasts. This difference may be related to the concentration and site of action of succinate. Additionally, research indicates that succinate-induced metabolic remodeling accelerates hypertrophic differentiation of chondrocytes, thereby disrupting the homeostasis of cartilage. Succinate is also associated with the inflammatory phenotype of synovial fibroblasts.

**Fig. 2 Mechanisms of action and related signaling pathways of succinate.**

Under stress conditions, such as LPS, hypoxia, ischemia, and hyperglycemia, succinate accumulates in mitochondria. With the accumulation of succinate in mitochondria, it can then be exported to the cytoplasm via the dicarboxylic acid translocator SLC25A10 on the inner mitochondrial membrane and the voltage-dependent anion channel (VDAC/porin) on the outer mitochondrial membrane. Succinate in the cytoplasm stabilizes HIF-1 $\alpha$  by inhibiting PHD activity, thereby promoting the transcription of HIF-1 $\alpha$ -dependent hypoxia-inducible genes, such as VEGF. The succinate in cytoplasm can be exported outside from the cell through the transporter MCT1. Extracellular succinate enters the cell through the transporter NaDC3 with the help of Na<sup>+</sup>. Extracellular succinate binds to SUCNR1 through autocrine or paracrine mechanisms and activates downstream signaling pathways. The cascade reactions mediated by SUCNR1 is similar to the Gi and Gq signaling cascades, leading to the accumulation of IP3, calcium mobilization, downregulation of cAMP, and phosphorylation of ERK, which activates the NF- $\kappa$ B pathway and leads to IL-1 $\beta$  production. Activation of SUCNR1 by succinate can also promote calcium-dependent NO and PGE2 production. Lysine succinylation participates in the regulation of various metabolic signaling pathways by the desuccinylase SIRT5. (Image created with BioRender.com, with permission).

### 3 Role of succinate in bone-related diseases

Bone homeostasis is a dynamic balance between bone formation and bone resorption, and is crucial for normal bone development and bone-defect repair. Osteoclasts (OCs), derived from hematopoietic progenitor cells, participate in bone resorption, playing a vital role in bone homeostasis. When OCs are overactive, the rate of bone resorption exceeds that of bone formation, disrupting bone homeostasis and often leading to metabolic bone diseases. A large number of metabolomic studies have demonstrated an association between elevated levels of succinate and bone-related diseases such as OP, OA, and RA, suggesting that succinate and related factors may serve as therapeutic targets for these conditions. Clarifying the specific role of succinate in bone-related diseases could provide new insights for treatment.

#### 3.1 Role of succinate in OP

OP is a systemic metabolic bone disease characterized by reduced bone density and bone-structure


deterioration, which is a typical manifestation of bone-homeostasis imbalance. The association between elevated succinate levels and OP is quite evident. Numerous metabolomic studies have observed significantly higher succinate levels in the metabolites of OP patients (Zhao et al., 2018; Yu et al., 2019; Deng et al., 2021). Additionally, a decrease in SDH activity has been observed in the articular cartilage of patients with femoral neck fracture caused by primary OP (Nahir et al., 1990). Studies have found a negative correlation between succinate and bone density (Wang et al., 2023). However, contradictory results exist, with a 4-week dietary supplement treatment based on succinate leading to increased bone mass and calcium content in aging mice (Maevsky et al., 2008). Nevertheless, this study had limitations such as insufficiently controlled variables and a small sample size. Evidence also suggests that succinylation contributes to the development of OP (Su et al., 2017).

In OP patients, OC activity is abnormally high. Studies by Guo et al. found that the succinate-SUCNR1 axis is involved in the activation of OCs and in enhanced bone resorption mediated by OCs in type 2 diabetes (T2D) mice (Guo, et al., 2017). In T2D mice with concomitant OP, succinate was increased 24-fold in BMSCs. Further research revealed that succinate can stimulate OC generation, differentiation, and maturation, promoting bone resorption. Besides directly stimulating OC generation, succinate can induce OC generation by downregulating levels of IL-4 and IL-13 and activating the expression of TNF and IL-1 $\beta$ . The specific mechanism by which accumulated succinate in BMSCs induces OC generation involves the succinate-SUCNR1 axis. As SUCNR1 is not expressed on BMSCs but is highly expressed in hematopoietic lineage cells (including OCs and OC precursor cells), accumulated succinate in BMSCs acts on SUCNR1 on the membrane of OC precursor cells through paracrine action, thereby promoting their transformation into OCs via stimulation of the NF- $\kappa$ B signaling pathway (p50/p65).

Mitochondrial dysfunction is associated with aging-related phenotypes such as OP (Guo, et al., 2020b). Succinate mediates mitochondrial dysfunction, and mitochondrial dysfunction leads to the accumulation of succinate (Nair et al., 2019). The succinate-SUCNR1 axis can promote mitochondrial fission through the downstream ERK1/2-DRP1 pathway (Nguyen et al., 2022). In mitochondria with impaired function, metabolic remodeling occurs, with decreased ATP synthesis and increased production of ROS. The transition from oxidative metabolism to glycolytic metabolism in aging bones is a major determinant of bone loss in mice (Liu et al., 2024). During mitochondrial electron transport, complexes I and III generate ROS, whose toxic byproducts damage the mitochondria. This damage then triggers adjacent mitochondria to release more ROS, creating a vicious cycle. Accumulated ROS can act as second messengers in OC generation (Callaway and Jiang, 2015), inducing bone resorption and thereby playing a role in the onset and progression of OP.

Reduced osteogenic differentiation of osteoprogenitor cells is also one of the contributing factors to OP. Mitochondrial transfer is a novel mode of intercellular communication. Macrophages play an important role in bone homeostasis (Gao et al., 2019). Under normal circumstances, macrophages can promote the osteogenic differentiation of MSCs by transferring mitochondria to MSCs. MSCs that receive functional mitochondria exhibit enhanced osteogenic potential, while those that receive damaged mitochondria experience impaired osteogenic potential. However, in OP, the phenotype and metabolic state of macrophages change, transitioning them from M2-type macrophages to M1-type macrophages. Inflammatory activated macrophages transfer oxidatively damaged mitochondria to MSCs via microvesicles, causing cellular damage and metabolic reshaping of MSCs. This metabolic disorder leads to the accumulation of succinate in MSCs by inhibiting the activity of SDH, thereby promoting the release of proinflammatory factors (such as IL-1 $\beta$ ) and ROS through the Hif-1 $\alpha$  signaling pathway and affecting the osteogenic differentiation of MSCs (Cai et al., 2023). Furthermore, mitochondria-derived oxidative damage from M1-type macrophages can increase apoptosis in MSCs and impair their proliferation. In addition to its effect on MSCs, succinate can inhibit the proliferation and osteogenic differentiation of mouse osteoprogenitor cells MC3T3-E1, which may be related to the downregulation of RUNX2 and OCN (Tanjun et al., 2019). The role of succinate in OP is illustrated in **Fig. 3**.





**Fig. 3 Role of succinate in OP.**

(A) When SDH activity is inhibited, accumulated succinate in BMSCs acts on SUCNR1 on the membrane of OC precursor cells through paracrine action, thereby promoting their transformation into OCs via stimulation of the NF- $\kappa$ B signaling pathway. (B) During OP, inflammatory activated macrophages transfer oxidatively damaged mitochondria to MSCs via macrovesicles, causing cellular damage and metabolic reshaping of MSCs. This leads to accumulation of succinate in MSCs by inhibiting the activity of SDH, thereby promoting release of IL-1 $\beta$  and ROS through the Hif-1 $\alpha$  signaling pathway. This promotes apoptosis of MSCs and inhibits proliferation and osteogenic differentiation of MSCs. (C) The succinate-SUCNR1 axis promotes mitochondrial fission via the ERK1/2-DRP1 pathway. In addition, mitochondrial dysfunction can lead to accumulation of succinate, inflammation, and ROS generation, creating a vicious cycle. These ROS act as second messengers, inducing osteoclast generation and bone resorption. (Image created with BioRender.com, with permission).

### 3.2 Succinate and RA

RA is a chronic progressive autoimmune disease characterized by inflammation and angiogenesis. Metabolomic analysis has identified succinate as a potential biomarker for RA (Li et al., 2016). RA patients have abundant succinate in their synovial fluid (Littlewood-Evans et al., 2016). Studies have also shown a correlation between levels of succinate in synovial fibroblasts (SF) and the degree of paw swelling in arthritic mice, suggesting that succinate levels could serve as a biological indicator of RA severity (Littlewood-Evans, et al., 2016). Succinate was found to be the most differentially expressed metabolite between RA and other joint diseases (Kim et al., 2014).

Angiogenesis in the synovium in RA promotes inflammation and pannus formation, leading to destruction of cartilage and bone (Szekanecz et al., 2010). Succinate can promote synovial angiogenesis, thereby driving the progression of RA. This process involves the HIF-1 $\alpha$ -VEGF axis, the succinate-SUCNR1 axis, transforming growth factor-beta1 (TGF- $\beta$ 1), and a reversal of SDH activity (Li, et al., 2018). In the synovium of RA, due to local hypoxia, TGF- $\beta$ 1 increases succinate production by reversing SDH activity in synovial fibroblasts (Li, et al., 2016). Accumulated succinate then induces VEGF expression by stabilizing HIF-1 $\alpha$  in the cytoplasm. VEGF can mediate angiogenesis and also promote synovial endothelial-cell activation to induce inflammation, thereby establishing crosstalk between angiogenesis and joint inflammation in RA. Additionally, accumulated succinate can be transported to the extracellular space, activating SUCNR1 on endothelial cells to induce VEGF production, further exacerbating angiogenesis. These two pathways work together to exacerbate inflammation and angiogenesis in RA synovium. Moreover, succinate can induce dysfunctional angiogenesis by inducing basic fibroblast growth factor (Biniacka et al., 2016). Inhibitors of SDH such as dimethyl malonate can

attenuate angiogenesis and inflammation in RA synovium by reducing succinate production.

The proinflammatory effect of succinate is also significant in the progression of RA. Synovial hypoxia in RA leads to succinate accumulation in SF. IL-1 $\beta$  is released from macrophages triggered by succinate in a SUCNR1-dependent manner, and thus serves as a core driver of arthritis inflammation. Maturation of IL-1 $\beta$  is mediated by the NLRP3 inflammasome (Wen et al., 2011). Accumulated succinate in SF can activate the NLRP3 inflammasome with the involvement of HIF-1 $\alpha$ , inducing IL-1 $\beta$  secretion and promoting inflammation. IL-1 $\beta$ -mediated inflammation, in turn, induces fibroblast activation by inducing TGF- $\beta$ 1. Activated fibroblasts further stimulate the generation of IL-1 $\beta$ , thus perpetuating a cycle that leads to sustained polarization of fibroblasts, ultimately resulting in fibrosis of the synovial tissue. Therefore, accumulation of succinate serves as a metabolic signal that connects IL-1 $\beta$  and TGF- $\beta$ 1 and establishes crosstalk between inflammation and fibrosis, further exacerbating tissue damage in RA (Li, et al., 2016). Additionally, succinate can exacerbate synovial tissue hyperplasia and inflammation by promoting the invasion and migration capacity of fibroblast-like synoviocytes (FLS) (Hu et al., 2024). The role of succinate in RA is illustrated in **Fig. 4**.

### 3.3 Succinate and OA

As an inflammatory joint disease, OA progresses significantly through inflammation-driven degeneration of joint cartilage (Jones et al., 2009). Some evidence suggests a link between succinate and OA. Reduced activity of SDH and alterations in the TCA cycle have been observed in OA (Nahir, et al., 1990; Van Pevenage et al., 2023), indicating changes in succinate-related metabolic pathways. However, when it comes to the changes of succinate levels in OA, v metabolomic studies have failed to reach a consensus. While Huang et al. reported decreased succinate levels in OA patients compared to healthy individuals (Huang et al., 2020), most studies affirm the crucial role of excess succinate in OA. Qualitative analyses of metabolic profiles in OA patients have revealed elevated levels of succinate in urine and blood samples (Liao et al., 2023).

Synovitis is an early manifestation of OA, and synovial fibroblasts are the primary effectors in synovitis (Nanus et al., 2020). Many lines of evidence suggest that the inflammation-activated phenotype of fibroblasts is closely related to their metabolic status (Yin et al., 2018; Farah et al., 2021). For instance, obese OA patients have significantly higher levels of succinate in their synovial fluid compared to those with normal BMI (Farah et al., 2022). While glycolysis is increased in obesity, this metabolic remodeling induces a phenotype of inflammation-activated synovial fibroblasts, which in turn secrete more IL-6, thereby promoting synovial inflammation (Xie et al., 2015). Additionally, succinate treatment in synovial fibroblasts can induce activation of the NLRP3 inflammasome and increase intracellular inflammatory responses (Li, et al., 2016).

Chondrocyte homeostasis is related to the occurrence and development of OA, and involves changes in metabolism. While healthy articular chondrocytes primarily utilize glycolysis for energy, hypertrophic chondrocytes rely more on oxidative phosphorylation and mitochondrial respiration (Pollesello et al., 1991). Increased mitochondrial respiration can cause catabolism and progression of OA in mouse articular cartilage (Shen et al., 2017). Abat is an enzyme that metabolizes GABA to succinate. Overexpression of Abat was found to cause chondrocyte hypertrophy in vitro and accelerate OA progression in mice. Treatment with vigabatrin, an FDA-approved small-molecule Abat inhibitor, can prevent OA progression caused by injury (Shen et al., 2019). Elevated levels of Abat lead to increased succinate levels, thereby enhancing the TCA cycle and mitochondrial respiration, which disrupts cartilage homeostasis and promotes OA progression. The role of succinate in OA is illustrated in **Fig. 4**.

#### Fig. 4 Role of succinate in OA and RA.

Abat is an enzyme that metabolizes GABA to succinate. Increased succinate levels enhance the TCA cycle and mitochondrial respiration, which disrupts cartilage homeostasis, thereby promoting cartilage catabolism and chondrocyte hypertrophy. Synovial hypoxia in RA leads to reversed SDH activity, resulting in succinate accumulation in SF. Succinate promotes angiogenesis via the HIF-1 $\alpha$ -VEGF and succinate- SUCNR1 pathways. Accumulated succinate triggers macrophages to release IL-1 $\beta$  via the SUCNR1 pathway. IL-1 $\beta$  release is mediated by the NLRP3 inflammasome and activated by succinate with HIF-1 $\alpha$  involvement. This process promotes inflammation and fibroblast activation through TGF- $\beta$ 1, creating a cycle of inflammation and fibrosis. Succinate also enhances synovial-tissue hyperplasia and inflammation by increasing the invasiveness and migration of fibroblast-like synoviocytes (FLS). Thus, succinate links IL-1 $\beta$  and TGF- $\beta$ 1, exacerbating RA progression. (Image created with BioRender.com, with permission).

### 3.4 Succinate and fractures

Bone regeneration after fracture is a complex process, and successful bone regeneration depends on the balanced interaction of anabolic and catabolic processes, including inflammation and angiogenesis, matrix formation, and tissue remodeling, which has high dynamic energy requirements (Schmidt-Bleek et al., 2014). However, the influence of metabolism on the progression and outcomes of bone healing remains insufficiently studied. Several studies suggest that succinate may contribute to fracture healing. Scientists have found that during the early stages of bone-healing inflammation, the levels of succinate are higher in rats with successful bone regeneration compared to those with impaired bone regeneration (Löffler, et al., 2023). Additionally, succinate has been discovered to accelerate the healing of mandibular fractures in guinea pigs (Faustov et al., 2001).

Angiogenesis is crucial in early osteoinduction. Succinate is a safer pseudohypoxia agent than cobalt chloride, and pseudohypoxia induced by succinate contributes to bone regeneration by promoting angiogenesis. An *in vitro* study found that the cultivation of saos-2 (osteoblasts) on succinate-alginate xerogel films promoted the proliferation of osteoblasts, with the effective succinate concentration ranging from 10  $\mu$ mol/L to 100 mmol/L (Deering et al., 2022). 1  $\mu$ mol/L and 10  $\mu$ mol/L of succinate can promote angiogenesis and maintain long-term vascular stability (Deering, et al., 2022). This indicates that succinate at a lower concentration (10  $\mu$ mol/L) can promote angiogenesis and early osteoinduction without damaging osteoblasts.

Succinate plays a significant role in successful bone healing, stimulating hMSCs migration in the fracture gap and supporting osteogenesis and bone formation. It can activate macrophages via SUCNR1, inducing expression of IL-1 $\beta$ . It has been found that sustained exposure to IL-1 $\beta$  in osteogenic cultures of hMSCs leads to attenuated osteogenic differentiation, whereas stimulation with IL-1 $\beta$  at the beginning of differentiation can promote osteogenic differentiation of hMSCs (Löffler, et al., 2023). The specific effects of succinate are concentration-dependent. In Löffler's study, 50  $\mu\text{mol/L}$  succinate acted as a significant vascular endothelial-growth-factor inductor, mediated by the succinate-SUCNR1 axis (Löffler, et al., 2023). The process involved upregulation of STAT3 and ERK 1/2, and expression of VEGF, independently of HIF-1a. Moreover, higher concentrations of succinate (200-800  $\mu\text{mol/L}$ ) were found to promote tubular structure formation in vitro and in zebrafish (Mu et al., 2017). Treatment with 400  $\mu\text{mol/L}$  or 800  $\mu\text{mol/L}$  of succinate significantly increased the vitality of human umbilical-vein endothelial cells. Addition of succinate (50 or 500  $\mu\text{mol/L}$ ) in vitro promoted the migration of primary hMSCs, with 500  $\mu\text{mol/L}$  of succinate particularly inducing osteogenic differentiation and intense mineralization of hMSCs. Notably, in the absence of other osteogenic stimuli, paracrine signals from succinate-stimulated macrophages, especially M2-type macrophages stimulated with 50  $\mu\text{mol/L}$  succinate, directly induced osteogenesis in hMSCs. There is a direct connection between succinate, SUCNR1, and tissue healing, as treatment with 50  $\mu\text{mol/L}$  of succinate accelerated the healing of skin incisions in mice (Ko et al., 2017). In addition, succinate can play a positive role in fracture healing by activating macrophage responses and neovascularization, as well as reducing fibrosclerosis in damaged muscle areas (Faustov, et al., 2001).

### 3.5 Succinate and periodontitis

Succinate plays a significant role in the progression of periodontitis. Elevated levels of succinate have been found in the gingival crevicular fluid of periodontal patients (Ohwaki, 1988; Lu et al., 2015). Targeted metabolomic analysis of 28 gingival plaque samples found that individuals with severe periodontitis had elevated succinate levels and dysregulated oral microbiota in their dental plaque (Guo et al., 2022). Additionally, succinate levels in periodontal pockets are significantly correlated with clinical indicators of disease severity (Lu et al., 2014). These pieces of evidence suggest potential involvement of succinate in the progression of periodontitis.

The development of periodontitis begins with colonization by pathogenic bacteria, and microbial metabolism is a major source of abnormal succinate accumulation (Gao et al., 2014). Many periodontitis-related bacteria produce succinate through metabolism, for example *P. gingivalis*, *P. intermedia*, and *Treponema denticola* (Shah and Williams, 1987; Takahashi, et al., 2000; Takahashi and Yamada, 2000; Kin, et al., 2020). Succinate can mediate the pro-inflammatory response of periodontal ligament stem cells (PDLSCs) triggered by *P. gingivalis* (Su et al., 2020). PDLSCs undergo metabolic reprogramming in response to *P. gingivalis*, from oxidative phosphorylation to glycolysis, and succinate accumulation, which triggers the inflammatory response of PDLSCs through the succinate-SDH-Hif-1 $\alpha$  axis and ROS production. However, in contrast to LPS-stimulated macrophages (Reduced SDH activity), PDLSCs infected with *P. gingivalis* show increased SDH production, SDH activity, and succinate levels. Therefore, the source of succinate accumulation in PDLSCs infected with *P. gingivalis* still requires further investigation. Succinate also enhances the expression of inflammatory cytokines, particularly TNF- $\alpha$  and IL-1 $\beta$ , by activating SUCNR1, thereby inducing alveolar bone loss (Guo, et al., 2022). Given that dysbiosis of the oral microbiome is a key phenotype and cause of periodontitis, the impact of succinate on the periodontal microbiota also contributes to the development of periodontitis. In vitro experiments have shown that succinate can directly stimulate the growth and virulence gene expression of *Fusobacterium nucleatum*, leading to an imbalance in the oral microbiota (Guo, et al., 2022; Li, 2023). Guo et al. found through 16S sequencing that systemic application of succinate significantly increased the abundance of periodontal-associated bacteria such as *Bacteroidetes*, whereas these changes were not evident in SUCNR1 knockout mice. The use of a SUCNR1 antagonist reduced the abundance of periodontal-related bacteria, indicating that succinate affects the microbiota composition through the SUCNR1

receptor (Guo, et al., 2022). Additionally, a metabolic cooperation has been found between *P. gingivalis* and *T. denticola*, and is known as succinate cross-feeding. *T. denticola* produces succinate, which is utilized by *P. gingivalis* and converted into toxic metabolites. This cooperation may exacerbate the progression of periodontal disease and help bacteria more efficiently acquire energy and carbon sources in nutrient-limited environments (Kin, et al., 2020).

Nevertheless, in the absence of bacteria-mediated effects, exogenous addition of succinate can act as a pseudo-hypoxic agent to promote proliferation, migration, and osteogenesis of human periodontal ligament stem cells (hPDLSCs) by activating the HIF-1 $\alpha$  pathway (Mao, et al., 2020). The role of hypoxia in osteogenic differentiation of MSCs remains controversial. The reason why succinate promotes osteogenesis of hPDLSCs may be that in addition to serving as a hypoxia mimetic, it can also enhance oxidative capacity, thereby enhancing bone-matrix protein biosynthesis in later stages. It is worth noting that hypoxia-mediated proliferation of hPDLSCs depends on the SUCNR1 pathway, while hPDLSC proliferation mediated by exogenous succinate includes SUCNR1-dependent and SUCNR1-independent pathways. High levels of exogenous succinate can bypass SUCNR1 and directly induce cellular effects through the cell membrane. The metabolic changes induced by succinate in cell metabolism depend on its concentration. Exogenous succinate at concentrations of 1 and 5 mmol/L (mainly 5 mmol/L) can promote proliferation, migration, and osteogenic ability of hPDLSCs, while high concentrations of succinate (25 mmol/L) can decrease cell viability (Ko, et al., 2017). This inhibitory effect of high-concentration succinate may be associated with succinate-induced apoptotic pathways (caspase3/7) (Wentzel et al., 2017), induction of DNA hypermethylation, epigenetic silencing (Jiang and Yan, 2017), protein succinylation, and induction of oxidative damage by ROS (García-Prat et al., 2017). The role of succinate in periodontitis is illustrated in **Fig. 5**.

#### **Fig. 5 Role of succinate in periodontitis.**

Succinate can mediate the pro-inflammatory response of *P. gingivalis* to PDLSCs. PDLSCs respond by shifting from oxidative phosphorylation to glycolysis, accumulating succinate, and triggering inflammation via the succinate-SDH-HIF-1 $\alpha$  axis and ROS production. *P. gingivalis*-infected PDLSCs show increased SDH production and activity. Succinate also supports the growth of periodontal pathogens through SUCNR1, disrupting the balance of oral microbiota. Additionally, SA activates SUCNR1, enhancing TNF- $\alpha$  and IL-1 $\beta$  expression and inducing alveolar bone loss. (Image created with BioRender.com, with permission).

### 3.6 Other findings

Jin et al. found that succinate can also stably bind to collagen fibers through hydrogen bonds, enhancing the ability of the collagen matrix to attract more calcium ions and thus accelerating the mineralization of collagen fiber. This process is crucial for both tooth and bone formation. Succinate promotes dentin restoration and remineralization of demineralized dentin, and enhances its mechanical properties (Jin et al., 2022), which suggests potential for application of succinate in dental caries restoration.

## 4 Treatment targets

Abnormal accumulation of succinate is associated with changes in SDH activity, and the succinate-SUCNR1 axis and succinate-HIF-1 $\alpha$  axis mediate the pathogenesis of bone-related diseases. Succinate also mediates the mitochondrial transfer from macrophages to MSCs, promoting OP development. Moreover, the multifaceted actions of succinate as a signaling molecule vary depending on its cellular location and concentration. Therefore, targeting key points in succinate-induced pathogenesis may offer new insights into the treatment of bone-related diseases.

### 4.1 SDH inhibitors

Activation of the reverse activity of SDH is known to lead to succinate accumulation (Chouchani et al., 2014b), which plays a role in synovial inflammation in the joints (Li, et al., 2018). SDH inhibitors (such as dimethyl malonate) can prevent abnormal succinate accumulation and inhibit synovial angiogenesis by blocking the HIF-1 $\alpha$ /VEGF axis (Li, et al., 2018), thus they serve as a therapeutic target for RA.

### 4.2 SUCNR1 antagonists

The succinate-SUCNR1 axis mediates various pathological processes, and the use of SUCNR1 antagonists can effectively block this pathway, hindering succinate-driven effects on OP, OA, RA, and periodontitis. In the K/BxN serum transfer-induced arthritis model and the antigen-induced mouse arthritis model, the absence of SUCNR1 has been shown to alleviate arthritic inflammation and lesions (Rubić-Schneider et al., 2017). Currently, the compound “4c” appears to be the most effective SUCNR1 antagonist in vitro (Ariza, et al., 2012), and has the potential to become a drug for treating inflammation-related diseases and cancer metastasis. Local application of the SUCNR1 antagonist gel formulation of 7a effectively inhibits inflammation, microbial dysbiosis, and bone loss in mice with periodontitis (Li, et al., 2018). This effect is mediated by inhibiting host SUCNR1 signaling, rather than through direct interaction with the host microbiota (Thomas et al., 2024).

### 4.3 Mitochondrial transfer

Mitochondrial transfer plays a crucial role in maintaining bone homeostasis and preserving bone-cell vitality. Artificial mitochondrial transfer to damaged cells or tissues can effectively repair the damage (Cai, et al., 2023). Guo et al. reported that mitochondrial transplantation therapy enhanced the proliferation, migration, and osteogenic differentiation of bone-marrow MSCs, improving bone-defect healing (Guo et al., 2020a). In OP, the mitochondrial transfer from pro-inflammatory M1 macrophages to MSCs inhibits MSC osteogenic differentiation. Utilizing artificial mitochondrial transfer by transferring healthy mitochondria into MSCs could serve as an effective therapeutic target for enhancing osteogenic differentiation in OP treatment.

### 4.4 Abat inhibitors

Abat is a key enzyme responsible for the metabolism of GABA to succinate (Osei and Churchich, 1995). Elevated Abat levels can lead to succinate accumulation, thereby enhancing the TCA cycle and increasing mitochondrial respiration. This affects chondrocyte homeostasis and causes OA. Treatment with Abat inhibitors (such as gabaculine) can prevent OA progression caused by cartilage injury in mice (Shen, et al., 2019).

### 4.5 Exogenous addition of succinate

The effects of succinate vary depending on the concentration. Exogenously adding specific concentrations of succinate has potential as a microbial therapy. Lower concentrations have a positive effect on fracture healing, with 10  $\mu\text{mol/L}$  of succinate promoting early bone induction and 50-500  $\mu\text{mol/L}$  of succinate aiding in the migration and osteogenic differentiation of hMSCs within the fracture gap (Löffler, et al., 2023). However, lower concentrations (primarily 5  $\text{mmol/L}$ ) of exogenous succinate, like hypoxia, can act as a pseudohypoxic agent by activating the HIF-1 $\alpha$  pathway to promote proliferation, migration, and osteogenic capacity of hPDLSCs, while higher concentrations of succinate (25  $\text{mmol/L}$ ) may decrease cell viability (Mao, et al., 2020).

## 5 Discussion

Succinate plays a significant role in bone-related diseases. Abnormal accumulation of it is closely associated with pathological processes such as OP, OA, RA, and periodontal diseases through the succinate-SUCNR1 axis and succinate-HIF-1 $\alpha$  axis, and links mitochondrial dysfunction to the pathogenesis of OP. Current research suggests that inhibiting succinate generation or modulating its metabolism, for example with Abat inhibitors, SDH inhibitors, or SUCNR1 antagonists, may provide new avenues for treating these diseases. However, translating the results of these novel compounds from in-vitro and animal models to human trials is challenging. There are issues such as low drug exposure at target sites and clinical safety concerns. Systemic administration of SUCNR-1 antagonists may attenuate the anti-inflammatory properties of macrophages, potentially leading to other adverse reactions (Keiran, et al., 2019). Additionally, exploring the regulatory role of succinate in stem-cell function and developing more precise drug-targeting strategies may bring further breakthroughs in the treatment of bone diseases in the future.

### Data availability statement

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

### Acknowledgments

This work is supported by the grants from the National Natural Science Foundation of China (82170984) and Zhejiang Provincial Natural Science Foundation of China under Grant No. Q24H140009. We want to express our gratitude for the drawing materials provided by BioRender (Agreement number: WY27MZUMKT, JO27MZUS7I, ES27MZULHJ, MH27MZUSYX, UR27MZUH3E).

### Author contributions

Zuping WU and Qiaoli DAI were involved in conceptualization, visualization, writing – original draft, and writing review & editing. Ying WANG was involved in writing – review & editing. Na WU was involved in writing – review & editing. Chenyu WANG was involved in writing – review & editing. Jiejun SHI was involved in writing – review & editing, supervision, and conceptualization. All authors read and approved the final manuscript and, therefore, had full access to all the data in the study and take responsibility for the integrity and security of the data.

### Compliance with ethics guidelines

Zuping WU, Qiaoli DAI, Ying WANG, Na WU, Chenyu WANG, and Jiejun SHI declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by any of the authors

### References

- Abdelmoez AM, Dmytriyeva O, Zurke YX, et al., 2023. Cell selectivity in succinate receptor *sucnr1/gpr91* signaling in skeletal muscle. *Am J Physiol Endocrinol Metab*, 324(4):E289-e298.  
<https://doi.org/10.1152/ajpendo.00009.2023>
- Agani FH, Pichiule P, Chavez JC, et al., 2000. The role of mitochondria in the regulation of hypoxia-inducible factor 1 expression during hypoxia. *J Biol Chem*, 275(46):35863-35867.  
<https://doi.org/10.1074/jbc.M005643200>
- Ariza AC, Deen PM, Robben JH, 2012. The succinate receptor as a novel therapeutic target for oxidative and metabolic stress-related conditions. *Front Endocrinol (Lausanne)*, 3:22.  
<https://doi.org/10.3389/fendo.2012.00022>

- Bardella C, Pollard PJ, Tomlinson I, 2011. Sdh mutations in cancer. *Biochim Biophys Acta*, 1807(11):1432-1443.  
<https://doi.org/10.1016/j.bbabi.2011.07.003>
- Bhuniya D, Umrani D, Dave B, et al., 2011. Discovery of a potent and selective small molecule hgr91 antagonist. *Bioorg Med Chem Lett*, 21(12):3596-3602.  
<https://doi.org/10.1016/j.bmcl.2011.04.091>
- Biniecka M, Canavan M, McGarry T, et al., 2016. Dysregulated bioenergetics: A key regulator of joint inflammation. *Ann Rheum Dis*, 75(12):2192-2200.  
<https://doi.org/10.1136/annrheumdis-2015-208476>
- Cai W, Zhang J, Yu Y, et al., 2023. Mitochondrial transfer regulates cell fate through metabolic remodeling in osteoporosis. *Adv Sci (Weinh)*, 10(4):e2204871.  
<https://doi.org/10.1002/adv.202204871>
- Callaway DA, Jiang JX, 2015. Reactive oxygen species and oxidative stress in osteoclastogenesis, skeletal aging and bone diseases. *J Bone Miner Metab*, 33(4):359-370.  
<https://doi.org/10.1007/s00774-015-0656-4>
- Chouchani ET, Pell VR, Gaude E, et al., 2014b. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature*, 515(7527):431-435.  
<https://doi.org/10.1038/nature13909>
- Connors J, Dawe N, Van Limbergen J, 2019. The role of succinate in the regulation of intestinal inflammation. *Nutrients*, 11(1)  
<https://doi.org/ARTN2510.3390/nu11010025>
- Dalla Pozza E, Dando I, Pacchiana R, et al., 2020. Regulation of succinate dehydrogenase and role of succinate in cancer. *Semin Cell Dev Biol*, 98:4-14.  
<https://doi.org/10.1016/j.semcdb.2019.04.013>
- De Vadder F, Mithieux G, 2018. Gut-brain signaling in energy homeostasis: The unexpected role of microbiota-derived succinate. *Journal of Endocrinology*, 236(2):R105-R108.  
<https://doi.org/10.1530/Joe-17-0542>
- Deering J, Lin DSY, D'elia A, et al., 2022. Fabrication of succinate-alginate xerogel films for in vitro coupling of osteogenesis and neovascularization. *Biomater Adv*, 141:213122.  
<https://doi.org/10.1016/j.bioadv.2022.213122>
- Deng D, Pan C, Wu Z, et al., 2021. An integrated metabolomic study of osteoporosis: Discovery and quantification of hyocholic acids as candidate markers. *Front Pharmacol*, 12:725341.  
<https://doi.org/10.3389/fphar.2021.725341>
- Farah H, Young SP, Mauro C, et al., 2021. Metabolic dysfunction and inflammatory disease: The role of stromal fibroblasts. *Febs j*, 288(19):5555-5568.  
<https://doi.org/10.1111/febs.15644>
- Farah H, Wijesinghe SN, Nicholson T, et al., 2022. Differential metabolotypes in synovial fibroblasts and synovial fluid in hip osteoarthritis patients support inflammatory responses. *Int J Mol Sci*, 23(6)  
<https://doi.org/10.3390/ijms23063266>
- Faustov LA, Nedel'ko NA, Morozova MV, 2001. [pathomorphology of regenerative processes in mandibular fracture after sodium succinate treatment and laser magnetotherapy in an experimental setting]. *Stomatologiya (Mosk)*, 80(6):8-11.
- Fernández-Veledo S, Vendrell J, 2019. Gut microbiota-derived succinate: Friend or foe in human metabolic diseases? *Rev Endocr Metab Disord*, 20(4):439-447.  
<https://doi.org/10.1007/s11154-019-09513-z>
- Gao B, Deng R, Chai Y, et al., 2019. Macrophage-lineage trap+ cells recruit periosteum-derived cells for periosteal osteogenesis and regeneration. *J Clin Invest*, 129(6):2578-2594.  
<https://doi.org/10.1172/jci98857>
- Gao YD, Zhao Y, Huang J, 2014. Metabolic modeling of common escherichia coli strains in human gut microbiome. *Biomed Res Int*, 2014:694967.  
<https://doi.org/10.1155/2014/694967>
- García-Prat L, Sousa-Victor P, Muñoz-Cánoves P, 2017. Proteostatic and metabolic control of stemness. *Cell Stem Cell*, 20(5):593-608.  
<https://doi.org/10.1016/j.stem.2017.04.011>
- Gilissen J, Jouret F, Pirote B, et al., 2016. Insight into sucnr1 (gpr91) structure and function. *Pharmacol Ther*, 159:56-65.  
<https://doi.org/10.1016/j.pharmthera.2016.01.008>
- Guo Y, Xie C, Li X, et al., 2017. Succinate and its G-protein-coupled receptor stimulates osteoclastogenesis. *Nat Commun*, 8:15621.  
<https://doi.org/10.1038/ncomms15621>



- Guo Y, Chi X, Wang Y, et al., 2020a. Mitochondria transfer enhances proliferation, migration, and osteogenic differentiation of bone marrow mesenchymal stem cell and promotes bone defect healing. *Stem Cell Res Ther*, 11(1):245. <https://doi.org/10.1186/s13287-020-01704-9>
- Guo Y, Cho SW, Saxena D, et al., 2020b. Multifaceted actions of succinate as a signaling transmitter vary with its cellular locations. *Endocrinol Metab (Seoul)*, 35(1):36-43. <https://doi.org/10.3803/EnM.2020.35.1.36>
- Guo Y, Xu F, Thomas SC, et al., 2022. Targeting the succinate receptor effectively inhibits periodontitis. *Cell Rep*, 40(12):111389. <https://doi.org/10.1016/j.celrep.2022.111389>
- Hakak Y, Lehmann-Bruinsma K, Phillips S, et al., 2009. The role of the gpr91 ligand succinate in hematopoiesis. *J Leukoc Biol*, 85(5):837-843. <https://doi.org/10.1189/jlb.1008618>
- He W, Miao FJ, Lin DC, et al., 2004. Citric acid cycle intermediates as ligands for orphan g-protein-coupled receptors. *Nature*, 429(6988):188-193. <https://doi.org/10.1038/nature02488>
- Hogberg C, Gidlöf O, Tan C, et al., 2011. Succinate independently stimulates full platelet activation via camp and phosphoinositide 3-kinase-beta signaling. *J Thromb Haemost*, 9(2):361-372. <https://doi.org/10.1111/j.1538-7836.2010.04158.x>
- Hohl C, Oestreich R, Rosen P, et al., 1987. Evidence for succinate production by reduction of fumarate during hypoxia in isolated adult rat heart cells. *Arch Biochem Biophys*, 259(2):527-535. [https://doi.org/10.1016/0003-9861\(87\)90519-4](https://doi.org/10.1016/0003-9861(87)90519-4)
- Hu Z, Li Y, Zhang L, et al., 2024. Metabolic changes in fibroblast-like synoviocytes in rheumatoid arthritis: State of the art review. *Front Immunol*, 15:1250884. <https://doi.org/10.3389/fimmu.2024.1250884>
- Huang Z, He Z, Kong Y, et al., 2020. Insight into osteoarthritis through integrative analysis of metabolomics and transcriptomics. *Clin Chim Acta*, 510:323-329. <https://doi.org/10.1016/j.cca.2020.07.010>
- Jiang S, Yan W, 2017. Succinate in the cancer-immune cycle. *Cancer Lett*, 390:45-47. <https://doi.org/10.1016/j.canlet.2017.01.019>
- Jin W, Jin Y, Duan P, et al., 2022. Promotion of collagen mineralization and dentin repair by succinates. *J Mater Chem B*, 10(30):5826-5834. <https://doi.org/10.1039/d2tb01005d>
- Jones SW, Brockbank SM, Clements KM, et al., 2009. Mitogen-activated protein kinase-activated protein kinase 2 (mk2) modulates key biological pathways associated with oa disease pathology. *Osteoarthritis Cartilage*, 17(1):124-131. <https://doi.org/10.1016/j.joca.2008.05.001>
- Kaufhold M, Schulz K, Breljak D, et al., 2011. Differential interaction of dicarboxylates with human sodium-dicarboxylate cotransporter 3 and organic anion transporters 1 and 3. *American Journal of Physiology-Renal Physiology*, 301(5):F1026-F1034. <https://doi.org/10.1152/ajprenal.00169.2011>
- Keiran N, Ceperuelo-Mallafre V, Calvo E, et al., 2019. Sucnr1 controls an anti-inflammatory program in macrophages to regulate the metabolic response to obesity. *Nat Immunol*, 20(5):581-592. <https://doi.org/10.1038/s41590-019-0372-7>
- Kim S, Hwang J, Xuan J, et al., 2014. Global metabolite profiling of synovial fluid for the specific diagnosis of rheumatoid arthritis from other inflammatory arthritis. *PLoS One*, 9(6):e97501. <https://doi.org/10.1371/journal.pone.0097501>
- Kin LX, Butler CA, Slakeski N, et al., 2020. Metabolic cooperativity between porphyromonas gingivalis and treponema denticola. *Journal of Oral Microbiology*, 12(1) <https://doi.org/10.1080/20002297.2020.1808750>
- Ko SH, Choi GE, Oh JY, et al., 2017. Succinate promotes stem cell migration through the gpr91-dependent regulation of drp1-mediated mitochondrial fission. *Sci Rep*, 7(1):12582. <https://doi.org/10.1038/s41598-017-12692-x>
- Lee KT, Liao HS, Hsieh MH, 2023. Glutamine metabolism, sensing and signaling in plants. *Plant Cell Physiol*, 64(12):1466-1481. <https://doi.org/10.1093/pcp/pcad054>
- Li XIN, 2023. Succinate signaling in periodontitis induced neuroinflammation and dementia. *NIH RePORTER*,
- Li Y, Zheng JY, Liu JQ, et al., 2016. Succinate/nlrp3 inflammasome induces synovial fibroblast activation: Therapeutic effects of clematichinenoside ar on arthritis. *Front Immunol*, 7:532. <https://doi.org/10.3389/fimmu.2016.00532>

- Li Y, Liu Y, Wang C, et al., 2018. Succinate induces synovial angiogenesis in rheumatoid arthritis through metabolic remodeling and hif-1 $\alpha$ /vegf axis. *Free Radic Biol Med*, 126:1-14.  
<https://doi.org/10.1016/j.freeradbiomed.2018.07.009>
- Liao Z, Han X, Wang Y, et al., 2023. Differential metabolites in osteoarthritis: A systematic review and meta-analysis. *Nutrients*, 15(19)  
<https://doi.org/10.3390/nu15194191>
- Littlewood-Evans A, Sarret S, Apfel V, et al., 2016. Gpr91 senses extracellular succinate released from inflammatory macrophages and exacerbates rheumatoid arthritis. *J Exp Med*, 213(9):1655-1662.  
<https://doi.org/10.1084/jem.20160061>
- Liu J, Gao Z, Liu X, 2024. Mitochondrial dysfunction and therapeutic perspectives in osteoporosis. *Front Endocrinol (Lausanne)*, 15:1325317.  
<https://doi.org/10.3389/fendo.2024.1325317>
- Löffler J, Noom A, Ellinghaus A, et al., 2023. A comprehensive molecular profiling approach reveals metabolic alterations that steer bone tissue regeneration. *Commun Biol*, 6(1):327.  
<https://doi.org/10.1038/s42003-023-04652-1>
- Lu R, Meng H, Gao X, et al., 2014. Effect of non-surgical periodontal treatment on short chain fatty acid levels in gingival crevicular fluid of patients with generalized aggressive periodontitis. *J Periodontol Res*, 49(5):574-583.  
<https://doi.org/10.1111/jre.12137>
- Lu RF, Feng XH, Xu L, et al., 2015. [clinical and putative periodontal pathogens' features of different sites with probing depth reduction after non-surgical periodontal treatment of patients with aggressive periodontitis]. *Beijing Da Xue Xue Bao Yi Xue Ban*, 47(1):13-18 (in Chinese)
- Maevsky EI, Peskov AB, Uchitel ML, et al., 2008. A succinate-based composition reverses menopausal symptoms without sex hormone replacement therapy. *Adv Gerontol*, 21(2):298-305.
- Mao H, Yang A, Zhao Y, et al., 2020. Succinate supplement elicited "pseudohypoxia" condition to promote proliferation, migration, and osteogenesis of periodontal ligament cells. *Stem Cells Int*, 2020:2016809.  
<https://doi.org/10.1155/2020/2016809>
- Mu X, Zhao T, Xu C, et al., 2017. Oncometabolite succinate promotes angiogenesis by upregulating vegf expression through gpr91-mediated stat3 and erk activation. *Oncotarget*, 8(8):13174-13185.  
<https://doi.org/10.18632/oncotarget.14485>
- Murphy MP, O'Neill LaJ, 2018. Krebs cycle reimaged: The emerging roles of succinate and itaconate as signal transducers. *Cell*, 174(4):780-784.  
<https://doi.org/10.1016/j.cell.2018.07.030>
- Nahir AM, Vitis N, Silbermann M, 1990. Cellular enzymatic activities in the articular cartilage of osteoarthritic and osteoporotic hip joints of humans: A quantitative cytochemical study. *Aging (Milano)*, 2(4):363-369.  
<https://doi.org/10.1007/BF03323952>
- Nair S, Sobotka KS, Joshi P, et al., 2019. Lipopolysaccharide-induced alteration of mitochondrial morphology induces a metabolic shift in microglia modulating the inflammatory response in vitro and in vivo. *Glia*, 67(6):1047-1061.  
<https://doi.org/10.1002/glia.23587>
- Nanus DE, Wijesinghe SN, Pearson MJ, et al., 2020. Regulation of the inflammatory synovial fibroblast phenotype by metastasis-associated lung adenocarcinoma transcript 1 long noncoding rna in obese patients with osteoarthritis. *Arthritis Rheumatol*, 72(4):609-619.  
<https://doi.org/10.1002/art.41158>
- Nguyen G, Park SY, Do DV, et al., 2022. Gemigliptin alleviates succinate-induced hepatic stellate cell activation by ameliorating mitochondrial dysfunction. *Endocrinol Metab (Seoul)*, 37(6):918-928.  
<https://doi.org/10.3803/EnM.2022.1530>
- Ohwaki K, 1988. [high carboxylic acid level in the gingival crevicular fluid (gcf) of the patients with advanced periodontal disease]. *Nihon Shishubyo Gakkai Kaishi*, 30(4):985-995.  
<https://doi.org/10.2329/perio.30.985>
- Osei YD, Churchich JE, 1995. Screening and sequence determination of a cDNA encoding the human brain 4-aminobutyrate aminotransferase. *Gene*, 155(2):185-187.  
[https://doi.org/10.1016/0378-1119\(94\)00858-P](https://doi.org/10.1016/0378-1119(94)00858-P)
- Pajor AM, 2014. Sodium-coupled dicarboxylate and citrate transporters from the slc13 family. *Pflugers Archiv-European Journal of Physiology*, 466(1):119-130.  
<https://doi.org/10.1007/s00424-013-1369-y>
- Palmieri F, 2013. The mitochondrial transporter family slc25: Identification, properties and physiopathology. *Mol Aspects Med*, 34(2-3):465-484.

- <https://doi.org/10.1016/j.mam.2012.05.005>
- Park J, Chen Y, Tishkoff DX, et al., 2013. Sirt5-mediated lysine desuccinylation impacts diverse metabolic pathways. *Mol Cell*, 50(6):919-930.  
<https://doi.org/10.1016/j.molcel.2013.06.001>
- Pell VR, Chouchani ET, Frezza C, et al., 2016. Succinate metabolism: A new therapeutic target for myocardial reperfusion injury. *Cardiovasc Res*, 111(2):134-141.  
<https://doi.org/10.1093/cvr/cvw100>
- Pellerin L, Magistretti PJ, 1994. Glutamate uptake into astrocytes stimulates aerobic glycolysis: A mechanism coupling neuronal activity to glucose utilization. *Proc Natl Acad Sci U S A*, 91(22):10625-10629.  
<https://doi.org/10.1073/pnas.91.22.10625>
- Peng B, Li H, Peng XX, 2015. Functional metabolomics: From biomarker discovery to metabolome reprogramming. *Protein Cell*, 6(9):628-637.  
<https://doi.org/10.1007/s13238-015-0185-x>
- Peruzzotti-Jametti L, Bernstock JD, Vicario N, et al., 2018. Macrophage-derived extracellular succinate licenses neural stem cells to suppress chronic neuroinflammation. *Cell Stem Cell*, 22(3):355-368.e313.  
<https://doi.org/10.1016/j.stem.2018.01.020>
- Pollesello P, De Bernard B, Grandolfo M, et al., 1991. Energy state of chondrocytes assessed by 31p-nmr studies of proosseous cartilage. *Biochem Biophys Res Commun*, 180(1):216-222.  
[https://doi.org/10.1016/s0006-291x\(05\)81279-3](https://doi.org/10.1016/s0006-291x(05)81279-3)
- Prag HA, Gruszczuk AV, Huang MM, et al., 2021. Mechanism of succinate efflux upon reperfusion of the ischaemic heart. *Cardiovascular Research*, 117(4):1188-1201.  
<https://doi.org/10.1093/cvr/cvaa148>
- Reddy A, Bozi LHM, Yaghi OK, et al., 2020. Ph-gated succinate secretion regulates muscle remodeling in response to exercise. *Cell*, 183(1):62-75.e17.  
<https://doi.org/10.1016/j.cell.2020.08.039>
- Rubić-Schneider T, Carballido-Perrig N, Regairaz C, et al., 2017. Gpr91 deficiency exacerbates allergic contact dermatitis while reducing arthritic disease in mice. *Allergy*, 72(3):444-452.  
<https://doi.org/10.1111/all.13005>
- Rubic T, Lametschwandtner G, Jost S, et al., 2008. Triggering the succinate receptor gpr91 on dendritic cells enhances immunity. *Nat Immunol*, 9(11):1261-1269.  
<https://doi.org/10.1038/ni.1657>
- Sadagopan N, Li W, Roberds SL, et al., 2007. Circulating succinate is elevated in rodent models of hypertension and metabolic disease. *Am J Hypertens*, 20(11):1209-1215.  
<https://doi.org/10.1016/j.amjhyper.2007.05.010>
- Schlessinger A, Sun NN, Colas C, et al., 2014. Determinants of substrate and cation transport in the human na<sup>+</sup>/dicarboxylate cotransporter nadc3. *Journal of Biological Chemistry*, 289(24):16998-17008.  
<https://doi.org/10.1074/jbc.M114.554790>
- Schmidt-Bleek K, Schell H, Lienau J, et al., 2014. Initial immune reaction and angiogenesis in bone healing. *J Tissue Eng Regen Med*, 8(2):120-130.  
<https://doi.org/10.1002/term.1505>
- Selak MA, Armour SM, Mackenzie ED, et al., 2005. Succinate links tca cycle dysfunction to oncogenesis by inhibiting hif-alpha prolyl hydroxylase. *Cancer Cell*, 7(1):77-85.  
<https://doi.org/10.1016/j.ccr.2004.11.022>
- Shah HN, Williams RaD, 1987. Catabolism of aspartate and asparagine by bacteroides-intermedius and bacteroides-gingivalis. *Current Microbiology*, 15(6):313-318.  
<https://doi.org/10.1007/bf01577587>
- Shen J, Wang C, Li D, et al., 2017. DNA methyltransferase 3b regulates articular cartilage homeostasis by altering metabolism. *JCI Insight*, 2(12)  
<https://doi.org/10.1172/jci.insight.93612>
- Shen J, Wang C, Ying J, et al., 2019. Inhibition of 4-aminobutyrate aminotransferase protects against injury-induced osteoarthritis in mice. *JCI Insight*, 4(18)  
<https://doi.org/10.1172/jci.insight.128568>
- Su H, Lou Y, Fu Y, et al., 2017. Involvement of the vitamin d receptor in energy metabolism revealed by profiling of lysine succinylome of white adipose tissue. *Sci Rep*, 7(1):14132.  
<https://doi.org/10.1038/s41598-017-14477-8>
- Su W, Shi J, Zhao Y, et al., 2020. Porphyromonas gingivalis triggers inflammatory responses in periodontal ligament cells by

- succinate-succinate dehydrogenase-hif-1 $\alpha$  axis. *Biochem Biophys Res Commun*, 522(1):184-190. <https://doi.org/10.1016/j.bbrc.2019.11.074>
- Szekanecz Z, Besenyei T, Szentpetery A, et al., 2010. Angiogenesis and vasculogenesis in rheumatoid arthritis. *Curr Opin Rheumatol*, 22(3):299-306. <https://doi.org/10.1097/BOR.0b013e328337c95a>
- Takahashi N, Sato T, Yamada T, 2000. Metabolic pathways for cytotoxic end product formation from glutamate- and aspartate-containing peptides by porphyromonas gingivalis. *J Bacteriol*, 182(17):4704-4710. <https://doi.org/10.1128/jb.182.17.4704-4710.2000>
- Takahashi N, Yamada T, 2000. Glucose metabolism by prevotella intermedia and prevotella nigrescens. *Oral Microbiol Immunol*, 15(3):188-195. <https://doi.org/10.1034/j.1399-302x.2000.150307.x>
- Tan Jun W, Hao H, Feng X, 2019. Effect of succinic acid on osteogenic differentiation of mouse mc3t3-e1 osteogenic precursor cells. *Mil Med J S Chin*, 33(10):669-672+703 (in Chinese). <https://doi.org/10.13730/j.issn.1009-2595.2019.10.003>
- Tannahill GM, Curtis AM, Adamik J, et al., 2013. Succinate is an inflammatory signal that induces il-1 $\beta$  through hif-1 $\alpha$ . *Nature*, 496(7444):238-242. <https://doi.org/10.1038/nature11986>
- Thomas SC, Guo Y, Xu F, et al., 2024. A novel sucnr1 inhibitor alleviates dysbiosis through inhibition of host responses without direct interaction with host microbiota. *Mol Oral Microbiol*, 39(2):80-90. <https://doi.org/10.1111/omi.12431>
- Toma I, Kang JJ, Sipos A, et al., 2008. Succinate receptor gpr91 provides a direct link between high glucose levels and renin release in murine and rabbit kidney. *J Clin Invest*, 118(7):2526-2534. <https://doi.org/10.1172/jci33293>
- Tomitsuka E, Kita K, Esumi H, 2010. The nadh-fumarate reductase system, a novel mitochondrial energy metabolism, is a new target for anticancer therapy in tumor microenvironments. *Ann N Y Acad Sci*, 1201:44-49. <https://doi.org/10.1111/j.1749-6632.2010.05620.x>
- Uhlen M, Fagerberg L, Hallstrom BM, et al., 2015. Proteomics. Tissue-based map of the human proteome. *Science*, 347(6220):1260419. <https://doi.org/10.1126/science.1260419>
- Valls-Lacalle L, Barba I, Miro-Casas E, et al., 2016. Succinate dehydrogenase inhibition with malonate during reperfusion reduces infarct size by preventing mitochondrial permeability transition. *Cardiovasc Res*, 109(3):374-384. <https://doi.org/10.1093/cvr/cvv279>
- Van Diepen JA, Robben JH, Hooiveld GJ, et al., 2017. Sucnr1-mediated chemotaxis of macrophages aggravates obesity-induced inflammation and diabetes. *Diabetologia*, 60(7):1304-1313. <https://doi.org/10.1007/s00125-017-4261-z>
- Van Pevenage PM, Birchmier JT, June RK, 2023. Utilizing metabolomics to identify potential biomarkers and perturbed metabolic pathways in osteoarthritis: A systematic review. *Semin Arthritis Rheum*, 59:152163. <https://doi.org/10.1016/j.semarthrit.2023.152163>
- Wang T, Xu YQ, Yuan YX, et al., 2019. Succinate induces skeletal muscle fiber remodeling via sucnr1 signaling. *EMBO Rep*, 20(9): e47892. <https://doi.org/10.15252/embr.201947892>
- Wang Y, Han X, Shi J, et al., 2023. Distinct metabolites in osteopenia and osteoporosis: A systematic review and meta-analysis. *Nutrients*, 15(23) <https://doi.org/10.3390/nu15234895>
- Wen H, Gris D, Lei Y, et al., 2011. Fatty acid-induced nlr3-asc inflammasome activation interferes with insulin signaling. *Nat Immunol*, 12(5):408-415. <https://doi.org/10.1038/ni.2022>
- Wentzel JF, Lewies A, Bronkhorst AJ, et al., 2017. Exposure to high levels of fumarate and succinate leads to apoptotic cytotoxicity and altered global DNA methylation profiles in vitro. *Biochimie*, 135:28-34. <https://doi.org/10.1016/j.biochi.2017.01.004>
- Wu KK, 2023. Extracellular succinate: A physiological messenger and a pathological trigger. *Int J Mol Sci*, 24(13) <https://doi.org/10.3390/ijms241311165>
- Wu W, Zhao S, 2013. Metabolic changes in cancer: Beyond the warburg effect. *Acta Biochim Biophys Sin (Shanghai)*, 45(1):18-26. <https://doi.org/10.1093/abbs/gms104>
- Xie N, Tan Z, Banerjee S, et al., 2015. Glycolytic reprogramming in myofibroblast differentiation and lung fibrosis. *Am J Respir Crit Care Med*, 192(12):1462-1474. <https://doi.org/10.1164/rccm.201504-0780OC>

- Yin XN, Wang J, Cui LF, et al., 2018. Enhanced glycolysis in the process of renal fibrosis aggravated the development of chronic kidney disease. *Eur Rev Med Pharmacol Sci*, 22(13):4243-4251.  
[https://doi.org/10.26355/eurrev\\_201807\\_15419](https://doi.org/10.26355/eurrev_201807_15419)
- Yu L, Qi H, An G, et al., 2019. Association between metabolic profiles in urine and bone mineral density of pre- and postmenopausal chinese women. *Menopause*, 26(1):94-102.  
<https://doi.org/10.1097/gme.0000000000001158>
- Zeng R, Fan X, Yang J, et al., 2023. Sdh mutations, as potential predictor of chemotherapy prognosis in small cell lung cancer patients. *Discov Oncol*, 14(1):89.  
<https://doi.org/10.1007/s12672-023-00685-4>
- Zhang W, Lang R, 2023. Succinate metabolism: A promising therapeutic target for inflammation, ischemia/reperfusion injury and cancer. *Front Cell Dev Biol*, 11:1266973.  
<https://doi.org/10.3389/fcell.2023.1266973>
- Zhao Q, Shen H, Su KJ, et al., 2018. Metabolomic profiles associated with bone mineral density in us caucasian women. *Nutr Metab (Lond)*, 15:57.  
<https://doi.org/10.1186/s12986-018-0296-5>
- Zhunossova A, Sen B, Friedman L, et al., 2015. Tumor microenvironment promotes dicarboxylic acid carrier-mediated transport of succinate to fuel prostate cancer mitochondria. *Am J Cancer Res*, 5(5):1665-1679.

unedited